MAGNESIUM CITRATE- leader magnesium citrate oral solution - grape liquid CARDINAL HEALTH, INC. 063997360

Leader Magnesium Citrate Oral Solution - Grape

Active Ingredient

Magnesium citrate 1.745g per fl oz

Purpose

Saline Laxtive

Use

for relief of occasional constipation (irregularity). Generally produces bowel movement in 1/2 to 6 hours

Warnings

kidney disease a magnesium restricted diet abdorominal pain, nausea, or vomiting noticed a sudden change in bowel habits that persists over a period of 2 weeks already used a laxative for a period longer than 1 week

ASK A DOCTOR OR PHARMACIST BEFORE USE IF YOU ARE

taking any other drug. Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.

STOP USE AND ASK A DOCTOR IF

you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition.

IF PREGNANT OR BREAST-FEEDING

ask health professional before use.

KEEP OUT OF REACH OF CHILDREN.

In case of overdose, get medical help or contact a Poison Control Center right away.

Other Information

Store at temperatures between 46° and 86°F (8° and 30° C)

Directions

shake well before using drink a full glass (8 ounces) of liquid with each dose may be taken as a single daily dose or in divided doses

adults and children 12 years of age and over - 6.5 to 10 fl oz maximum 10 fl oz in 24 hours

children 6 to under 12 years of age - 3 to 7 fl oz maximum 7 fl oz in 24 hours

children 2 to under 6 years of age - 2 to 3 fl oz in 24 hours maximum 3 fl oz in 24 hours

children under 2 years of age - ask a doctor

Other Information

each fl oz contains: magnesium 290 mg each fl oz contains: sodium 1 mg

Inactive Ingredients

benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water

Questions?

1-800-200-6313

_	Drug Facts	Drug Facts (continued)	
LEADER	Active ingredient (in each fl oz) Purpose Magnesium citrate 1.745 g	Store at temperatures between 46° and 86°F (8° and 3 °C)	
NDC 70000-0659-1	Uses: For relief of occasional constipation (irregularity). Generally produces bowel movement in ½ to 6 hours	Directions: shake well before using drin full glass (8 ounces) of liquid with each dose mi	
Sugar-Free	Warnings.	adults and children 12 6.5 to 10 fl oz maximu	
Magnesium	Ask a doctor before use if you have ■ Kidney disease ■ a magnesium restricted diet ■ Addominal pain, nausea, or vomiting ■ Noticed a sudden change in bowel habits that persists over a period of 2 weeks	children 6 to under 12 years of age 7 fl oz maximum 7 fl oz in 24 hours	
Citrate	 Noticed a sudgen change in bowel nabits that persists over a period of 2 weeks Already used a laxative for a period longer than 1 week 		
Saline Laxative	Ask a doctor or pharmacist before use if you are Taking any other drug. Take this product 2 or more hours before or after othe drugs. Laxatives may affect how other drugs work.	of age ask a doctor ask a doctor ask a doctor	
Crono Color		each fl oz contains: sodium 1 mg	
Grape Flavor	Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition. If prepnant or breast facting ask a health	Inactive ingredients: benzoic acid, citric acid, disodium EDTA, flavor, sucralose, wate	
Alcohol Content 0.0125% 100% Money Back Guarantee	If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	Questions? 1-833-551-0932 DIST. BY CAH DUBLIN, OH 43017 ©2023 Cardinal He	
(From Flavoring)	Do not use if tamper evident twist-off cap is missing, broken, or separated from neckring	www.myleader.com 1-800-200-6313	
10 FL OZ (296 mL)			
		o "96295"14311	

MAGNESIU			1		
eader magnesi	um citrate ora	l solution - grape liquid	1		
Product Info	rmation				
Product Type		HUMAN OTC DRUG	Item Code (Source)		NDC:70000-0659
Route of Admir	istration	ORAL			
Active Ingred	lient/Active	Moiety			
Ingredient Name Basis of Strength				Strength	
MAGNESIUM CITI UNII:T6V3LHY838)	RATE (UNII: RHO	2601T9V) (MAGNESIUM CA	TION -	MAGNESIUM CITRATE	1.745 g in 29.6 mL
Inactive Ingr	edients				
		Ingredient Name			Strength
EDETATE DISODIUM ANHYDROUS (UNII: 8NLQ36F6MM)					
BENZOIC ACID (U	INII: 85KN0B0MIN	1)			
WATER (UNII: 059	QF0KO0R)				
CITRIC ACID MOI	NOHYDRATE (UN	III: 2968PHW8QP)			
SUCRALOSE (UNII	: 96K6UQ3ZD4)				
Packaging					
# Item Code	Pa	ackage Description		Marketing Start Date	: Marketing End Date

	96 mL in 1 BOTTLE, PLASTIC; Type 0: Not a ombination Product	02/06/2024		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Dru	g M007	02/06/2024		

Labeler - CARDINAL HEALTH, INC. 063997360 (063997360)

Registrant - Pharma Nobis, LLC (118856114)

Establishment

Name	Address	ID/FEI	Business Operations
Pharma Nobis, LLC		118564114	label(70000-0659) , manufacture(70000-0659) , analysis(70000-0659) , pack(70000-0659)

Revised: 2/2024

CARDINAL HEALTH, INC. 063997360