

**ISOPROPYL ALCOHOL- isopropyl alcohol spray spray**  
**Dynarex Corporation**

-----  
**1166 Isopropyl Alcohol Spray**

***Active Ingredient***

Isopropyl Alcohol 70%

***Purpose***

First Aid Antiseptic

***Use(s)***

- First aid to help prevent infection in minor cuts, scrapes, and burns.

***Warnings***

**For External Use Only**

***Do not use***

- Near eyes or mucous membranes • Over large areas of the body • Longer than 1 week unless directed by a doctor • With electrocautery procedures

***Ask a doctor before use if you have***

Deep or puncture wounds, animal bites, or serious burns.

***Stop use and ask a doctor if***

Condition persists for more than 72 hours or get worse.

***Keep out of reach of children***

If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

***Directions***

- Clean the affected area and spray 1 to 3 times daily • May be covered with a sterile bandage • If bandaged, let dry first
- **Children under 12 years of age:** Consult a doctor before use

***Other Information***

- Store at 20°-25°C
- Avoid excessive heat

### Inactive Ingredient

Purified Water

### Questions?

1-888-Dynarex Monday - Friday, 9AM - 5PM EST

### Label 1166



**LOT** LLLLLLLLLL

**YYYY-MM-DD**

**YYYY-MM-DD**



**Manufactured for:**  
 Dynarex Corporation  
 11 Dynarex Drive  
 Middletown, NY 10941  
 USA • www.dynarex.com

Symbol Glossary:  
[dynarex.com/symbols.php](http://dynarex.com/symbols.php)

Made in China  
 R240201  
 NDC# 67777-011-02



## Isopropyl Alcohol Spray

**2 FL. OZ.**  
**Item 1166**

**Drug Facts**

Active Ingredient	Purpose
Isopropyl Alcohol USP 70% .....	First Aid Antiseptic

**Use(s)** ■ First aid to help prevent infection in minor cuts, scrapes, and burns.

**Warnings** For External Use Only

Flammable. Keep away from fire or flame.

**Do not use** ■ Near eyes or mucous membranes  
 ■ Over large areas of the body  
 ■ Longer than 1 week unless directed by a doctor  
 ■ With electrocautery procedures

Ask a doctor before use if you have deep or puncture wounds, animal bites, or serious burns.

Stop use and ask a doctor if condition persists for more than 72 hours or gets worse.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**

■ Clean the affected area and spray 1 to 3 times daily ■ May be covered with a sterile bandage  
 ■ If bandaged, let dry first ■ Children under 12 years of age: Consult a doctor before use

**Other Information**

Store at 20°-25°C ■ Avoid excessive heat

**Inactive Ingredient** Purified Water

**Questions?**  
 1-888-Dynarex Monday-Friday, 9AM – 5PM EST

### Isopropyl Alcohol Spray

<b>ISOPROPYL ALCOHOL</b> isopropyl alcohol spray spray			
Product Information			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:67777-011
<b>Route of Administration</b>	TOPICAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>ISOPROPYL ALCOHOL</b> (UNII: ND2M416302) (ISOPROPYL ALCOHOL - UNII:ND2M416302)	ISOPROPYL ALCOHOL	0.7 mL in 1 mL

**Inactive Ingredients**

Ingredient Name	Strength
<b>WATER</b> (UNII: 059QF0KO0R)	

**Product Characteristics**

<b>Color</b>		<b>Score</b>	
<b>Shape</b>	FREEFORM	<b>Size</b>	
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:67777-011-01	24 in 1 CASE	02/29/2024	
1	NDC:67777-011-02	59.1 mL in 1 BOTTLE, SPRAY; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M003	02/29/2024	

**Labeler** - Dynarex Corporation (008124539)

Revised: 2/2024

Dynarex Corporation