

EXCEDRIN EXTRA STRENGTH GELTABS- acetaminophen, aspirin, caffeine tablet, coated
GlaxoSmithKline Consumer Healthcare Holdings (US) LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredients (in each GelTab)

Acetaminophen 250 mg

Aspirin 250 mg (NSAID)*

Caffeine 65 mg

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever

Pain reliever

Pain reliever aid

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - a cold
 - arthritis
 - muscular aches
 - toothache
 - premenstrual & menstrual cramps

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 gels in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

Do not use

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if

- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are

taking

- a prescription drug for diabetes, gout, or arthritis
- any other drug, or are under a doctor's care for any serious condition

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
 - you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - ringing in the ears or loss of hearing occurs
 - painful area is red or swollen
 - pain gets worse or lasts for more than 10 days
 - fever gets worse or lasts for more than 3 days
 - any new symptoms appear
- These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not use more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 gels every 6 hours; not more than 8 gels in 24 hours
- children under 12 years: ask a doctor

Other information

- store at 20°-25°C (68°-77°F)
- close cap tightly after use
- read all product information before using. Keep this box for important information.

Inactive ingredients

benzoic acid, D&C yellow #10 aluminum lake, edetate disodium, FD&C blue #1, FD&C blue #1 aluminum lake, FD&C red #40 aluminum lake, gelatin, glycerin, hypromellose, iron oxide black, isopropyl alcohol, lactose monohydrate, light mineral oil, low-substituted hydroxypropyl cellulose, maltitol syrup, medium-chain triglycerides, microcrystalline cellulose, pepsin, polysorbate 20, povidone, propyl gallate, propylene glycol, simethicone, sorbitan monolaurate, stearic acid, titanium dioxide

Questions?

1-800-452-0051

Principal Display Panel

NDC 0067-2021-20

EXCEDRIN

EXTRA STRENGTH

Acetaminophen, Aspirin (NSAID) and Caffeine

Pain Reliever/Pain Reliever Aid

20 GELTABS

TAMPER-EVIDENT BOTTLE

DO NOT USE IF INNER FOIL SELA IMPRINTED WITH “SEALED for YOUR PROTECTION” IS BROKEN OR MISSING


Distributed by: **GSK Consumer Healthcare**

Warren, NJ 07059

©2015 GSK or its licensor.

Visit us at www.excedrin.com

46172149



EXCEDRIN EXTRA STRENGTH GELTABS

acetaminophen, aspirin, caffeine tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0067-2021
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg

Inactive Ingredients

Ingredient Name	Strength
BENZOIC ACID (UNII: 8SKN0B0MIM)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C BLUE NO. 1 (UNII: HBR47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
FERROSO FERRIC OXIDE (UNII: XM0M87F357)	
ISOPROPYL ALCOHOL (UNII: ND2M416302)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
LIGHT MINERAL OIL (UNII: N6K5787QVP)	
LOW-SUBSTITUTED HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 2165RE0K14)	
MALTITOL (UNII: D65DG142WK)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
PEPSIN (UNII: GID333S43J)	
POLYSORBATE 20 (UNII: 7T1F30V5YH)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
PROPYL GALLATE (UNII: 8D4SNN7V92)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SORBITAN MONOLAURATE (UNII: 6W9PS8B71J)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	GREEN (dark green) , WHITE	Score	no score
Shape	ROUND	Size	18mm
Flavor		Imprint Code	E
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0067-2021-20	1 in 1 CARTON	09/13/2011	
1		20 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:0067-2021-24	1 in 1 CARTON	09/13/2011	
2		24 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:0067-2021-80	1 in 1 CARTON	09/13/2011	
3		80 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	09/13/2011	

Labeler - GlaxoSmithKline Consumer Healthcare Holdings (US) LLC (079944263)

Revised: 12/2020

GlaxoSmithKline Consumer Healthcare Holdings (US) LLC