

**NICOTINE- nicotine polacrilex gum, chewing  
Dolgenercorp, Inc. (DOLLAR GENERAL & REXALL)**

-----  
**Nicotine Polacrilex Gum Coated Ice Mint 4 mg ANDA204833 - Equate**

***Drug Facts***

***Active ingredient (in each chewing piece)***

Nicotine Polacrilex 4 mg (nicotine)

***Purpose***

Stop smoking aid

***Use***

- reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

***Warnings***

**If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider.** Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

**Ask a doctor before use if you have**

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

**Ask a doctor or pharmacist before use if you are**

- using a non-nicotine stop smoking drug
- taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

**Stop use and ask a doctor if**

- mouth, teeth, or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness,

diarrhea, weakness, and rapid heartbeat

- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

**Keep out of reach of children and pets.**

Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**

- **if you are under 18 years of age, ask a doctor before use**
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- **if you smoke your first cigarette more than 30 minutes after waking up,** use 2 mg nicotine gum
- **if you smoke your first cigarette within 30 minutes of waking up,** use 4 mg nicotine gum according to the following 12 week schedule:

<b>Weeks 1 to 6</b>	<b>Weeks 7 to 9</b>	<b>Weeks 10 to 12</b>
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

**other information**

- **each piece contains:** calcium 115 mg, sodium 8 mg
- store at 20-25°C (68-77°F)
- protect from light

## ***Inactive ingredients***

acacia, acesulfame potassium, carnauba wax, D&C yellow #10, FD&C blue #2, FD&C red #40, FD&C yellow #6, flavors, gum base, hydroxypropyl cellulose, magnesium oxide, sodium bicarbonate, sodium carbonate, sucralose, talc, titanium dioxide, xylitol.

## ***Questions or comments?***

Call **1-888-309-9030**

## **Principal Display Panel**

Compare to the active ingredient in Nicorette® Gum\*\*

Coated

Nicotine Gum

Nicotine Polacrilex Gum 4 mg (Nicotine)

Stop Smoking Aid

### **FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE WITHIN 30 MINUTES OF WAKING UP**

If you smoke your first cigarette MORE THAN 30 MINUTES after waking up, use Nicotine Polacrilex gum, 2 mg

Sugar Free

Coated Fruit Flavor

### **TO INCREASE YOUR SUCCESS IN QUITTING:**

1. You must be motivate to quit
2. Use Enough - Chew at least 9 pieces of Nicotine Polacrilex Gum per day during the first six weeks
3. Use long enough - Use Nicotine Polacrilex Gum for the full 12 weeks
4. Use with a Support Program as directed in the enclosed User's Guide

To remove the gum, tear off single unit

Peel off backing starting at corner with loose edge

Push gum through foil

- not for sale to those under 18 years of age
- proof of age required
- not for sale in vending machines or from any source where proof of age cannot be verified

**BLISTER PACKAGED FOR YOUR PROTECTION. DO NOT USE IF INDIVIDUAL SEALS ARE OPEN OR TORN.**

\*\*This product is not manufactured or distributed by GlaxoSmithKline Consumer

Healthcare, L.P., distributor of Nicorette Gum®.

DISTRIBUTED BY

OLD EAST MAIN CO.

100 MISSION RIDGE

GOODLETTSVILLE, TN 37072

**Package Label**

**TO INCREASE YOUR SUCCESS IN QUITTING:**

1. You must be motivated to quit.
2. **Use Enough** - Chew at least 9 pieces of Nicotine Polacrilex Gum per day during the first six weeks.
3. **Use Long Enough** - Use Nicotine Polacrilex Gum for the full 12 weeks.
4. **Use With a Support Program** as directed in the enclosed User's Guide.

**Drug Facts (continued)**

**Other information**

- each piece contains: calcium 115 mg, sodium 8 mg
- store at 20-25°C (68 -77°F)
- protect from light

**Inactive ingredients**

acacia, acesulfame potassium, carnauba wax, D&C yellow #10 lake, FD&C blue #2 lake, FD&C red #40, FD&C yellow #6 lake, flavors, gum base, hydroxypropyl cellulose, magnesium oxide, sodium bicarbonate, sodium carbonate, sucralose, talc, titanium dioxide, xylitol

**Questions or comments?**  
Call 1-888-309-9030

\*\*This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare, L.P., distributor of Nicorette® Gum.

**BLISTER PACKAGED FOR YOUR PROTECTION. DO NOT USE IF INDIVIDUAL SEALS ARE OPEN OR TORN.**

**100% Satisfaction Guaranteed!**  
(888) 309-9030

DISTRIBUTED BY  
OLD EAST MAIN CO.  
100 MISSION RIDGE  
GOODLETTSVILLE, TN 37072

Product of Ireland



Compare to the active ingredient of Nicorette® Gum\*\*

**Coated Nicotine Gum**

Nicotine Polacrilex Gum  
4 mg (Nicotine)  
Stop Smoking Aid

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE WITHIN 30 MINUTES OF WAKING UP.

If you smoke your first cigarette **MORE THAN 30 MINUTES** after waking up, use Nicotine Polacrilex Gum, 2 mg

**Sugar Free**

**Coated Fruit Flavor**



**20 Pieces**

Actual Gum Size, 4 mg EACH

**4 mg**

■ not for sale to those under 18 years of age  
■ proof of age required  
■ not for sale in vending machines or from any source where proof of age cannot be verified



Push gum through foil.



Peel off backing at starting point, tear off corner with loose edge.



To remove the gum, tear off single unit.

**Drug Facts**

**Active ingredient (in each chewing piece)** Nicotine polacrilex..... 4 mg (nicotine)  
**Purpose** Stop smoking aid

**Use** ■ reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

**Warnings**

If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

**Ask a doctor before use if you have**

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

**Ask a doctor or pharmacist before use if you are**

- using a non-nicotine stop smoking drug
- taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

**Stop use and ask a doctor if**

- mouth, teeth, or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness, and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

**Keep out of reach of children and pets.** Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

A1663



3 59726 93120 5

PLD-A621A FC008687

# DOLLAR GENERAL HEALTH Nicotine Gum Fruit Flavor

## NICOTINE

nicotine polacrilex gum, chewing

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:55910-918
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>NICOTINE</b> (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R)	NICOTINE	4 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>ACACIA</b> (UNII: 5C5403N26O)	
<b>ACESULFAME POTASSIUM</b> (UNII: 23OV73Q5G9)	
<b>CARNAUBA WAX</b> (UNII: R12CBM0EIZ)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>HYDROXYPROPYL CELLULOSE, UNSPECIFIED</b> (UNII: 9XZ8H6N6OH)	
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>SODIUM BICARBONATE</b> (UNII: 8MDF5V39QO)	
<b>SODIUM CARBONATE</b> (UNII: 45P3261C7T)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>XYLITOL</b> (UNII: VCQ006KQ1E)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>MAGNESIUM OXIDE</b> (UNII: 3A3U0GI71G)	
<b>FD&amp;C BLUE NO. 2</b> (UNII: L06K8R7DQK)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	

### Product Characteristics

<b>Color</b>	yellow (Brown (light tan))	<b>Score</b>	no score
<b>Shape</b>	SQUARE	<b>Size</b>	14mm
<b>Flavor</b>	FRUIT	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55910-918-20	20 in 1 CARTON	05/01/2023	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination		

Product			
<b>Marketing Information</b>			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079038	05/01/2023	

**Labeler** - Dolgencorp, Inc. (DOLLAR GENERAL & REXALL) (068331990)

Revised: 5/2023

Dolgencorp, Inc. (DOLLAR GENERAL & REXALL)