

NEOMED 325- neomycin sulfate powder, for solution
Bimeda, Inc.

100 g pouch

NeoMed® 325

(neomycin sulfate)

Soluble Powder (Commercial Grade)

FOR ORAL USE IN CATTLE, SWINE, SHEEP, GOATS & TURKEYS

For the treatment and control of colibacillosis (bacterial enteritis) caused by *Escherichia coli* susceptible to neomycin sulfate in cattle (excluding veal calves), swine, sheep and goats.

For the control of mortality associated with *Escherichia coli* organisms susceptible to neomycin sulfate in growing turkeys.

Antibacterial

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

Each Packet Contains: 71.4 g neomycin sulfate (commercial grade) equivalent to 50 g neomycin.

NOT FOR USE IN HUMANS

KEEP OUT OF REACH OF CHILDREN

Restricted Drug (California)- Use Only as Directed

Approved by FDA under ANADA # 200-050

NET WEIGHT:

100 g (3.5 oz)

100 g pouch

Add to drinking water - Not for use in liquid supplements.

DOSAGE AND ADMINISTRATION: Administer to cattle (excluding veal calves), swine, sheep and goats at a dose of 10 mg of neomycin sulfate per pound of body weight per day in divided doses for a maximum of 14 days. Administer to turkeys at a dose of 10 mg neomycin sulfate per pound of body weight per day for 5 days.

Herd/Flock Treatment: Each packet will treat 7,150 pounds body weight. Therefore, estimate the total number of pounds of body weight of the animals to be treated and administer one (1) packet (or portion thereof) for each 7,150 pounds. The product should be added to the amount of drinking water estimated to be consumed in 12-24 hours. Provide medicated water as the sole source of water each day until consumed, followed by non-medicated water as required. Fresh medicated water should be prepared each day.

Individual Animal Treatment: To provide 10 mg neomycin sulfate per pound of body weight, mix one (1) level teaspoon in water or milk for each 160 pounds of body weight. Administer daily either as a drench in divided doses or in the drinking water to be

consumed in 12-24 hours.

Drinking Water: Swine Use the number of packets indicated below in 256 gallons of water, or in two gallons of stock solution used in proportioner set to meter one ounce per gallon.

Pigs Weighing 25 to 50 Pounds	2 Packets
Pigs Weighing 50 to 100 Pounds	3 Packets
Pigs Weighing Over 100 Pounds	4 Packets

200 g Pouch

NeoMed® 325

(neomycin sulfate)

Soluble Powder (Commercial Grade)

FOR ORAL USE IN CATTLE, SWINE, SHEEP, GOATS & TURKEYS

For the treatment and control of colibacillosis (bacterial enteritis) caused by *Escherichia coli* susceptible to neomycin sulfate in cattle (excluding veal calves), swine, sheep and goats.

For the control of mortality associated with *Escherichia coli* organisms susceptible to neomycin sulfate in growing turkeys.

Antibacterial

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

Each Packet Contains: 142.8 g neomycin sulfate (commercial grade) equivalent to 100 g neomycin.

NOT FOR USE IN HUMANS

KEEP OUT OF REACH OF CHILDREN

Restricted Drug (California)- Use Only as Directed

Approved by FDA under ANADA # 200-050

NET WEIGHT:

200 g (7 oz)

200 g Pouch

Add to drinking water - Not for use in liquid supplements.

DOSAGE AND ADMINISTRATION: Administer to cattle (excluding veal calves), swine, sheep and goats at a dose of 10 mg of neomycin sulfate per pound of body weight per day in divided doses for a maximum of 14 days. Administer to turkeys at a dose of 10 mg neomycin sulfate per pound of body weight per day for 5 days.

Herd/Flock Treatment: Each packet will treat 14,300 pounds body weight. Therefore, estimate the total number of pounds of body weight of the animals to be treated and

administer one (1) packet (or portion thereof) for each 14,300 pounds. The product should be added to the amount of drinking water estimated to be consumed in 12-24 hours. Provide medicated water as the sole source of water each day until consumed, followed by non-medicated water as required. Fresh medicated water should be prepared each day.

Individual Animal Treatment: To provide 10 mg neomycin sulfate per pound of body weight, mix one (1) level teaspoon in water or milk for each 160 pounds of body weight. Administer daily either as a drench in divided doses or in the drinking water to be consumed in 12-24 hours.

Drinking Water: Swine Use the number of packets indicated below in 256 gallons of water, or in two gallons of stock solution used in proportioner set to meter one ounce per gallon.

Pigs Weighing 25-50 Pounds	1 Packet
Pigs Weighing 50 to 100 Pounds	1.5 Packets
Pigs Weighing Over 100 Pounds	2 Packets

50 Pounds

NeoMed® 325

(neomycin sulfate)

Soluble Powder (Commercial Grade)

FOR ORAL USE IN CATTLE, SWINE, SHEEP, GOATS & TURKEYS

For the treatment and control of colibacillosis (bacterial enteritis) caused by *Escherichia coli* susceptible to neomycin sulfate in cattle (excluding veal calves), swine, sheep and goats.

For the control of mortality associated with *Escherichia coli* organisms susceptible to neomycin sulfate in growing turkeys.

Antibacterial

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

Each Pound Contains: 325 g neomycin sulfate (commercial grade) equivalent to 227.5 g neomycin.

NOT FOR USE IN HUMANS

KEEP OUT OF REACH OF CHILDREN

Restricted Drug (California)- Use Only as Directed

Approved by FDA under ANADA # 200-050

NET WEIGHT:

50 Pounds (22.7 kg)

50 Pounds

Add to drinking water - Not for use in liquid supplements.

Dosage and Administration: Administer to cattle (excluding veal calves), swine, sheep and goats at a dose of 10 mg of neomycin sulfate per pound of body weight per day in divided doses for a maximum of 14 days. Administer to turkeys at a dose of 10 mg neomycin sulfate per pound of body weight per day for 5 days. Using measuring spoons, the recommended dosage can be determined as follows:

Daily Schedule For Drinking Water:

SWINE

One level tablespoon of NeoMed 325 added to water consumed in one day will treat	Weight of each pig
50 pigs	10 pounds
20 pigs	25 pounds
10 pigs	50 pounds

CATTLE

One level tablespoon of NeoMed 325 added to water consumed in one day will treat	Weight of each calf
10 calves	50 pounds
7 calves	75 pounds
4 calves	125 pounds

TURKEYS

One level tablespoon of NeoMed 325 added to water consumed in one day will treat	Weight of each bird
1000 birds	0.5 pounds
500 birds	1 pounds
100 birds	5 pounds
50 birds	10 pounds

The product should be added to the amount of drinking water estimated to be consumed in 12-24 hours. Provide medicated water as the sole source of water each day until consumed, followed by non-medicated water as required. Fresh medicated water should be prepared each day. For a **stock solution**, add six level tablespoons to one gallon of water. Each pint of this stock solution will medicate 5 gallons of drinking water.

For use in **Automatic Proportioners** delivering 2 ounces of stock solution per gallon of drinking water, dissolve 9 level tablespoons in a gallon of water to make the stock solution.

Individual Animal Treatment: To provide 10 mg neomycin sulfate per pound of body weight, mix one (1) level teaspoon in water or milk for each 160 pounds of body weight. Administer daily either as a drench in divided doses or in the drinking water to be consumed in 12-24 hours.

Caution: To administer the stated dosage, the concentration of neomycin required in medicated water must be adjusted to compensate for variation in age and weight of animal, the nature and severity of disease signs, and environmental temperature and humidity, each of which affects water consumption.

If symptoms persist after using this preparation for 2 or 3 days, consult a veterinarian. If symptoms such as fever, depression, or going off feed develop, oral neomycin is not indicated as the sole treatment since systemic levels of neomycin are not obtained due to low absorption from the gastrointestinal tract.

Important: Treatment should continue 24 to 48 hours beyond remission of disease symptoms, but not to exceed a total of 14 consecutive days for cattle, swine, sheep and goats and 5 days for turkeys. Animals not drinking or eating should be treated individually by drench.

Warning: Not for human use. Keep out of reach of children.

Residue Warnings: Discontinue treatment prior to slaughter by at least the number of days listed below for appropriate species:

Cattle	1 day
Swine and Goats	3 days
Sheep	2 days
Turkeys	0 days

A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal. A milk discard period has not been established for this product in lactating dairy cattle. Do not use in female dairy cattle 20 months of age or older. Use of more than one product containing neomycin or failure to follow withdrawal times may result in illegal drug residues.

The Safety Data Sheet (SDS) contains more detailed occupational safety information. To report suspected adverse drug events, for technical assistance or to obtain a copy of the SDS, contact Bimeda, Inc. at 1-888-524-6332. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or online at www.fda.gov/reportanimalae.

100 g and 200 g Pouches


When storing partially used packet, securely close protect from moisture. Store in a dry place.

Store at controlled room temperature 20°C-25°C (68°F-77°F), with excursions permitted to 15°C- 30°C (59°-86°F).

50 Pounds

When storing partially used containers, securely close bags and keep container tightly closed to prevent contents from caking. Store in a dry place.

Store at controlled room temperature 20°C-25°C (68°F-77°F), with excursions permitted to 15°C- 30°C (59°-86°F).



NeoMed[®] 325

(neomycin sulfate)
Soluble Powder (Commercial Grade)

FOR ORAL USE IN CATTLE, SWINE, SHEEP, GOATS & TURKEYS

For the treatment and control of colibacillosis (bacterial enteritis) caused by *Escherichia coli* susceptible to neomycin sulfate in cattle (excluding veal calves), swine, sheep and goats.

For the control of mortality associated with *Escherichia coli* organisms susceptible to neomycin sulfate in growing turkeys.

Antibacterial

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

EACH PACKET CONTAINS: 71.4 g neomycin sulfate (commercial grade) equivalent to 50 g neomycin.

NOT FOR USE IN HUMANS

KEEP OUT OF REACH OF CHILDREN

Restricted Drug (California) - Use Only as Directed

Approved by FDA under ANADA # 200-050



**Net
Weight: 100 g (3.5 oz)**

Manufactured By:
Teva Inc.
Lakewood, CO 80015
WWW.TEVAINC.COM

NeoMed[®] 325 (neomycin sulfate) **Soluble Powder (Commercial Grade)**

Add to drinking water - Not for use in liquid supplements.

Dosage and Administration: Administer to cattle (excluding weal calves), swine, sheep and goats at a dose of 10 mg of neomycin sulfate per pound of body weight per day in divided doses for a maximum of 14 days. Administer to turkeys at a dose of 10 mg neomycin sulfate per pound of body weight per day for 5 days.

Herb/Pack Treatment: Each packet will treat 7,150 pounds body weight. Therefore, estimate the total number of pounds of body weight of the animals to be treated and administer one (1) packet for portion thereof for each 7,150 pounds. The product should be added to the amount of drinking water estimated to be consumed in 12-24 hours. Provide medicated water as the sole source of water each day until consumed, followed by non-medicated water as required. Fresh indicated water should be prepared each day.

Individual Animal Treatment: To provide 10 mg neomycin sulfate per pound of body weight, mix one (1) level teaspoon in water or milk for each 100 pounds body weight. Administer daily either as a drench in divided doses or in the drinking water to be consumed in 12-24 hours.

Drinking Water: Swine - Use the number of packets indicated below in 256 gallons of water, or in two gallons of stock solution used in proportioner set to meter one ounce per gallon.

Pigs Weighing 25 to 50 Pounds 2 Packets
Pigs Weighing 50 to 100 Pounds 3 Packets
Pigs Weighing Over 100 Pounds 4 Packets

Caution: To administer the stated dosage, the concentration of neomycin required in medicated water must be adjusted to compensate for variation in age and weight of animal, the nature and severity of disease signs, and environmental temperature and humidity, each of which affects water consumption.

If symptoms persist after using this preparation for 2 or 3 days, consult a veterinarian. If symptoms such as fever, depression, or going off feed develop, oral neomycin is not indicated as the sole treatment since systemic levels of neomycin are not obtained due to low absorption from the gastrointestinal tract.

Important: Treatment should continue 24 to 48 hours beyond remission of disease symptoms, but not to exceed a total of 14 consecutive days for cattle, swine, sheep and goats and 5 days for turkeys. Animals not drinking or eating should be treated individually by drench.

Warning: Not for human use. Keep out of reach of children.

Residue Warnings: Discontinue treatment prior to slaughter by at least the number of days listed below for the appropriate species:


Cattle 1 day Sheep 2 days
Swine and goats 3 days Turkeys 0 days

A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal. A milk discard period has not been established for this product in lactating dairy cattle. Do not use in female dairy cattle 20 months of age or older. Use of more than one product containing neomycin or failure to follow withdrawal times may result in illegal drug residues.


The Safety Data Sheet (SDS) contains more detailed occupational safety information. To report suspected adverse drug events, for technical assistance or to obtain a copy of the SDS, contact Bimeda, Inc. at 1-888-524-6332. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or online at www.fda.gov/medwatch.

When storing partially used packet, securely close to protect from moisture. Store in a dry place.

Store at controlled room temperature 20°C-25°C (68°F-77°F), with excursions permitted to 15°C-30°C (59°F-86°F).



NeoMed[®] is a registered trademark of Bimeda, Inc.
NEOMC 942015 Rev 1/21



NeoMed[®] 325

(neomycin sulfate)
Soluble Powder (Commercial Grade)

FOR ORAL USE IN CATTLE, SWINE, SHEEP, GOATS & TURKEYS

For the treatment and control of colibacillosis (bacterial enteritis) caused by *Escherichia coli* susceptible to neomycin sulfate in cattle (excluding veal calves), swine, sheep and goats.

For the control of mortality associated with *Escherichia coli* organisms susceptible to neomycin sulfate in growing turkeys.

Antibacterial

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

EACH PACKET CONTAINS: 142.8 g neomycin sulfate (commercial grade) equivalent to 100 g neomycin.

NOT FOR USE IN HUMANS
KEEP OUT OF REACH OF CHILDREN

Restricted Drug (California) - Use Only as Directed
Approved by FDA under ANADA # 200-050



Net
Weight: 200 g (7 oz)

Manufactured By:
Bimeda, Inc.
Le Sueur, MN 56088
www.bimeda.com

NeoMed[®] 325 (neomycin sulfate) **Soluble Powder (Commercial Grade)**

Add to drinking water - Not for use in liquid supplements.

Dosage and Administration: Administer to cattle (including veal calves), swine, sheep and goats at a dose of 10 mg of neomycin sulfate per pound of body weight per day in divided doses for a maximum of 14 days. Administer to turkeys at a dose of 10 mg neomycin sulfate per pound of body weight per day for 5 days.

Herd/Flock Treatment: Each packet will treat 14,300 pounds body weight. Therefore, estimate the total number of pounds of body weight of the animals to be treated and administer one (1) packet (or portion thereof) for each 14,300 pounds. The product should be added to the amount of drinking water estimated to be consumed in 12-24 hours. Provide medicated water as the sole source of water each day until consumed, followed by non-medicated water as required. Fresh medicated water should be prepared each day.

Individual Animal Treatment: To provide 10 mg neomycin sulfate per pound of body weight, mix one (1) level teaspoon in water or milk for each 100 pounds body weight. Administer daily either as a drench in divided doses or in the drinking water to be consumed in 12-24 hours.

Drinking Water: Swine - Use the number of packets indicated below in 256 gallons of water, or in two gallons of stock solution used in proportioner and to water one cubic per gallon.

Pigs Weighing 25 to 50 Pounds	1 Packet
Pigs Weighing 50 to 100 Pounds	1.5 Packets
Pigs Weighing Over 100 Pounds	2 Packets

Caution: To administer the stated dosage, the concentration of neomycin required in medicated water must be adjusted to compensate for variation in age and weight of animal, the nature and severity of disease signs, and environmental temperature and humidity, each of which affects water consumption.

If symptoms persist after using this preparation for 2 or 3 days, consult a veterinarian. If symptoms such as fever, depression, or going off feed develop, oral neomycin is not indicated as the sole treatment since systemic levels of neomycin are not obtained due to low absorption from the gastrointestinal tract.

Important: Treatment should continue 24 to 48 hours beyond remission of disease symptoms, but not to exceed a total of 14 consecutive days for cattle, swine, sheep and goats and 5 days for turkeys. Animals not drinking or eating should be treated individually by drench.

Warning: Not for human use. Keep out of reach of children.

Residue Warnings: Discontinue treatment prior to slaughter by at least the number of days listed below for the appropriate species.			
Cattle	1 day	Sheep	2 days
Swine and goats	3 days	Turkeys	0 days

A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal. A milk discard period has not been established for this product in lactating dairy cattle. Do not use in female dairy cattle 20 months of age or older. Use of more than one product containing neomycin or failure to follow withdrawal times may result in illegal drug residues.

The Safety Data Sheet (SDS) contains more detailed occupational safety information. To report suspected adverse drug events, for technical assistance or to obtain a copy of the SDS, contact Bimeda, Inc. at 1-888-524-6232. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VEIS or online at www.fda.gov/veis/index.html.

When storing partially used packet, securely close to protect from moisture. Store in a dry place.
Store at controlled room temperature 20°C-25°C (68°F-77°F), with excursions permitted to 15°C-30°C (59°F-86°F).



NeoMed[®] is a registered trademark of Bimeda, Inc.
V02010 080008 Rev. 12/11

Add to drinking water - Not for use in liquid supplements.

Dosage and Administration: Administer to cattle (excluding veal calves), swine, sheep and goats at a dose of 10 mg of neomycin sulfate per pound of body weight per day in divided doses for a maximum of 14 days. Administer to turkeys at a dose of 10 mg neomycin sulfate per pound of body weight per day for 5 days. Using measuring spoons, the recommended dosage can be determined as follows:

Daily Schedule For Drinking Water:

SWINE: NeoMed 325	
One level tablespoon of NeoMed 325 added to water consumed in one day will treat	Weight of each pig
50 pigs	10 pounds
20 pigs	25 pounds
10 pigs	50 pounds
CATTLE: NeoMed 325	
One level tablespoon of NeoMed 325 added to water consumed in one day will treat	Weight of each calf
10 calves	50 pounds
7 calves	75 pounds
4 calves	125 pounds
TURKEYS: NeoMed 325	
One level tablespoon of NeoMed 325 added to water consumed in one day will treat	Weight of each bird
1000 birds	0.5 pounds
500 birds	1 pound
100 birds	5 pounds
50 birds	10 pounds

The product should be added to the amount of drinking water estimated to be consumed in 12-24 hours. Provide medicated water as the sole source of water each day until consumed, followed by non-medicated water as required. Fresh medicated water should be prepared each day. For a **stock solution**, add six level tablespoons to one gallon of water. Each pint of this stock solution will medicate 5 gallons of drinking water.

For use in **Automatic Proportioners** delivering 2 ounces of stock solution per gallon of drinking water, dissolve 9 level tablespoons in a gallon of water to make the stock solution.



NeoMed® 325

(neomycin sulfate)
Soluble Powder (Commercial Grade)

FOR ORAL USE IN CATTLE, SWINE, SHEEP, GOATS & TURKEYS

For the treatment and control of colibacillosis (bacterial enteritis) caused by *Escherichia coli* susceptible to neomycin sulfate in cattle (excluding veal calves), swine, sheep and goats.

For the control of mortality associated with *Escherichia coli* organisms susceptible to neomycin sulfate in growing turkeys.


Antibacterial

CAUTION: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

Each Pound Contains: 325 g neomycin sulfate (commercial grade) equivalent to 227.5 g neomycin.

NOT FOR USE IN HUMANS
KEEP OUT OF REACH OF CHILDREN

Restricted Drug (California) - Use Only as Directed
Approved by FDA under ANADA # 200-050



Net Weight:
50 Pounds (22.7 kg)

Manufactured By:
Bimeda, Inc.
Le Sueur, MN 56058
www.bimeda.com

Individual Animal Treatment: To provide 10 mg neomycin sulfate per pound of body weight, mix one (1) level teaspoon in water or milk for each 160 pounds of body weight. Administer daily either as a drench in divided doses or in the drinking water to be consumed in 12-24 hours.

Caution: To administer the stated dosage, the concentration of neomycin required in medicated water must be adjusted to compensate for variation in age and weight of animal, the nature and severity of disease signs, and environmental temperature and humidity, each of which affects water consumption.

If symptoms persist after using this preparation for 2 or 3 days, consult a veterinarian. If symptoms such as fever, depression, or going off feed develop, oral neomycin is not indicated as the sole treatment since systemic levels of neomycin are not obtained due to low absorption from the gastrointestinal tract.

Important: Treatment should continue 24 to 48 hours beyond remission of disease symptoms, but not to exceed a total of 14 consecutive days for cattle, swine, sheep and goats and 5 days for turkeys. Animals not drinking or eating should be treated individually by drench.

Warning: Not for human use. Keep out of reach of children.

Residue Warnings: Discontinue treatment prior to slaughter by at least the number of days listed below for the appropriate species:

Cattle.....	1 day
Swine and goats.....	3 days
Sheep.....	2 days
Turkeys.....	0 days

A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

A milk discard period has not been established for this product in lactating dairy cattle. Do not use in female dairy cattle 20 months of age or older.

Use of more than one product containing neomycin or failure to follow withdrawal times may result in illegal drug residues.

The Safety Data Sheet (SDS) contains more detailed occupational safety information. To report suspected adverse drug events, for technical assistance or to obtain a copy of the SDS, contact Bimeda, Inc. at 1-888-524-6332. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or online at www.fda.gov/reportanimal.

When storing partially used containers, securely close bags and keep container tightly closed to prevent contents from caking. Store in a dry place.

Store at controlled room temperature 20°C-25°C (68°F-77°F), with excursions permitted to 15°C-30°C (59°F-86°F).

NeoMed® is a registered trademark of Bimeda, Inc.



1NE007 BNE003 Rev. 12/21

LOT:
EXP:

NEOMED 325

neomycin sulfate powder, for solution

Product Information

Product Type	PRESCRIPTION ANIMAL DRUG	Item Code (Source)	NDC:61133-5442
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NEOMYCIN SULFATE (UNII: 057Y626693) (Neomycin - UNII:I16QD7X297)	Neomycin	227.5 g in 0.454 kg

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:61133-5442-2	22.7 kg in 1 PAIL		
2	NDC:61133-5442-3	0.2 kg in 1 POUCH		
3	NDC:61133-5442-4	0.1 kg in 1 POUCH		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANADA	ANADA200050	01/12/2010	

Labeler - Bimeda, Inc. (060492923)

Registrant - Bimeda, Inc. (060492923)

Establishment

Name	Address	ID/FEI	Business Operations
Bimeda, Inc.		060492923	manufacture

Revised: 3/2024

Bimeda, Inc.