

# VASOPRESSIN - vasopressin injection

## Eagle Pharmaceuticals, Inc.

-----

### HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use VASOPRESSIN INJECTION, safely and effectively. See full prescribing information for VASOPRESSIN INJECTION.

**VASOPRESSIN Injection, for intravenous use**  
**Initial U.S. Approval: 2014**

#### INDICATIONS AND USAGE

- Vasopressin Injection is indicated to increase blood pressure in adults with vasodilatory shock who remain hypotensive despite fluids and catecholamines. (1)

#### DOSAGE AND ADMINISTRATION

- Dilute 20 units/mL multiple dose vial content with normal saline (0.9% sodium chloride) or 5% dextrose in water (D5W) to either 0.1 units per mL or 1 unit per mL for intravenous administration. Discard unused diluted solution after 18 hours at room temperature or 24 hours under refrigeration. (2.1)
- Post-cardiotomy shock: 0.03 to 0.1 units/minute (2.2)
- Septic shock: 0.01 to 0.07 units/minute (2.2)

#### DOSAGE FORMS AND STRENGTHS

- Injection: 20 units per mL in a multiple dose vial. To be used after dilution. (3)

#### CONTRAINDICATIONS

- Vasopressin Injection is contraindicated in patients with known allergy or hypersensitivity to 8-L-arginine vasopressin or chlorobutanol. (4)

#### WARNINGS AND PRECAUTIONS

- Can worsen cardiac function. (5.1)
- Reversible diabetes insipidus. (5.2)

#### ADVERSE REACTIONS

The most common adverse reactions include decreased cardiac output, bradycardia, tachyarrhythmias, hyponatremia and ischemia (coronary, mesenteric, skin, digital). (6)

**To report SUSPECTED ADVERSE REACTIONS, contact Eagle Pharmaceuticals, Inc. at 1-855-318-2170 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch)**

#### DRUG INTERACTIONS

- Pressor effects of catecholamines and Vasopressin Injection are expected to be additive. (7.1)
- Indomethacin may prolong effects of Vasopressin Injection. (7.2)
- Co-administration of ganglionic blockers or drugs causing SIADH may increase the pressor response. (7.3, 7.4)
- Co-administration of drugs causing diabetes insipidus may decrease the pressor response. (7.5)

#### USE IN SPECIFIC POPULATIONS

**Pregnancy:** May induce uterine contractions. (8.1)

**Pediatric Use:** Safety and effectiveness have not been established. (8.4)

**Geriatric Use:** No safety issues have been identified in older patients. (8.5)

**Revised: 6/2021**

---

## FULL PRESCRIBING INFORMATION: CONTENTS\*

### 1 INDICATIONS AND USAGE

### 2 DOSAGE AND ADMINISTRATION

#### 2.1 Preparation of Solution

2.2 Administration

### **3 DOSAGE FORMS AND STRENGTHS**

### **4 CONTRAINDICATIONS**

### **5 WARNINGS AND PRECAUTIONS**

5.1 Worsening Cardiac Function

5.2 Reversible Diabetes Insipidus

### **6 ADVERSE REACTIONS**

### **7 DRUG INTERACTIONS**

7.1 Catecholamines

7.2 Indomethacin

7.3 Ganglionic Blocking Agents

7.4 Drugs Suspected of Causing SIADH

7.5 Drugs Suspected of Causing Diabetes Insipidus

### **8 USE IN SPECIFIC POPULATIONS**

8.1 Pregnancy

8.2 Lactation

8.4 Pediatric Use

8.5 Geriatric Use

### **10 OVERDOSAGE**

### **11 DESCRIPTION**

### **12 CLINICAL PHARMACOLOGY**

12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics

### **13 NONCLINICAL TOXICOLOGY**

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

13.2 Animal Toxicology and/or Pharmacology

### **14 CLINICAL STUDIES**

### **16 HOW SUPPLIED/STORAGE AND HANDLING**

\* Sections or subsections omitted from the full prescribing information are not listed.

---

## **FULL PRESCRIBING INFORMATION**

### **1 INDICATIONS AND USAGE**

Vasopressin Injection is indicated to increase blood pressure in adults with vasodilatory shock who remain hypotensive despite fluids and catecholamines.

### **2 DOSAGE AND ADMINISTRATION**

#### **2.1 Preparation of Solution**

Inspect parenteral drug products for particulate matter and discoloration prior to use, whenever solution and container permit.

#### **Vasopressin Injection Solution for Dilution, 20 units/mL**

Dilute Vasopressin Injection in normal saline (0.9% sodium chloride) or 5% dextrose in water (D5W) prior to use for intravenous administration. Discard unused diluted solution after 18 hours at room temperature or 24 hours under refrigeration.

**Table 1 Preparation of diluted solutions**

Fluid restriction?	Final concentration	Mix	
		Vasopressin Injection	Diluent
No	0.1 units/mL	2.5 mL (50 units)	500 mL
Yes	1 unit/mL	5 mL (100 units)	100 mL

## 2.2 Administration

In general, titrate to the lowest dose compatible with a clinically acceptable response.

The recommended starting dose is:

*Post-cardiotomy shock:* 0.03 units/minute

*Septic Shock:* 0.01 units/minute

Titrate up by 0.005 units/minute at 10- to 15-minute intervals until the target blood pressure is reached. There are limited data for doses above 0.1 units/minute for post-cardiotomy shock and 0.07 units/minute for septic shock. Adverse reactions are expected to increase with higher doses.

After target blood pressure has been maintained for 8 hours without the use of catecholamines, taper vasopressin injection by 0.005 units/minute every hour as tolerated to maintain target blood pressure.

## 3 DOSAGE FORMS AND STRENGTHS

Vasopressin injection, USP is a clear, practically colorless solution available as 20 units/mL in a multiple dose vial. To be used after dilution.

## 4 CONTRAINDICATIONS

Vasopressin Injection is contraindicated in patients with known allergy or hypersensitivity to 8-L-arginine vasopressin or chlorobutanol.

## 5 WARNINGS AND PRECAUTIONS

### 5.1 Worsening Cardiac Function

A decrease in cardiac index may be observed with the use of vasopressin.

### 5.2 Reversible Diabetes Insipidus

Patients may experience reversible diabetes insipidus, manifested by the development of polyuria, a dilute urine, and hypernatremia, after cessation of treatment with vasopressin. Monitor serum electrolytes, fluid status and urine output after vasopressin discontinuation. Some patients may require readministration of vasopressin or administration of desmopressin to correct fluid and electrolyte shifts.

## 6 ADVERSE REACTIONS

The following adverse reactions associated with the use of vasopressin were identified in

the literature. Because these reactions are reported voluntarily from a population of uncertain size, it is not possible to estimate their frequency reliably or to establish a causal relationship to drug exposure.

Bleeding/lymphatic system disorders: Hemorrhagic shock, decreased platelets, intractable bleeding

Cardiac disorders: Right heart failure, atrial fibrillation, bradycardia, myocardial ischemia

Gastrointestinal disorders: Mesenteric ischemia

Hepatobiliary: Increased bilirubin levels

Renal/urinary disorders: Acute renal insufficiency

Vascular disorders: Distal limb ischemia

Metabolic: Hyponatremia

Skin: Ischemic lesions

## **Postmarketing Experience**

Reversible diabetes insipidus [see *Warnings and Precautions (5.2)*].

## **7 DRUG INTERACTIONS**

### **7.1 Catecholamines**

Use with *catecholamines* is expected to result in an additive effect on mean arterial blood pressure and other hemodynamic parameters. Hemodynamic monitoring is recommended; adjust the dose of vasopressin as needed.

### **7.2 Indomethacin**

Use with *indomethacin* may prolong the effect of Vasopressin Injection on cardiac index and systemic vascular resistance. Hemodynamic monitoring is recommended; adjust the dose of vasopressin as needed [see *Clinical Pharmacology (12.3)*].

### **7.3 Ganglionic Blocking Agents**

Use with *ganglionic blocking agents* may increase the effect of Vasopressin Injection on mean arterial blood pressure. Hemodynamic monitoring is recommended; adjust the dose of vasopressin as needed [see *Clinical Pharmacology (12.3)*].

### **7.4 Drugs Suspected of Causing SIADH**

Use with *drugs suspected of causing SIADH* (e.g., SSRIs, tricyclic antidepressants, haloperidol, chlorpropamide, enalapril, methyldopa, pentamidine, vincristine, cyclophosphamide, ifosfamide, felbamate) may increase the pressor effect in addition to the antidiuretic effect of Vasopressin Injection. Hemodynamic monitoring is recommended; adjust the dose of vasopressin as needed.

### **7.5 Drugs Suspected of Causing Diabetes Insipidus**

Use with *drugs suspected of causing diabetes insipidus* (e.g., demeclocycline, lithium, fosfarnet, clozapine) may decrease the pressor effect in addition to the antidiuretic effect of Vasopressin Injection. Hemodynamic monitoring is recommended; adjust the

dose of vasopressin as needed.

## **8 USE IN SPECIFIC POPULATIONS**

### **8.1 Pregnancy**

#### Risk Summary

There are no available data on Vasopressin Injection use in pregnant women to inform a drug associated risk of major birth defects, miscarriage, or adverse maternal or fetal outcomes. Animal reproduction studies have not been conducted.

#### Clinical Considerations

*Dose adjustments during pregnancy and the postpartum period:* Because of increased clearance of vasopressin in the second and third trimester, the dose of Vasopressin Injection may need to be increased [see *Dosage and Administration (2.2)* and see *Clinical Pharmacology (12.3)*].

*Maternal adverse reactions:* Vasopressin Injection may produce tonic uterine contractions that could threaten the continuation of pregnancy.

### **8.2 Lactation**

There are no data on the presence of vasopressin injection in either human or animal milk, the effects on the breastfed infant, or the effects on milk production.

### **8.4 Pediatric Use**

Safety and effectiveness of Vasopressin Injection in pediatric patients with vasodilatory shock have not been established.

### **8.5 Geriatric Use**

Clinical studies of vasopressin did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy [see *Warnings and Precautions (5)*, *Adverse Reactions (6)*, and *Clinical Pharmacology (12.3)*].

## **10 OVERDOSAGE**

Overdosage with Vasopressin Injection can be expected to manifest as consequences of vasoconstriction of various vascular beds (peripheral, mesenteric, and coronary) and as hyponatremia. In addition, overdosage may lead less commonly to ventricular tachyarrhythmias (including Torsade de Pointes), rhabdomyolysis, and non-specific gastrointestinal symptoms.

Direct effects will resolve within minutes of withdrawal of treatment.

## **11 DESCRIPTION**

Vasopressin is a polypeptide hormone. Vasopressin Injection, USP is a sterile, aqueous solution of synthetic arginine vasopressin for intravenous administration.



vasopressin is 9 to 25 mL/min/kg in patients with vasodilatory shock. The apparent  $t_{1/2}$  of vasopressin at these levels is  $\leq 10$  minutes.

#### Metabolism

Serine protease, carboxipeptidase and disulfide oxido-reductase cleave vasopressin at sites relevant for the pharmacological activity of the hormone. Thus, the generated metabolites are not expected to retain important pharmacological activity.

#### Excretion

Vasopressin is predominantly metabolized and only about 6% of the dose is excreted unchanged into urine.

#### Specific Populations

*Pregnancy:* Because of a spillover into blood of placental vasopressinase the clearance of exogenous and endogenous vasopressin increases gradually over the course of a pregnancy. During the first trimester of pregnancy, the clearance is only slightly increased. However, by the third trimester the clearance of vasopressin is increased about 4-fold and at term up to 5-fold. After delivery the clearance of vasopressin returns to preconception baseline within two weeks.

Drug Interactions Indomethacin more than doubles the time to offset for vasopressin's effect on peripheral vascular resistance and cardiac output in healthy subjects [see *Drug Interactions (7.2)*].

The ganglionic blocking agent tetra-ethylammonium increases the pressor effect of vasopressin by 20% in healthy subjects [see *Drug Interactions (7.3)*].

Halothane, morphine, fentanyl, alfentanyl and sufentanyl do not impact exposure to endogenous vasopressin.

## **13 NONCLINICAL TOXICOLOGY**

### **13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility**

No formal carcinogenicity or fertility studies with vasopressin have been conducted in animals. Vasopressin was found to be negative in the *in vitro* bacterial mutagenicity (Ames) test and the *in vitro* Chinese hamster ovary (CHO) cell chromosome aberration test. In mice, vasopressin has been reported to have an effect on function and fertilizing ability of spermatozoa.

### **13.2 Animal Toxicology and/or Pharmacology**

No toxicology studies were conducted with vasopressin.

## **14 CLINICAL STUDIES**

Increases in systolic and mean blood pressure following administration of vasopressin were observed in 7 studies in septic shock and 8 in post-cardiotomy vasodilatory shock.

## **16 HOW SUPPLIED/STORAGE AND HANDLING**

Vasopressin Injection, USP is a clear, colorless to practically colorless solution for intravenous administration available as:

NDC 42367-570-87: A carton containing 25 x 1 mL multiple dose vials. Each vial contains

vasopressin 1 mL at 20 units/mL.

Store between 2°C and 8°C (36°F and 46°F). Do not freeze.

Vials may be held up to 12 months upon removal from refrigeration to room temperature storage conditions (20°C to 25°C [68°F to 77°F], USP Controlled Room Temperature), anytime within the labeled shelf life. Once removed from refrigeration, unopened vial should be marked to indicate the revised 12 month expiration date. If the manufacturer's original expiration date is shorter than the revised expiration date, then the shorter date must be used. Do not use Vasopressin Injection beyond the manufacturer's expiration date stamped on the vial.

Discard vial after 48 hours after first puncture.

The storage conditions and expiration periods are summarized in the following table.

	Unopened Refrigerated 2°C to 8°C (36°F to 46°F)	Unopened Room Temperature 20°C to 25°C (68°F to 77°F) Do not store above 25°C (77°F)	Opened (After First Puncture)
1 mL Vial	Until manufacturer expiration date	12 months or until manufacturer expiration date, whichever is earlier	48 hours

Marketed by:  
**Eagle Pharmaceuticals, Inc.**  
Woodcliff Lake, NJ 07677

Rev. 06/2021

**PRINCIPAL DISPLAY PANEL - NDC: 42367-570-87 - Vial Carton**



**PRINCIPAL DISPLAY PANEL - NDC: 42367-570-22 - Vial Label**

<p>NDC 42367-570-22  <b>Vasopressin</b>          Injection, USP  <b>20 Units per mL</b>          For Intravenous Infusion          Must be diluted          prior to use          1 mL Multiple Dose Vial          Rx Only</p>	<p>Store between 2°C and 8°C (36°F and 46°F). Vials may be held at 20°C to 25°C (68°F to 77°F) for up to 12 months. Do not freeze. <b>Usual Dosage:</b> See full prescribing information. After initial entry into the vial, the remaining contents must be used within 48 hours.          Marketed by: Eagle Pharmaceuticals, Inc.          Woodcliff Lake, NJ 07677</p>		<p>124520</p>	<p>LOT          EXP</p>
--	---	--	---------------	-----------------------------

**VASOPRESSIN**  
 vasopressin injection

**Product Information**

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:42367-570
---------------------	-------------------------	---------------------------	---------------

**Route of Administration** PARENTERAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
VASOPRESSIN (UNII: Y4907O6MFD) (VASOPRESSIN - UNII:Y4907O6MFD)	VASOPRESSIN	20 U in 1 mL

### Inactive Ingredients

Ingredient Name	Strength
CHLOROBUTANOL HEMIHYDRATE (UNII: 3X4P6271OX)	
WATER (UNII: 059QF0K00R)	
ACETIC ACID (UNII: Q40Q9N063P)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:42367-570-87	25 in 1 CARTON	01/10/2022	06/30/2024
1	NDC:42367-570-22	1 mL in 1 VIAL, MULTI-DOSE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA211538	01/10/2022	06/30/2024

**Labeler** - Eagle Pharmaceuticals, Inc. (849818161)

### Establishment

Name	Address	ID/FEI	Business Operations
Curia New Mexico, LLC		078268825	MANUFACTURE(42367-570) , ANALYSIS(42367-570)

Revised: 3/2023

Eagle Pharmaceuticals, Inc.