

MUCINEX SINUS-MAX SEVERE CONGESTION RELIEF- acetaminophen, guaifenesin, and phenylephrine hydrochloride tablet, coated
RB Health (US) LLC

Mucinex® Sinus-Max®
Severe Congestion Relief

Drug Facts

Active ingredients (in each caplet)	Purpose
Acetaminophen 325 mg	Pain reliever
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves:
 - nasal congestion
 - headache
 - minor aches and pains
 - sinus congestion and pressure
- temporarily promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blister
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen**(prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain

drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see Overdose warning)**
- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years of age and older: take 2 caplets every 4 hours
- children under 12 years of age: do not use

Other information

- store between 20-25°C (68-77°F)

Inactive ingredients

croscarmellose sodium, crospovidone, FD&C red no. 40 aluminum lake, FD&C yellow no. 6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, talc, titanium dioxide

Questions?

1-866-MUCINEX (1-866-682-4639) You may also report side effects to this phone number.

Dist. by: Reckitt Benckiser, Parsippany, NJ 07054-0224
Made in England

PRINCIPAL DISPLAY PANEL - 20 Caplet Blister Pack Carton

MAXIMUM STRENGTH*

NDC 63824-201-20

Mucinex®

SINUS- MAX®

SEVERE CONGESTION RELIEF

Acetaminophen– Pain Reliever • Guaifenesin – Expectorant
Phenylephrine HCl – Nasal Decongestant

- **Clears Sinus Congestion**
- **Relieves Headache**
- **Thins & Loosens Mucus**

FOR AGES 12+

20 CAPLETS

LOT
EXP.

Drug Facts

Active Ingredients **Purposes**
(in each caplet) (in each caplet)

Acetaminophen 325 mg.....Pain reliever
Guaifenesin 200 mg.....Expectorant
Phenylephrine HCl 5 mg.....Nasal decongestant

Uses

- temporarily relieves:
 - nasal congestion
 - headache
 - minor aches and pains
 - sinus congestion and pressure
- temporarily promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passages of bothersome mucus and make coughs more productive

Warnings

Liver warnings: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 12 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin redness
- blister
- rash

 If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- kidney disease
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland

Drug Facts (continued)

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin. When using this product do not use more than directed

Stop use and ask a doctor if

- feverishness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- ever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or hives that last. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years of age and older: take 2 caplets every 4 hours
- children under 12 years of age: do not use

Other information

- store between 20°-25°C (68°-77°F)

Inactive ingredients

croscarmellose sodium, crospovidone, FD&C red no. 40 aluminum lake, FD&C yellow no. 5 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, talc, titanium dioxide

MAXIMUM STRENGTH*

Mucinex
SINUS-MAX

SEVERE CONGESTION RELIEF

MAXIMUM STRENGTH*

Mucinex
SINUS-MAX

SEVERE CONGESTION RELIEF

Acetaminophen – Pain Reliever • Guaifenesin – Expectorant
Phenylephrine HCl – Nasal Decongestant

- ✓ Clears Sinus Congestion
- ✓ Relieves Headache
- ✓ Thins & Loosens Mucus

FOR AGES 12+

20 CAPLETS

Questions? 1-866-MUCINEX (1-866-682-4639)
You may also report side effects to this phone number.

Dist. by: Reckitt Benckiser, Parsippany, NJ 07054-0224
Made in England ©2016 RB www.mucinex.com
070916 3022055

3 63824 20120 7

Per 4-hour dose

Tamper evident: Do not use if carton is damaged or if printed seal on blister is broken or missing.

Keep carton for complete product information.

NDC 63824-201-20

MUCINEX SINUS-MAX SEVERE CONGESTION RELIEF
acetaminophen, guaifenesin, and phenylephrine hydrochloride tablet, coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63824-201
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
CROSPROVIDONE (15 MPAS AT 5%) (UNII: 68401960MK)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	orange	Score	no score
Shape	OVAL	Size	20mm
Flavor		Imprint Code	VV;CS
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63824-201-20	2 in 1 CARTON	07/23/2012	11/15/2018
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:63824-201-30	3 in 1 CARTON	07/23/2012	11/15/2018
2		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:63824-201-72	2 in 1 POUCH; Type 0: Not a Combination Product	07/23/2012	11/15/2018
4	NDC:63824-201-44	4 in 1 CARTON	07/23/2012	
4	NDC:63824-201-72	2 in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/23/2012	

Labeler - RB Health (US) LLC (081049410)

Revised: 11/2023

RB Health (US) LLC