

**EXCEDRIN TENSION HEADACHE- acetaminophen, caffeine tablet, film coated**  
**GlaxoSmithKline Consumer Healthcare Holdings (US) LLC**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Drug Facts**

***Active ingredients (in each caplet)***

Acetaminophen 500 mg

Caffeine 65 mg

***Purposes***

Pain reliever

Pain reliever aid

***Uses***

- temporarily relieves minor aches and pain due to:
- headache
- muscular aches

***Warnings***

**Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:**

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take**

- more than 6 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and occasionally, rapid heartbeat.**

**Do not use**

- if you are allergic to acetaminophen
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not

sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if**

- you have liver disease

**Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin

**Stop use and ask a doctor if**

- any new symptoms occur
- painful area is red or swollen
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days

These could be signs of a serious condition

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

***Directions***

- **do not use more than directed**
- adults and children 12 years and over: take 2 caplets every 6 hours; not more than 6 caplets in 24 hours
- children under 12 years: ask a doctor

***Other information***

- store at 20°-25°C (68°-77°F)
- close cap tightly after use
- read all product information before using. Keep this box for important information.

***Inactive ingredients***

benzoic acid, D&C red #27 aluminum lake, FD&C blue #2 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, stearic acid, talc, titanium dioxide

***Questions or comments?***

**1-800-452-0051**

**Principal Display Panel**

NDC 0067-8139-02

**EXCEDRIN®**

**TENSION HEADACHE**

*Acetaminophen and Caffeine*

*Pain Reliever / Pain Reliever Aid*

*Aspirin Free*

**100 CAPLETS**

**TAMPER-EVIDENT BOTTLE**

**DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH “SEALED for YOUR PROTECTION” IS BROKEN OR MISSING**

Distributed by: **GSK Consumer Healthcare,**

Warren, NJ 07059

©2016 GSK group of companies or its licensor. All rights reserved. Trademarks are owned by or licensed to the GSK group of companies

**Visit us at [www.excedrin.com](http://www.excedrin.com)**

46177300

NDC 0067-8139-02

# EXCEDRIN®

## TENSION HEADACHE

*Aspirin Free*

*Acetaminophen and Caffeine*  
*Pain Reliever / Pain Reliever Aid*

**100**  
**CAPLETS**



### EXCEDRIN TENSION HEADACHE

acetaminophen, caffeine tablet, film coated

#### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0067-8139
Route of Administration	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg

**Inactive Ingredients**

Ingredient Name	Strength
BENZOIC ACID (UNII: 8SKN0B0MM)	
D&C RED NO. 27 ALUMINUM LAKE (UNII: ZK64F7XSTX)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TALC (UNII: 7SEV7J4R1U)	

**Product Characteristics**

Color	RED	Score	no score
Shape	OVAL ((Caplet))	Size	17mm
Flavor		Imprint Code	ETH
Contains			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0067-8139-01	1 in 1 CARTON	12/15/2016	
1		24 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:0067-8139-02	1 in 1 CARTON	12/15/2016	
2		100 in 1 BOTTLE; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	12/15/2016	

**Labeler** - GlaxoSmithKline Consumer Healthcare Holdings (US) LLC (079944263)