

CURATUSSIN DM- dextromethorphan hbr/guaifenesin liquid
Wittman Pharma, Inc.

CuraTUSSIN DM

Directions:

- follow dosage below or use as directed by a physician
- do not take more than 6 doses in any 24-hour period

| Age | Dose |
|--|---------------------------------|
| adults and children 12 years and over | 2 teaspoonfuls every 4 hours |
| children 6 years to under 12 years | 1 teaspoonfuls every 4 hours |
| children 2 years to under 6 years | ½ teaspoonfuls every 4 hours |
| under 2 years | ask a doctor |

Warnings: Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI)

(certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2

weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI; ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic, such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Stop use and ask doctor if

■ cough lasts more than 7 days., comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

■ you are hypersensitive to any of the ingredients.

If pregnant or breastfeeding, ask a health professional before use. Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Inactive Ingredients: Citric Acid, Flavor, Methylparaben, Monoammonium Glycyrhizinate, Potassium Citrate, Propylene Glycol, Propylparaben, Purified Water, Sorbitol, Sucralose

Uses:

- helps loosen phlegm(mucus) and thin bronchial secretions to make coughs more productive

■ temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Purpose:

Cough Suppressant and Expectorant

Active Ingredient (in each 5 mL tsp)

Dextromethorphan HBr 10mg

Guaifenesin 100mg

| Drug Facts | |
|--|-------------------|
| Active Ingredient (in each 5 mL tsp) | Purpose |
| Dextromethorphan HBr 10 mg..... | Cough Suppressant |
| Guaifenesin 100 mg..... | Expectorant |
| Uses: ■ helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive ■ temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold | |
| Warnings: Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. | |
| Ask a doctor before use if you have ■ cough that occurs with too much phlegm (mucus) ■ cough that lasts or is chronic, such as occurs with smoking, asthma, chronic bronchitis, or emphysema | |
| Stop use and ask doctor if ■ cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition. ■ you are hypersensitive to any of the ingredients. | |
| If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. ▶ | |
|  3 83335 10216 9 | |

MADE IN USA NDC 83335-102-16



CuraTUSSIN DM

Cough Suppressant
Expectorant

Each teaspoonful (5 mL) for oral administration contains:
 Dextromethorphan HBr 10 mg
 Guaifenesin 100 mg

Compare to the active ingredient in *Robitussin®DM

Loosens & Relieves Chest Congestion

✓SUGAR FREE ✓ALCOHOL FREE
 ✓DYE FREE ✓GLUTEN FREE
 ✓NON-NARCOTIC

Grape Flavor

 **Wittman
Pharma**

Net Content: 16 fl oz (473 mL)

| Drug Facts (continued) | |
|---|------------------------------|
| Directions: ■ follow dosage below or use as directed by a physician ■ do not take more than 6 doses in any 24-hour period | |
| Age | Dose |
| adults and children 12 years and over | 2 teaspoonfuls every 4 hours |
| children 6 years to under 12 years | 1 teaspoonfuls every 4 hours |
| children 2 years to under 6 years | ½ teaspoonfuls every 4 hours |
| under 2 years | ask a doctor |
| Other information ■ store at 15°-30°C (59°-86°F) ■ protect from light ■ tamper evident, do not use if safety seal under cap is broken or missing | |
| Inactive Ingredients: Citric Acid, Flavor, Methylparaben, Monoammonium Glycyrhizinate, Potassium Citrate, Propylene Glycol, Propylparaben, Purified Water, Sorbitol, Sucralose | |
| Questions? Call 1-352-549-9917 M-F 9am - 5pm EST Manufactured for: Wittman Pharma, Inc. Brooksville, FL 34604 | |

*Robitussin®DM is a registered trademark of Wyeth Consumer Healthcare. Rev. 3/2024

CURATUSSIN DM

dextromethorphan hbr/guaifenesin liquid

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:83335-102 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|-------------------------------|-------------------|
| GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ) | GUAIFENESIN | 100 mg in 5 mL |
| DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS) | DEXTROMETHORPHAN HYDROBROMIDE | 10 mg in 5 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|--|----------|
| ANHYDROUS CITRIC ACID (UNII: XF417D3PSL) | |
| AMMONIUM GLYCYRRHIZATE (UNII: 3VRD35U26C) | |
| PROPYLPARABEN (UNII: Z8IX2SC1OH) | |
| POTASSIUM CITRATE (UNII: EE90ONI6FF) | |
| METHYLPARABEN (UNII: A2I8C7HI9T) | |
| WATER (UNII: 059QF0KO0R) | |
| SUCRALOSE (UNII: 96K6UQ3ZD4) | |
| SORBITOL (UNII: 506T60A25R) | |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3) | |

Product Characteristics

| | | | |
|-----------------|-------|---------------------|--|
| Color | | Score | |
| Shape | | Size | |
| Flavor | GRAPE | Imprint Code | |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:83335-102-16 | 473 mL in 1 BOTTLE, PUMP; Type 0: Not a Combination Product | 03/01/2024 | |
| 2 | NDC:83335-102-08 | 237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | 03/01/2024 | |
| 3 | NDC:83335-102-04 | 118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | 03/01/2024 | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M012 | 03/01/2024 | |

Labeler - Wittman Pharma, Inc. (830980947)

Establishment

| Name | Address | ID/FEI | Business Operations |
|----------------------|---------|-----------|---|
| Wittman Pharma, Inc. | | 830980947 | analysis(83335-102) , manufacture(83335-102) , label(83335-102) |

Revised: 4/2024

Wittman Pharma, Inc.