

MAXI-TUSS TR- pseudoephedrine hydrochloride and triprolidine hydrochloride liquid
MCR American Pharmaceuticals, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Maxi-Tuss TR

Drug Facts

<i>Active Ingredients (in each 5 mL teaspoonful)</i>	<i>Purpose</i>
Pseudoephedrine HCl 30 mg	Nasal Decongestant
Triprolidine HCl 1.25 mg	Antihistamine

Uses

temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other respiratory allergies:

- runny nose
- sneezing
- itching of the nose or throat
- itchy, watery eyes
- nasal congestion
- reduces swelling of nasal passages

Warnings

Do not exceed recommended dosage.

Do not use this product

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product

Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- heart disease
- high blood pressure
- thyroid disease
- diabetes mellitus
- difficulty in urination due to enlargement of the prostate gland

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product

- excitability may occur, especially in children
- may cause drowsiness
- alcohol, sedatives and tranquilizers may increase drowsiness

- avoid alcoholic beverages
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not improve within 7 days or are accompanied by fever
- new symptoms occur

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of accidental overdose, seek professional help or contact a Poison Control Center immediately.

Directions

Do not exceed recommended dosage.

Adults and children 12 years of age and over:	2 teaspoonfuls (10 mL) every 4-6 hours, not to exceed 8 teaspoonfuls in 24 hours
Children 6 to under 12 years of age:	1 teaspoonful (5 mL) every 4-6 hours, not to exceed 4 teaspoonfuls in 24 hours
Children under 6 years of age:	Consult a physician

Other information

Store at 59°-86°F (15°-30°C) [see USP Controlled Room Temperature]

Inactive ingredients

Bubblegum flavor, citric acid, methylparaben, monoammonium glycyrrhizinate, potassium citrate, propylparaben, propylene glycol, purified water, sorbitol, sucralose

Questions or comments?

Call 352.754.8587

PRINCIPAL DISPLAY PANEL - 473 mL Bottle Label

NDC 58605-305-16

Maxi-Tuss TR

Nasal Decongestant □ Antihistamine

Sugar Free □ Alcohol Free □ Dye Free

Each teaspoonful (5 mL) for oral administration contains:

Pseudoephedrine HCl

30 mg

Triprolidine HCl

1.25 mg

Bubblegum Flavor

Tamper evident by foil seal under cap.
Do not use if foil seal is broken or missing.

Manufactured for:
MCR American Pharmaceuticals, Inc.
Brooksville, FL 34604

16 fl oz (473 mL)

NDC 58605-305-16

Maxi-Tuss TR

Nasal Decongestant ■ Antihistamine

Sugar Free ■ Alcohol Free ■ Dye Free

Each teaspoonful (5 mL) for oral administration contains:
Pseudoephedrine HCl 30 mg
Triprolidine HCl 1.25 mg

Bubblegum Flavor

Tamper evident by foil seal under cap.
Do not use if foil seal is broken or missing.

Manufactured for:
MCR American Pharmaceuticals, Inc.
Brooksville, FL 34604

Lot: Exp. Date:



N 3 58605 30516 8

16 fl oz (473 mL)

Drug Facts

Active Ingredients (in each 5 mL teaspoonful)

Pseudoephedrine HCl 30 mg	Nasal Decongestant
Tripolidine HCl 1.25 mg	Antihistamine

Uses temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other respiratory allergies:
 ■ runny nose ■ sneezing ■ itching of the nose or throat
 ■ itchy, watery eyes ■ nasal congestion ■ reduces swelling of nasal passages

Warnings
Do not exceed recommended dosage.
Do not use this product
 ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product

Ask a doctor before use if you have
 ■ a breathing problem such as emphysema or chronic bronchitis ■ glaucoma ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes mellitus ■ difficulty in urination due to enlargement of the prostate gland

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product
 ■ excitability may occur, especially in children
 ■ may cause drowsiness
 ■ alcohol, sedatives and tranquilizers may increase drowsiness
 ■ avoid alcoholic beverages
 ■ use caution when driving a motor vehicle or operating machinery

Drug Facts (continued)

Stop use and ask a doctor if
 ■ nervousness, dizziness, or sleeplessness occur
 ■ symptoms do not improve within 7 days or are accompanied by fever
 ■ new symptoms occur

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.
 In case of accidental overdose, seek professional help or contact a Poison Control Center immediately.

Directions
Do not exceed recommended dosage.

Adults and children 12 years of age and over:	2 teaspoonfuls (10 mL) every 4-6 hours, not to exceed 8 teaspoonfuls in 24 hours
Children 6 to under 12 years of age:	1 teaspoonful (5 mL) every 4-6 hours, not to exceed 4 teaspoonfuls in 24 hours
Children under 6 years of age:	Consult a physician

Other information
 Store at 59°-86°F (15°-30°C) [see USP Controlled Room Temperature]

Inactive ingredients
 Bubblegum flavor, citric acid, methylparaben, monoammonium glycyrrhizinate, potassium citrate, propylparaben, propylene glycol, purified water, sorbitol, sucralose

Questions or comments?
 Call 352.754.8587

Rev. 05/20

MAXI-TUSS TR

pseudoephedrine hydrochloride and triprolidine hydrochloride liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58605-305
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Pseudoephedrine Hydrochloride (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)	Pseudoephedrine Hydrochloride	30 mg in 5 mL
Tripolidine Hydrochloride (UNII: YAN7R5L890) (TRIPROLIDINE - UNII:2L8T9S52QM)	Tripolidine Hydrochloride	1.25 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
Methylparaben (UNII: A2I8C7HI9T)	
AMMONIUM GLYCYRRHIZATE (UNII: 3VRD35U26C)	
Potassium Citrate (UNII: EE90ONI6FF)	
Propylparaben (UNII: Z8IX2SC1OH)	
Propylene Glycol (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
Sorbitol (UNII: 506T60A25R)	
Sucralose (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	BUBBLE GUM	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58605-305-16	473 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	08/01/2020	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part341	08/01/2020	

Labeler - MCR American Pharmaceuticals, Inc. (783383011)

Establishment

Name	Address	ID/FEI	Business Operations
MCR American Pharmaceuticals, Inc.		783383011	MANUFACTURE(58605-305)

Revised: 6/2020

MCR American Pharmaceuticals, Inc.