

**CAREONE MILK OF MAGNESIA ORIGINAL MILK OF MAGNESIA ORIGINAL-
magnesium hydroxide liquid
Retail Business Services, LLC.**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

CAREone Milk of Magnesia original

ACTIVE INGREDIENT(S)

Magnesium hydroxide 1200 mg

PURPOSE

Saline laxative

USE(S)

relieves occasional constipation (irregularity). This product usually produces bowel movement in 1/2 to 6 hours.

WARNINGS

.

ASK A DOCTOR BEFORE USE IF YOU HAVE

- kidney disease
- a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts over 14 days

ASK A DOCTOR OR PHARMACIST BEFORE USE IF YOU ARE

taking a prescription drug. This product may interact with certain prescription drugs.

STOP USE AND ASK DOCTOR IF

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

IF PREGNANCY OR BREAST-FEEDING,

ask a health professional before use.

KEEP OUT OF REACH OF CHILDREN

In case of overdose, get medical help or contact a Poison Control Center right away.

DIRECTIONS

- shake well before use
- do not exceed the maximum recommended daily dose in a 24 hour period
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor. Drink a full glass (8 OZ) of liquid with each dose.
- for accurate dosing, only use the dosing cup provided
- mL=milliliter

adults and children 12 years and older	30 mL to 60 mL
children 6 to 11 years	15 mL to 30 mL
children under 6years	ask a doctor

OTHER INFORMATION

- **each 15 mL contains: magnesium 500 mg**
- store at 20 - 25°C (68 - 77°F), Avoid excessive heat above 40°C(104°F).
Avoid freezing. Close cap tightly after use.

INACTIVE INGREDIENTS

citric acid, glycerin, hydroxyethyl cellulose, purified water, sodium hypochlorite, sucrose.

PRINCIPAL DISPLAY PANEL

CAREone™

NDC : 72476-661-05

Compare to the active ingredients in Phillips^{1®} Milk of Magnesia*

MILK OF MAGNESIA

Saline Laxative

Magnesium Hydroxide 1200 mg per 15 mL

Original Flavor

Comfortable, Cramp Free Relief!

Stimulant Free

12FL OZ (355 mL)

Drug Facts (continued)

adults and children 12 years and older	30 mL to 60 mL
children 6 to 11 years	15 mL to 30 mL
children under 6 years	ask a doctor

Other information ■ each 15 mL contains: magnesium 500 mg ■ store at 20° - 25°C (68° - 77°F). Avoid excessive heat above 40°C (104°F). Avoid freezing. Close cap tightly after use.

Inactive ingredients citric acid, glycerin, hydroxyethyl cellulose, purified water, sodium hypochlorite, sucrose.

CAREoneTM NDC 72476-661-05

Compare to the active ingredient in
*Phillips'® Milk of Magnesia**

MILK OF MAGNESIA

Saline Laxative

Magnesium Hydroxide 1200 mg per 15 mL

Original Flavor

Comfortable, Cramp Free Relief!

Stimulant Free



12 FL OZ (355mL)

TAMPER-EVIDENT: DO NOT USE IF PRINTED NECKBAND IS TORN OR MISSING.

DISTRIBUTED BY:
ADUSA DISTRIBUTION, LLC
SALISBURY, NC 28147
1-833-992-3872
Quality guaranteed or your money back.

*This product is not manufactured by Bayer Corporation, the distributor of Phillips'® Milk of Magnesia.

MFR# 53041 REV 0022

LOT: EXP:


3 41520 00619 7

Drug Facts

Active ingredient (in each 15 mL)	Purpose
Magnesium hydroxide 1200 mg.....	Saline laxative

Use

■ relieves occasional constipation (irregularity). This product usually produces bowel movement in ½ to 6 hours.

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts over 14 days

Ask a doctor or pharmacist before use if you are taking a prescription drug. This product may interact with certain prescription drugs.

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ shake well before use ■ do not exceed the maximum recommended daily dose in a 24 hour period ■ dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor. Drink a full glass (8 oz) of liquid with each dose. ■ for accurate dosing, only use the dosing cup provided ■ mL = milliliter

CAREONE MILK OF MAGNESIA ORIGINAL MILK OF MAGNESIA ORIGINAL

magnesium hydroxide liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:72476-661
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Magnesium Hydroxide (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838)	Magnesium Hydroxide	1200 mg in 15 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
Glycerin (UNII: PDC6A3C0OX)	
HYDROXYETHYL CELLULOSE (2000 MPA.S AT 1%) (UNII: S38J6RZN16)	
water (UNII: 059QF0KO0R)	
Sodium Hypochlorite (UNII: DY38VHM5OD)	
Sucrose (UNII: C151H8M554)	

Product Characteristics

Color	WHITE	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72476-661-05	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	04/27/2022	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH NOT FINAL	part334	04/27/2022	

Labeler - Retail Business Services, LLC. (967989935)

Revised: 4/2022

Retail Business Services, LLC.