

KINEVAC- sincalide injection, powder, lyophilized, for solution Bracco Diagnostics Inc.

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use KINEVAC safely and effectively. See full prescribing information for KINEVAC

KINEVAC (sincalide for injection), for intravenous use Initial U.S. Approval: 1976

RECENT MAJOR CHANGES

Contraindications (4)	03/2018
Warnings and Precautions (5.1)	03/2018

INDICATIONS AND USAGE

Kinevac is a cholecystokinin (CCK) analog indicated in adults to:

- stimulate gallbladder contraction, as may be assessed by various methods of diagnostic imaging, or to obtain by duodenal aspiration a sample of concentrated bile for analysis of cholesterol, bile salts, phospholipids, and crystals. (1)
- stimulate pancreatic secretion in combination with secretin prior to obtaining a duodenal aspirate for analysis of enzyme activity, composition, and cytology. (1)
- accelerate the transit of a barium meal through the small bowel, thereby decreasing the time and extent of radiation associated with fluoroscopy and x-ray examination of the intestinal tract. (1)

DOSAGE AND ADMINISTRATION

Recommended Adult Dosage and Administration by Indication:

To Stimulate Contraction of the Gallbladder

- 0.02 mcg/kg as a single dose over 30 to 60 seconds via intravenous injection. If satisfactory contraction does not occur in 15 minutes, administer a dose of 0.04 mcg/kg over 30 to 60 seconds. (2.1)
- Alternatively consider an intravenous infusion to reduce gastrointestinal adverse reactions:* 0.12 mcg/kg diluted in 100 mL of 0.9% Sodium Chloride Injection USP and infused over 50 minutes at a rate of 2 mL per minute. (2.1, 2.2, 5.3)

To Stimulate Pancreatic Secretion in Combination with Secretin

- 30 minutes after initiation of secretin for injection, administer 0.02 mcg/kg diluted in 30 mL of 0.9% Sodium Chloride Injection USP and infused over 30 minutes at a rate of 1 mL per minute. (2.1, 2.2)

To Accelerate Transit of a Barium Meal Through the Small Intestine

- After the barium meal is beyond the proximal jejunum, administer 0.04 mcg/kg over 30 to 60 seconds via intravenous injection. (2.1)
- If satisfactory transit of the barium meal has not occurred in 30 minutes, administer a second dose of 0.04 mcg/kg over 30 to 60 seconds. (2.1)
- Alternatively consider an intravenous infusion to reduce gastrointestinal adverse reactions:* 0.12 mcg/kg diluted in 100 mL 0.9% Sodium Chloride Injection USP and infused over 30 minutes. (2.1, 2.2, 5.3)

DOSAGE FORMS AND STRENGTHS

For injection: 5 mcg of sincalide as a lyophilized powder in a single-dose vial for reconstitution (3)

CONTRAINDICATIONS

- History of hypersensitivity to sincalide. (4, 5.1)
- Intestinal obstruction. (4)

WARNINGS AND PRECAUTIONS

- Anaphylaxis, Anaphylactic Shock and Other Hypersensitivity Reactions: May occur during or soon after administration. If symptoms occur, discontinue the drug. (4, 5.1)
- Evacuation of Gallstones: Stimulation of gallbladder contraction in patients with small gallbladder stones could lead to the evacuation of the stones from the gallbladder, resulting in their lodging in the cystic duct or in the common bile duct. (5.2)
- Gastrointestinal Adverse Reactions with Intravenous Injection: Administration as an intravenous injection may cause transient nausea, vomiting, abdominal pain or cramping, dizziness or flushing. To reduce the risk of adverse reactions when used to stimulate contraction of the gallbladder or accelerate transit of a barium meal through the small intestine, administer as an intravenous infusion over 50 or 30 minutes, respectively. (2.1, 5.3)
- Preterm Labor or Spontaneous Abortion: Advise pregnant women of the potential risk for preterm labor and

spontaneous abortion. (5.4, 8.1)

----- **ADVERSE REACTIONS** -----

Most common adverse reactions ($\geq 20\%$) are: abdominal discomfort or pain, and nausea. (6)

To report SUSPECTED ADVERSE REACTIONS, contact Bracco Diagnostics Inc. at 1-800-257-5181 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

----- **DRUG INTERACTIONS** -----

Drugs that Affect Gallbladder Motility or Contractile Response: May interfere with response to sincalide. Consider discontinuing these drugs prior to administration of Kinevac, when used to stimulate contraction of the gallbladder. (7.1)

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 12/2018

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

2.1 Recommended Dosage and Administration Instructions

2.2 Preparation Instructions

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

5.1 Anaphylaxis, Anaphylactic Shock and Other Hypersensitivity Reactions

5.2 Evacuation of Gallstones

5.3 Gastrointestinal Adverse Reactions with Intravenous Injection

5.4 Preterm Labor or Spontaneous Abortion

6 ADVERSE REACTIONS

7 DRUG INTERACTIONS

7.1 Drugs that Affect Gallbladder Motility or Contractile Response

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

8.2 Lactation

8.4 Pediatric Use

8.5 Geriatric Use

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

12.2 Pharmacodynamics

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

Kinevac is indicated in adults to:

- to stimulate gallbladder contraction, as may be assessed by various methods of diagnostic imaging, or to obtain by duodenal aspiration a sample of concentrated bile for analysis of cholesterol, bile salts, phospholipids, and crystals;
- to stimulate pancreatic secretion in combination with secretin prior to obtaining a duodenal aspirate for analysis of enzyme activity, composition, and cytology;
- to accelerate the transit of a barium meal through the small bowel, thereby decreasing the time and extent of radiation associated with fluoroscopy and x-ray examination of the intestinal tract.

2 DOSAGE AND ADMINISTRATION

2.1 Recommended Dosage and Administration Instructions

The recommended dosage and administration of Kinevac by indication is shown in Table 1. For preparation instructions see *Dosage and Administration (2.2)*.

Table 1: Recommended Adult Dosage and Administration of Kinevac by Treatment Indication

Indication	Recommended Adult Dosage and Administration of KINEVAC
To stimulate contraction of the gallbladder	Kinevac 0.02 mcg/kg as a single dose over 30 to 60 seconds via intravenous injection. If satisfactory contraction does not occur in 15 minutes, administer a dose of 0.04 mcg/kg over 30 to 60 seconds. <u>Alternatively, Consider an Intravenous Infusion to Reduce Gastrointestinal Adverse Reactions [see Warnings and Precautions (5.3)]:</u> 0.12 mcg/kg diluted in 100 mL of 0.9% Sodium Chloride Injection USP and infused over 50 minutes at a rate of 2 mL per minute.
To stimulate pancreatic secretion in combination with secretin for injection	Secretin for Injection: 0.25 units/kg as intravenous infusion over 60 minutes Kinevac: 30 minutes after initiation of secretin infusion, administer Kinevac 0.02 mcg/kg diluted in 30 mL of 0.9% Sodium Chloride Injection USP and infused over 30 minutes at a rate of 1 mL per minute.
To accelerate the transit of a barium meal through the small intestine	After the barium meal is beyond the proximal jejunum, administer Kinevac 0.04 mcg/kg over 30 to 60 seconds via intravenous injection. If satisfactory transit of the barium meal has not occurred in 30 minutes, administer a second dose of 0.04 mcg/kg over 30 to 60 seconds. <u>Alternatively, Consider an Intravenous Infusion to Reduce Gastrointestinal Adverse Reactions [see Warnings and Precautions (5.3)]:</u> 0.12 mcg/kg diluted in 100 mL 0.9% Sodium Chloride Injection USP and infused over 30 minutes.

2.2 Preparation Instructions

For Intravenous Injection

- Reconstitute Kinevac aseptically by adding 5 mL of Sterile Water for Injection USP to the vial.
- Inspect the reconstituted solution visually for particulate matter and discoloration after reconstitution and prior to administration.
- Withdraw the prescribed dose of the reconstituted solution from the vial and administer as an intravenous injection over 30 to 60 seconds, as shown in Table 1. Discard the unused portion.
- Store the reconstituted solution at room temperature. Discard after 8 hours.
- For single use only; discard unused portion.

For Intravenous Infusion

- Reconstitute Kinevac aseptically by adding 5 mL of Sterile Water for Injection USP to the vial.
- After reconstitution, withdraw the prescribed dose of the solution from the vial. Discard unused portion.
- Dilute the reconstituted solution in 30 mL or 100 mL of 0.9% Sodium Chloride Injection USP,

depending on the indication, as described in Table 1.

- Inspect the Kinevac solutions visually for particulate matter and discoloration after reconstitution, dilution and prior to administration.
- Store the diluted solution at room temperature. *Discard after 1 hour.*

3 DOSAGE FORMS AND STRENGTHS

For injection: 5 mcg of sincalide as a lyophilized white powder for reconstitution in a single-dose vial.

4 CONTRAINDICATIONS

KINEVAC is contraindicated in patients with:

- a history of hypersensitivity to sincalide. Serious hypersensitivity reactions have included anaphylaxis and anaphylactic shock [see *Warnings and Precautions (5.1), Adverse Reactions (6)*].
- intestinal obstruction.

5 WARNINGS AND PRECAUTIONS

5.1 Anaphylaxis, Anaphylactic Shock and Other Hypersensitivity Reactions

In postmarketing experience, anaphylaxis, anaphylactic shock and other serious hypersensitivity reactions have been reported during and within one hour following administration of Kinevac [see *Adverse Reactions (6)*].

Due to the potential for anaphylaxis, appropriate medical support should be readily available when Kinevac is administered. If anaphylaxis or other hypersensitivity reactions occur, immediately discontinue the infusion and initiate appropriate medical treatment. Observe patients closely during and after the infusion. Do not reinitiate Kinevac in patients who have experienced symptoms of hypersensitivity [see *Contraindications (4)*].

5.2 Evacuation of Gallstones

Stimulation of gallbladder contraction in patients with small gallbladder stones could lead to the evacuation of the stones from the gallbladder, resulting in their lodging in the cystic duct or in the common bile duct.

5.3 Gastrointestinal Adverse Reactions with Intravenous Injection

Administration of Kinevac as an intravenous injection may cause adverse reactions such as nausea, vomiting, abdominal pain or cramping, dizziness, and flushing [see *Adverse Reactions (6)*]. These reactions are generally transient. To reduce the risk of adverse reactions with intravenous injection when used to simulate contraction of the gallbladder or accelerate transit of a barium meal through the small intestine, administer Kinevac as an intravenous infusion over 50 or 30 minutes, respectively [see *Dosage and Administration (2.1)*].

5.4 Preterm Labor or Spontaneous Abortion

Because of Kinevac's effect on smooth muscle, pregnant patients should be advised that spontaneous abortion or premature induction of labor may occur [see *Use in Specific Populations (8.1)*].

6 ADVERSE REACTIONS

The following adverse reactions associated with the use of Kinevac were identified in clinical trials or postmarketing reports. Because these reactions were reported voluntarily from a population of uncertain size, it is not always possible to estimate their frequency, reliably, or to establish a causal

relationship to drug exposure.

The most frequent adverse reactions (20% or greater) are gastrointestinal: abdominal discomfort or pain, and nausea; these may not necessarily indicate an abnormality of the biliary tract unless there is other clinical or radiologic evidence of disease.

Less common adverse reactions include:

Hypersensitivity reactions: anaphylaxis and anaphylactic shock, hypotension, throat tightness, bradycardia, shortness of breath, nausea, abdominal cramping, diaphoresis, hives, rash, itching; and numbness of face, lips and eyes [see *Contraindications (4), (5.1)*].

Neurological reactions: seizures, headache.

Vasovagal reactions: dizziness, loss of consciousness, nausea, diaphoresis, syncope and hypotension (generally self-limiting).

Other: nausea, vomiting, flushing, hypertension, urge to defecate, diarrhea, sneezing.

7 DRUG INTERACTIONS

7.1 Drugs that Affect Gallbladder Motility or Contractile Response

Drugs that may stimulate or inhibit gallbladder motility or contractile response may interfere with the response to sincalide. Consider discontinuing these drugs prior to administration of Kinevac, when used to simulate contraction of the gallbladder.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

Based on limited human data and mechanism of action, sincalide may cause preterm labor or spontaneous abortion [see *Warnings and Precautions (5.4)*]. Available data with sincalide for injection are insufficient to establish a drug-associated risk of major birth defects, miscarriage or adverse maternal or fetal outcomes. In animal embryo-fetal development studies in which sincalide was administered to hamsters and rats during the period of organogenesis, no effects were seen at doses comparable to the maximum recommended clinical dose on a mg/kg basis. However, in a prenatal development study in which rats were administered sincalide during organogenesis through parturition, decreased weight gain and developmental delays were observed at a dose 122 times higher than the maximum recommended human dose based on body surface area.

The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2 to 4% and 15 to 20%, respectively.

Data

Animal Data

There were no effects on embryo-fetal development in hamsters when sincalide was administered subcutaneously at 250 or 750 ng/kg during organogenesis (Gestation Days 7 to 13) at doses up to 0.8 times the maximum recommended dose of 120 ng/kg on a body surface area basis. No effects on embryo-fetal development were observed in Sprague-Dawley rats at subcutaneous doses of 250, 450, or 750 ng/kg from Gestation Days 6 to 16, representing 1.0 time the maximum recommended human dose on a body surface area basis. In a separate study at a higher dose of 90 mcg/kg administered subcutaneously to CFY rats from Gestation Day 10 through parturition (representing 122 times the

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

When injected intravenously, sincalide stimulates gallbladder contraction and reduction in size. The evacuation of bile that results is similar to that which occurs physiologically in response to endogenous cholecystikinin. Sincalide also stimulates pancreatic secretion and intestinal motility causing pyloric contraction and slows gastric emptying.

Concurrent administration of sincalide with secretin increases both the volume of pancreatic secretion and the out-put of bicarbonate and enzymes. This combined effect of secretin and sincalide permits the assessment of specific pancreatic function through measurement and analysis of the duodenal aspirate.

12.2 Pharmacodynamics

Following an intravenous (bolus) injection of 0.02 mcg/kg of sincalide, maximal contraction of the gallbladder occurred in 5 to 15 minutes. Sincalide reduced gallbladder radiographic size by at least 40%, which is generally considered satisfactory contraction.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Long-term studies in animals have not been performed to evaluate carcinogenic or mutagenic potential, or possible impairment of fertility in males or females.

16 HOW SUPPLIED/STORAGE AND HANDLING

Kinevac (sincalide for injection) is supplied as 5 mcg of sincalide as a lyophilized white powder for reconstitution in a single-dose vial; in packages of 10 vials (NDC 0270-0556-15).

Store at 25° C (77° F); excursions permitted to 15-30° C (59-86° F) [See USP Controlled Room Temperature].

17 PATIENT COUNSELING INFORMATION

Anaphylaxis, Anaphylactic Shock and Other Hypersensitivity Reactions

Inform patients that hypersensitivity reactions, including anaphylaxis and anaphylactic shock have been reported during or following administration of Kinevac. Advise patients to report immediately to a healthcare provider if they experience symptoms of a hypersensitivity reaction [see Warnings and Precautions (5.1)].

Gastrointestinal Adverse Reactions

Advise patients that Kinevac may cause transient gastrointestinal symptoms [see Warnings and Precautions (5.3)].

Pregnancy

Advise pregnant women of the potential risk for preterm labor and spontaneous abortion [see Warnings and Precautions (5.4), Use in Specific Populations (8.1)].

U.S. Patent 6,803,046

Rx only

Manufactured for
Bracco Diagnostics Inc.

Monroe Township, NJ 08831
By Jubilant HollisterStier LLC
Spokane, WA 99207

Kinevac 5 mcg Label
NDC: 0270-0556-15



Bracco Diagnostics

10 vials

NDC 0270-0556-15

5 mcg per vial

KINEVAC[®]

**Sincalide
for Injection**

Rx only

Manufactured for
Bracco Diagnostics Inc.
Monroe Twp., NJ 08831
by Jubilant HollisterStier LLC
Spokane, WA 99207 USA

Each vial contains a sterile, lyophilized powder providing 5 mcg sincalide with 170 mg mannitol, 30 mg arginine hydrochloride, 15 mg lysine hydrochloride, 9 mg potassium phosphate dibasic, 4 mg methionine, 2 mg pentetic acid, 0.04 mg sodium metabisulfite, and 0.005 mcg polysorbate 20. pH adjusted to 6.0 - 8.0 with hydrochloric acid and/or sodium hydroxide. When reconstituted with 5 mL of Sterile Water for Injection USP, each mL contains 1 mcg sincalide.

For Intravenous use

Usual dose: See insert

Single Dose Vials • Discard Unused Portion

Store at 25 °C (77 °F) See insert

After reconstitution the solution may be stored at room temperature for up to 8 hours.

U.S. Patent 6,803,046

Made in USA

876657-H05



NDC3-0270-0556-15-4

NON-VARNISH AREA



NDC3-0270-0556-15-4

Kinevac 5 mcg Label
NDC: 0270-0556-15

Intravenous • Vial contains a sterile, lyophilized powder providing 5 mcg sincalide (see insert for inactive ingredients); pH adjusted to 6.0 - 8.0 with hydrochloric acid and/or sodium hydroxide. US Patent 6,803,046

Usual dose: See insert



Bracco Diagnostics

5 mcg NDC 0270-0556-15

KINEVAC[®]
Sincalide for Injection

Rx only

873093-H06



Single Dose Vial • Discard Unused Portion
Store at 25° C (77° F) See insert

Manufactured for

Bracco Diagnostics Inc.

Monroe Twp., NJ 08831

by Jubilant HollisterStier LLC

Spokane, WA 99207 USA

KINEVAC

sincalide injection, powder, lyophilized, for solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0270-0556
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Sincalide (UNII: M03GIQ7Z6P) (Sincalide - UNII:M03GIQ7Z6P)	Sincalide	5 ug in 5 mL

Inactive Ingredients

Ingredient Name	Strength
mannitol (UNII: 3OWL53L36A)	170 mg in 5 mL
arginine hydrochloride (UNII: F7LTH1E20Y)	30 mg in 5 mL
lysine hydrochloride (UNII: JNJ23Q2COM)	15 mg in 5 mL
potassium phosphate, dibasic (UNII: CI71S98N1Z)	9 mg in 5 mL
methionine (UNII: AE28F7PNPL)	4 mg in 5 mL
sodium metabisulfite (UNII: 4VON5FNS3C)	0.04 mg in 5 mL
polysorbate 20 (UNII: 7T1F30V5YH)	0.005 ug in 5 mL

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0270-0556-15	10 in 1 PACKAGE	07/21/1976	
1		5 mL in 1 VIAL; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA017697	07/21/1976	

Labeler - Bracco Diagnostics Inc. (849234661)

Registrant - Bracco Diagnostics Inc. (849234661)

Establishment

Name	Address	ID/FEI	Business Operations
Jubilant HollisterStier LLC		069263643	ANALYSIS(0270-0556) , MANUFACTURE(0270-0556)

Establishment

Name	Address	ID/FEI	Business Operations
Bachem AG		482220311	API MANUFACTURE(0270-0556)

Revised: 12/2018

Bracco Diagnostics Inc.