

**NIGHTTIME HBP COLD AND FLU RELIEF- acetaminophen, dextromethorphan
hbr, doxylamine succinate solution
CVS PHARMACY**

CVS 44-013 HBP

Active ingredients (in each 30 mL)

Acetaminophen 650 mg
Dextromethorphan HBr 30 mg
Doxylamine succinate 12.5 mg

Purpose

Pain reliever/fever reducer
Cough suppressant
Antihistamine

Uses

- temporarily relieves common cold and flu symptoms:
 - runny nose and sneezing
 - sore throat
 - headache
 - minor aches and pains
 - cough due to minor throat and bronchial irritation
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- difficulty in urination due to enlargement of the prostate gland
- cough that occurs with too much phlegm (mucus)
- liver disease
- glaucoma
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- excitability may occur, especially in children
- avoid alcoholic beverages
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed**

- do not exceed 4 doses per 24 hours
- mL = milliliter
- only use the dose cup provided
- adults and children 12 years and over: 30 mL every 6 hours
- children under 12 years: do not use

Other information

- **each 30 mL contains:** sodium 13 mg
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients

anhydrous citric acid, FD&C blue #1, FD&C red #40, flavor, glycerin, high fructose corn syrup, polyethylene glycol 400, propylene glycol, purified water, sodium benzoate, sodium chloride, sodium citrate dihydrate, sodium saccharin, sucralose

Questions or comments?

1-800-426-9391

Principal display panel

♥CVS

Health®

Compare to the active ingredients
in Vicks® NyQuil® HBP Cold & Flu*

Decongestant Free

Multi-Symptom

**Nighttime
High Blood
Pressure
COLD & FLU**

ACETAMINOPHEN

Pain reliever, Fever reducer

DEXTROMETHORPHAN HBR

Cough suppressant

DOXYLAMINE SUCCINATE

Antihistamine

For people with high blood pressure

Relieves: Aches & fever, Runny nose,
Sneezing, Cough

Cherry Flavor

Ages 12 & Over

12 FL OZ (355 mL)

TAMPER EVIDENT: DO NOT USE IF PRINTED NECK WRAP IS BROKEN OR MISSING

*This product is not manufactured or distributed by The Procter & Gamble Company, owner of the registered trademark Vicks® NyQuil® HBP Cold & Flu.

50844 ORG032301302

Distributed by : CVS Pharmacy, Inc.

One CVS Drive, Woonsocket, RI 02895

©2024 CVS/pharmacy

CVS.com® 1-800-SHOP CVS

V-19849

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

CVS Health. Compare to the active ingredients in Vicks® NyQuil® HBP Cold & Flu*
Decongestant Free

Multi-Symptom Nighttime High Blood Pressure

COLD & FLU

ACETAMINOPHEN
Pain reliever, Fever reducer

DEXTROMETHORPHAN HBr
Cough suppressant

DOXYLAMINE SUCCINATE
Antihistamine

For people with high blood pressure
Relieves: Aches & fever, Runny nose, Sneezing, Cough

Cherry Flavor

Ages 12 Years & Over

12 FL OZ (355 mL)

Drug Facts PEEL BACK FOR COMPLETE DRUG FACTS
TAMPER EVIDENT: DO NOT USE IF PRINTED NECK WRAP IS BROKEN OR MISSING

Active ingredients (in each 30 mL) Purpose
Acetaminophen 650 mg Pain reliever/fever reducer
Dextromethorphan HBr 30 mg Cough suppressant
Doxylamine succinate 12.5 mg Antihistamine

Uses temporarily relieves common cold and flu symptoms:
headache minor aches and pains
sore throat runny nose and sneezing
cough due to minor throat and bronchial irritation
temporarily reduces fever

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
more than 4,000 mg of acetaminophen in 24 hours
with other drugs containing acetaminophen
3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions.
Symptoms may include: skin reddening blisters rash
If a skin reaction occurs, stop use and seek medical help right away.

*This product is not manufactured or distributed by The Procter & Gamble Company, owner of the registered trademark Vicks® NyQuil® HBP Cold & Flu.

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2024 CVS/pharmacy
CVS.com®
1-800-SHOP CVS
V-19849

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

#442927

FPO 80%
UPC# 050428082423
X XXXXXX XXXXXX X

50844
ORG032301302
B-013 ORG

no print / no varnish area
lot no. & exp. date

Drug Facts (continued)

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have
difficulty in urination due to enlargement of the prostate gland liver disease glaucoma
a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are
taking the blood thinning drug warfarin
taking sedatives or tranquilizers

When using this product
marked drowsiness may occur
excitability may occur, especially in children
avoid alcoholic beverages
alcohol, sedatives, and tranquilizers may increase drowsiness
use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if pain or cough gets worse or lasts more than 7 days

Drug Facts (continued)

- redness or swelling is present
- fever gets worse or lasts more than 3 days
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions
do not take more than directed
do not exceed 4 doses per 24 hours
mL = milliliter
only use the dose cup provided
adults and children 12 years and over: 30 mL every 6 hours
children under 12 years: do not use

Other information
each 30 mL contains: sodium 13 mg
store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
use by expiration date on package

Inactive ingredients
anhydrous citric acid, FD&C blue #1, FD&C red #40, flavor, glycerin, high fructose corn syrup, polyethylene glycol 400, propylene glycol, purified water, sodium benzoate, sodium chloride, sodium citrate dihydrate, sodium saccharin, sucralose

Questions or comments?
1-800-426-9391

CVS 44-013

NIGHTTIME HBP COLD AND FLU RELIEF

acetaminophen, dextromethorphan hbr, doxylamine succinate solution

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:69842-086

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL
DOXYLAMINE SUCCINATE (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color	red (maroon)	Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69842-086-02	355 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	12/04/2020	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	12/04/2020	

Labeler - CVS PHARMACY (062312574)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	manufacture(69842-086) , pack(69842-086)

Revised: 4/2024

CVS PHARMACY