

MEDI-FIRST PLUS EXTRA STRENGTH NON-ASPIRIN- acetaminophen tablet, film coated

MEDI-FIRST EXTRA STRENGTH NON-ASPIRIN- acetaminophen tablet, film coated

MEDIQUE EXTRA STRENGTH APAP- acetaminophen tablet, film coated

Unifirst First Aid Corporation

Acetaminophen 500 mg

Drug Facts

Active Ingredient (in each tablet)

Acetaminophen 500 mg

Purpose

Pain reliever / fever reducer

Uses

temporarily relieves minor aches and pains due to:

- headache
- muscular aches
- backache
- minor pain of arthritis
- the common cold
- toothache
- premenstrual and menstrual cramps

temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have

- liver disease

Ask a doctor or pharmacist before use if

- you are taking the blood thinning drug warfarin

Stop using and ask a doctor if

- pain gets worse or lasts more than for more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

Keep out of reach of children.

Overdose Warning:

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not use more than directed (see overdose warning)**

Adults and children 12 years and over

- take 2 tablets every 6 hours while symptoms last.
- do not take more than 6 tablets in 24 hours, unless directed by a doctor
- do not use for more than 10 days unless directed by a doctor

Children under 12 years

Ask a doctor

Other information

- store at room temperature 59°-86°F (15°-30°C)
- tamper-evident sealed packets
- do not use any opened or torn packets

Inactive ingredients

corn starch*, hypromellose*, polyethylene glycol*, povidone (K-30)*, pregelatinized starch*, purified water*, sodium starch glycolate, stearic acid, titanium dioxide*

* May contain

Medique

Extra Strength APAP

Non-Aspirin Tablets

THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.

Pain Reliever / Fever Reducer • Acetaminophen 500 mg

Compare active ingredient to Tylenol®

Registered McNeil Products

Tamper Evident Unit Dose Packets

24 Tablets

(12 x 2)

Drug Facts (continued) Directions Do not use more than directed (see overdose warning) Adults and children (12 years and over) ■ take 2 tablets every 6 hours while symptoms last ■ do not take more than 6 tablets in 24 hours, unless directed by a doctor ■ do not use for more than 10 days, unless directed by a doctor Children under 12 years: Ask a doctor Other information ■ store at room temperature 59°-86°F (15°-30°C) ■ tamper-evident sealed packets ■ do not use any opened or torn packets Inactive ingredients corn starch, hypromellose, polyethylene glycol, povidone (K-30), pregelatinized starch, purified water, stearic acid, titanium dioxide *may contain Questions or comments? 1-800-634-7680	Drug Facts Active ingredient (in each tablet) Acetaminophen 500 mg Pain reliever/fever reducer Uses temporarily relieves minor aches and pains due to: ■ headache ■ muscular aches ■ minor pain of arthritis ■ backache ■ the common cold ■ toothache ■ the common cold ■ premenstrual and menstrual cramps temporarily reduces fever Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ hives ■ rash If a skin reaction occurs, stop use and seek medical help right away. Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are allergic to acetaminophen or any of the inactive ingredients in this product Ask a doctor before use if you have ■ liver disease (continued on opposite panel) RETAIN CARTON FOR COMPLETE PRODUCT INFORMATION	Recorder #17564 4768217564	Medique® Extra Strength APAP Non-Aspirin Tablets THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN. Pain Reliever/Fever Reducer • Acetaminophen 500 mg 24 Tablets (12 x 2) Compare active ingredient to Tylenol® Registered McNeil Consumer Products Tamper Evident Unit Dose Packets Manufactured for: Medique Products Fort Myers, FL 33967 1-800-634-7680 www.mediqueproducts.com	Drug Facts (continued) Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are allergic to acetaminophen or any of the inactive ingredients in this product Ask a doctor before use if you have ■ liver disease Ask a doctor or pharmacist before use if ■ you are taking the blood thinning drug warfarin Stop use and ask a doctor if ■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. Directions Do not use more than directed (see overdose warning) Adults and children (12 years and over) ■ take 2 tablets every 6 hours while symptoms last ■ do not take more than 6 tablets in 24 hours, unless directed by a doctor ■ do not use for more than 10 days unless directed by a doctor Children under 12 years: Ask a doctor Other information ■ store at room temperature 59°-86°F (15°-30°C) ■ tamper-evident sealed packets ■ do not use any opened or torn packets Questions or comments? 1-800-634-7680
---	---	-------------------------------	---	--

Drug Facts Active ingredient (in each tablet) Acetaminophen 500 mg Pain reliever/fever reducer Uses temporarily relieves minor aches and pains due to: ■ headache ■ muscular aches ■ minor pain of arthritis ■ backache ■ the common cold ■ toothache ■ the common cold ■ premenstrual and menstrual cramps temporarily reduces fever Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ hives ■ rash If a skin reaction occurs, stop use and seek medical help right away. Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are allergic to acetaminophen or any of the inactive ingredients in this product Ask a doctor before use if you have ■ liver disease (continued on opposite panel) Retain carton for complete product information	MEDI-FIRST Plus Extra Strength Non-Aspirin Acetaminophen 500 mg Pull To Open THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN. Pain Reliever/Fever Reducer Compare active ingredient to: Extra Strength Tylenol® Registered Trademark of McNeil Consumer Products Tamper Evident Unit Dose Packets	Drug Facts (continued) Ask a doctor or pharmacist before use if ■ you are taking the blood thinning drug warfarin Stop use and ask a doctor if ■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. Directions Do not use more than directed (see overdose warning) Adults and children (12 years and over) ■ take 2 tablets every 6 hours while symptoms last ■ do not take more than 6 tablets in 24 hours, unless directed by a doctor ■ do not use for more than 10 days unless directed by a doctor Children under 12 years: Ask a doctor Other information ■ store at room temperature 59°-86°F (15°-30°C) ■ tamper-evident sealed packets ■ do not use any opened or torn packets Questions or comments? 1-800-634-7680	Recorder #95633 4768290433 Manufactured for Med First Plus Fort Myers, FL 33967 USA 1-800-634-7680 www.mediqueproducts.com
---	--	---	---

Drug Facts Active ingredient (in each tablet) Acetaminophen 500 mg Pain reliever/fever reducer Uses temporarily relieves minor aches and pains due to: ■ headache ■ muscular aches ■ minor pain of arthritis ■ backache ■ the common cold ■ toothache ■ the common cold ■ premenstrual and menstrual cramps temporarily reduces fever Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ hives ■ rash If a skin reaction occurs, stop use and seek medical help right away. Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are allergic to acetaminophen or any of the inactive ingredients in this product Ask a doctor before use if you have ■ liver disease (continued on opposite panel) Retain carton for complete product information	MEDI-FIRST® Extra Strength Non-Aspirin Pain Reliever/Fever Reducer Aches, Fever • Acetaminophen 500 mg Pull To Open Compare active ingredient to: Extra Strength Tylenol® Registered Trademark of McNeil Consumer Products Tamper Evident Unit Dose Packets THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN	Drug Facts (continued) Ask a doctor or pharmacist before use if ■ you are taking the blood thinning drug warfarin Stop use and ask a doctor if ■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. Directions Do not use more than directed (see overdose warning) Adults and children (12 years and over) ■ take 2 tablets every 6 hours while symptoms last ■ do not take more than 6 tablets in 24 hours, unless directed by a doctor ■ do not use for more than 10 days unless directed by a doctor Children under 12 years: Ask a doctor Other information ■ store at room temperature 59°-86°F (15°-30°C) ■ tamper-evident sealed packets ■ do not use any opened or torn packets Questions or comments? 1-800-634-7680	MEDI-FIRST® Recorder #80433 Manufactured for Medique Products Fort Myers, FL 33967 1-800-634-7680 www.mediqueproducts.com
---	--	---	--

MEDI-FIRST PLUS EXTRA STRENGTH NON-ASPIRIN

acetaminophen tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:47682-274
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STARCH, CORN (UNII: O8232NY3SJ)	
POVIDONE K30 (UNII: U725QWY32X)	

Product Characteristics

Color	white ((White to Off-White))	Score	no score
Shape	ROUND	Size	12mm
Flavor		Imprint Code	G552
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-274-33	50 in 1 BOX	03/28/2024	
1		2 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:47682-274-48	125 in 1 BOX	03/28/2024	
2		2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	03/28/2024	

MEDI-FIRST EXTRA STRENGTH NON-ASPIRIN

acetaminophen tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:47682-273
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

Inactive Ingredients

Ingredient Name	Strength
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE K30 (UNII: U725QWY32X)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

Product Characteristics

Color	white ((White to Off-White))	Score	no score
Shape	ROUND	Size	12mm
Flavor		Imprint Code	G552
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-273-13	250 in 1 BOX	03/28/2024	
1	NDC:47682-273-99	2 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:47682-273-33	50 in 1 BOX	03/28/2024	
2	NDC:47682-273-99	2 in 1 PACKET; Type 0: Not a Combination Product		
3	NDC:47682-273-48	125 in 1 BOX	03/28/2024	
3	NDC:47682-273-99	2 in 1 PACKET; Type 0: Not a Combination Product		
4	NDC:47682-273-50	25 in 1 BOX	03/28/2024	
4	NDC:47682-273-99	2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing	Application Number or Monograph	Marketing Start	Marketing End
------------------	--	------------------------	----------------------

Category	Citation	Date	Date
OTC Monograph Drug	M013	03/28/2024	

MEDIQUE EXTRA STRENGTH APAP

acetaminophen tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:47682-272
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

Inactive Ingredients

Ingredient Name	Strength
STEARIC ACID (UNII: 4ELV7Z65AP)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
STARCH, CORN (UNII: O8232NY3SJ)	
POVIDONE K30 (UNII: U725QWY32X)	

Product Characteristics

Color	white ((White to Off-White))	Score	no score
Shape	ROUND	Size	12mm
Flavor		Imprint Code	G552
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-272-13	250 in 1 BOX	03/28/2024	
1	NDC:47682-272-99	2 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:47682-272-33	50 in 1 BOX	03/28/2024	
2	NDC:47682-272-99	2 in 1 PACKET; Type 0: Not a Combination Product		
3	NDC:47682-272-48	125 in 1 BOX	03/28/2024	
3	NDC:47682-272-99	2 in 1 PACKET; Type 0: Not a Combination Product		
	NDC:47682-272			

4	NDC:47682-272-64	12 in 1 BOX	03/28/2024	
4	NDC:47682-272-99	2 in 1 PACKET; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013		03/28/2024	

Labeler - Unifirst First Aid Corporation (832947092)

Establishment			
Name	Address	ID/FEI	Business Operations
Prestige Packaging		080667761	pack(47682-272, 47682-273, 47682-274)

Establishment			
Name	Address	ID/FEI	Business Operations
Medique Products		086911794	pack(47682-272, 47682-273, 47682-274)

Revised: 3/2024

Unifirst First Aid Corporation