

MUCINEX FAST-MAX SEVERE CONGESTION AND COUGH MAXIMUM STRENGTH-
dextromethorphan hydrobromide, guaifenesin capsule, liquid filled
RB Health (US) LLC

Mucinex Fast-Max Severe Congestion & Cough

Drug Facts

Active ingredients (in each liquid gel)

Dextromethorphan HBr 10 mg
Guaifenesin 200 mg

Purposes

Cough suppressant
Expectorant

Keep out of reach of children.

In case of overdose, get medical help or contact a
Poison Control Center right away.

Dextromethorphan HBr 10 mg....Cough suppressant
Guaifenesin 200mg.....Expectorant

Uses

Uses:

■ helps loosen phlegm (mucus) and thin bronchial
secretions to rid the bronchial passageways of
bothersome mucus and make coughs more
productive

Temporarily relieves:

- cough due to minor throat and bronchial
irritation as may occur with the common cold or
inhaled irritants
- the intensity of coughing
- the impulse to cough to help you get to sleep

Do not use:

■ if you are now taking a prescription monoamine
oxidase inhibitor (MAOI) (certain drugs for
depression, psychiatric, or emotional conditions,
or Parkinson's disease), or for 2 weeks after
stopping the MAOI drug. If you do not know if your
prescription drug contains an MAOI, ask a doctor
or pharmacist before taking this product.

Ask a doctor before use if you have

- persistent or chronic cough such as occurs with
smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

When using this product do not use more than directed

Stop use and ask a doctor if

■ cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than 12 capsules in any 24-hour period
- adults and children 12 years of age and over: take 2 capsules every 4 hours
- children under 12 years of age: do not use
- store at 20-25°C (68-77°F)
- avoid excessive heat

Inactive ingredients

D&C yellow no.10, FD&C blue no. 1, gelatin, glycerin, hypromellose, lecithin, light mineral oil, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution, titanium dioxide

Questions? 1-866-MUCINEX (1-866-682-4639)

Fast Dissolving Liquid Gels!

NDC 72854-126-16

MAXIMUM STRENGTH

MUCINEX®

FAST-MAX®



MUCINEX FAST-MAX SEVERE CONGESTION AND COUGH MAXIMUM STRENGTH

dextromethorphan hydrobromide, guaifenesin capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:72854-153
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RT9KYH) (DEXTROMETHORPHAN - UNII: 7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg

Inactive Ingredients

Ingredient Name	Strength
POVIDONE (UNII: FZ989GH94E)	
WATER (UNII: 059QF0KO0R)	

GLYCERIN (UNII: PDC6A3C00X)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)	
LIGHT MINERAL OIL (UNII: N6K5787QVP)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
GELATIN (UNII: 2G86QN327L)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	

Product Characteristics

Color	green	Score	no score
Shape	OVAL	Size	16mm
Flavor		Imprint Code	AR08
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72854-153-16	2 in 1 CARTON	06/01/2024	
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/01/2024	

Labeler - RB Health (US) LLC (081049410)