MUCINEX FAST-MAX SEVERE CONGESTION AND COUGH MAXIMUM STRENGTH-
dextromethorphan hydrobromide, guaifenesin capsule, liquid filled
Reckitt Benckiser LLC

Mucinex Fast-Max Severe Congestion & Cough

Drug Facts

Active ingredients (in each liquid gel)	**Purposes**
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant

Keep out of reach of children.
In case of overdose, get medical help or contact a Poison Control Center right away.

Dextromethorphan HBr 10 mg....Cough suppressant
Guaifenesin 200mg.....Expectorant

Uses

Uses:

■ helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Temporarily relieves:
■ cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
■ the intensity of coughing
■ the impulse to cough to help you get to sleep

Do not use:
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have
■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
■ cough that occurs with too much phlegm (mucus)

When using this product do not use more than directed

Stop use and ask a doctor if
■ cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.
In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

■ do not take more than 12 capsules in any 24-hour period
■ adults and children 12 years of age and over: take 2 capsules every 4 hours
■ children under 12 years of age: do not use

■ store at 20-25°C (68-77°F)
■ avoid excessive heat

Inactive ingredients
D&C yellow no.10, FD&C blue no. 1, gelatin, glycerin, hypromellose, lecithin, light mineral oil, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution, titanium dioxide

Questions? 1-866-MUCINEX (1-866-682-4639)

Fast Dissolving Liquid Gels!

NDC 72854-153

MAXIMUM STRENGTH

MUCINEX®
FAST-MAX ®

MUCINEX FAST-MAX SEVERE CONGESTION AND COUGH MAXIMUM STRENGTH

dextromethorphan hydrobromide, guaifenesin capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	**Item Code (Source)**	NDC:72854-153
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg

Inactive Ingredients

Ingredient Name	Strength
POVIDONE (UNII: FZ989GH94E)	
WATER (UNII: 059QF0KO0R)	
GLYCERIN (UNII: PDC6A3C0OX)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WQ0SDW1A)	

FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)	
LIGHT MINERAL OIL (UNII: N6K5787QVP)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
GELATIN (UNII: 2G86QN327L)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	

Product Characteristics

Color	green	**Score**	no score
Shape	OVAL	**Size**	16mm
Flavor		**Imprint Code**	AR08
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72854-153-16	2 in 1 CARTON	06/01/2024	
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/01/2024	

Labeler - Reckitt Benckiser LLC (094405024)

Revised: 5/2026 Reckitt Benckiser LLC