

TUSSIN MULTI SYMPTOM COLD CF- dextromethorphan hbr, guaifenesin, phenylephrine hci liquid
Cardinal Health (Leader) 70000

Drug Facts

Active ingredients (in each 10 mL)

Dextromethorphan HBr 20 mg

Guaifenesin 200 mg

Phenylephrine HCl 10 mg

Purposes

Cough suppressant

Expectorant

Nasal decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold:
 - nasal congestion
 - cough due to minor throat and bronchial irritation

Warnings

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- diabetes
- high blood pressure
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

Ask a doctor or pharmacist before use if you are

taking any other oral nasal decongestant or stimulant.

**When using this product,
do not use more than directed.**

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL=milliliter
- this adult product is not intended for use in children under 12 years of age
- adults and children 12 years and over: 10 mL every 4 hours
- children under 12 years: do not use

Other information

- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

anhydrous citric acid, FD&C red #40, flavor, glycerin, lactic acid, menthol, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose

Questions or comments?

Call **1-877-753-3935 Monday-Friday 9AM-5PM EST**

Principal Display Panel

Adult

Tussin Multi-Symptom Cold CF

Dextromethorphan HBr

Guaifenesin

Phenylephrine HCl

Cough Suppressant

Expectorant

Nasal Decongestant

Regular Strength

Multi-Symptom Cold

Ages 12 Years and Over

Relieves:

Nasal Congestion

Cough

Mucus

Non-Drowsy

Alcohol Free

FL OZ (mL)

Dosing Cup Included

COMPARE TO ROBITUSSIN® MULTI-SYMPTOM COLD CF active ingredients*

*This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributors of Robitussin® Multi-Symptom Cold CF.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

DISTRIBUTED BY CARDINAL HEALTH

DUBLIN, OHIO 43017

www.myleader.com 1-800-200-6313

Package Label

Drug Facts (continued)

Inactive ingredients citric acid, FD&C red #40, flavor, glycerin, lactic acid, menthol, propylene glycol, purified water, sodium benzoate, sorbitol, sucralose

Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

*This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Robitussin® Multi-Symptom Cold CF.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org



Dosing Cup Included

LEADER™

LEADER™

NDC 70000-0133-2

Adult Tussin Multi-Symptom Cold CF

Dextromethorphan HBr
Guaifenesin
Phenylephrine HCl

Cough Suppressant
Expectorant
Nasal Decongestant

Regular Strength

Multi-Symptom Cold

Ages 12 Years and Over

Relieves:
Nasal Congestion
Cough
Mucus

Non-Drowsy
Alcohol Free

COMPARE TO
ROBITUSSIN®
MULTI-SYMPTOM
COLD CF
active ingredients*

100% Money
Back Guarantee

8 FL. OZ (237 mL)

© 2017 Cardinal Health. All Rights Reserved. CARDINAL HEALTH, the Cardinal Health LOGO, LEADER, and the Leader LOGO are trademarks or registered trademarks of Cardinal Health. All other marks are the property of their respective owners.



DISTRIBUTED BY CARDINAL HEALTH
DUBLIN, OHIO 43017

www.myleader.com 1-800-200-6313

Essential to Care® since 1979



All LEADER™ Brand Products Have A
100% Money Back Guarantee

Return to place of purchase if not satisfied.

Drug Facts

Active ingredients (in each 10 mL)

Dextromethorphan HBr 20 mg.....Cough suppressant
Guaifenesin 200 mg.....Expectorant
Phenylephrine HCl 10 mg.....Nasal decongestant

Purposes

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes
- temporarily relieves these symptoms occurring with a cold
 - nasal congestion
 - cough due to minor throat and bronchial irritation

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- diabetes
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

Ask a doctor or pharmacist before use if you are taking any other oral nasal decongestant or stimulant.

When using this product, do not use more than directed.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.


Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age
- adults and children 12 years and over: 10 mL every 4 hours
- children under 12 years: do not use

Other information

■ store between 20-25°C (68-77°F). Do not refrigerate.

CIN 5283676 REV. 2/17



0 96295 12940 3

PLD-4322C F0003775

Lot No.:

Exp. Date:

LEADER Adult Tussin Multi-Symptom Cold CF

TUSSIN MULTI SYMPTOM COLD CF

dextromethorphan hbr, guaifenesin, phenylephrine hci liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70000-0133
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 10 mL
GUAIFENESIN (UNII: 405W7451V0) (GUAIFENESIN - UNII:405W7451V0)	GUAIFENESIN	200 mg

GUAIFENESIN (UNII: 493W7431VQ) (GUAIFENESIN - UNII:493W7431VQ)	GUAIFENESIN	in 10 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 10 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0K00R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
LACTIC ACID (UNII: 33X04XA5AT)	
MENTHOL (UNII: L7T10EIP3A)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70000-0133-1	1 in 1 BOX	01/31/2017	12/31/2024
1		118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:70000-0133-2	1 in 1 BOX	01/31/2017	12/31/2024
2		237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	01/31/2017	12/31/2024

Labeler - Cardinal Health (Leader) 70000 (063997360)

Revised: 12/2023

Cardinal Health (Leader) 70000