

CONRX ALLERGY SINUS- acetaminophen, chlorpheniramine maleate, and phenylephrine hydrochloride tablet

Eagle Distributors, Inc

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

ConRx™ Allergy Sinus

Drug Facts

Active ingredients (in each tablet)	Purpose
Acetaminophen 325 mg	Pain reliever
Chlorpheniramine maleate 2 mg	Antihistamine
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves these symptoms of hay fever or other upper respiratory allergies:
 - headache
 - sinus congestion and pressure
 - nasal congestion
 - runny nose and sneezing
 - minor aches and pains
 - itching of the nose or throat
 - itchy, watery eyes
- helps decongest sinus openings and passages

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 tablets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes

- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- glaucoma

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- **do not exceed recommended dosage**
- excitability may occur, especially in children
- drowsiness may occur
- alcohol, sedatives and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over take 2 tablets every 4 hours, do not take more than 8 tablets in 24 hours, children under 12 years, consult a doctor

Other information

- avoid excessive heat and humidity
- store between 15-30°C (59-86°F)
- **do not use if pouch is torn or open**
- see side panel for lot number and expiration date

Inactive ingredients

colloidal silicon dioxide, sodium lauryl sulphate, edetaet disodium, dioctyl sodium sulphosuccinate, polyvinylpyrrolidone, gelatin, corn starch, magnesium stearate, microcrystalline cellulose, polyethylene glycol 6000, sodium benzoate, sodium starch glycolate, stearic acid, purified talc, titanium dioxide, yellow iron oxide.

Questions or comments?

1-800-570-8650

PRINCIPAL DISPLAY PANEL - 2 Caplets × 50 Pouch Box

See New Warnings Information & Directions

Compare to the Active Ingredients in

Tylenol® Allergy*
Multi-Symptom

ConRx™ Allergy Sinus

Sneezing - Runny Nose - Congestion

- **Acetaminophen**
Headache
- **Phenylephrine HCl**
Sinus Pressure/Nasal Congestion
- **Chlorpheniramine Maleate**
Watery Eyes/Runny Nose

TO OPEN

PUSH IN TAB AND PULL OUT

Compare to the Active Ingredients in

Tylenol® Allergy*
Multi-Symptom

50 Pouches of 2 Caplets Each

See New Warnings Information & Directions
Compare to the Active Ingredients in

Tylenol® Allergy*
Multi-Symptom

ConRx™ Allergy Sinus
Sneezing - Runny Nose - Congestion

- Acetaminophen Headache
- Phenylephrine HCl Sinus Pressure/Nasal Congestion
- Chlorpheniramine Maleate Watery Eyes/Runny Nose

See New Warnings Information & Directions
Compare to the Active Ingredients in

Tylenol® Allergy*
Multi-Symptom

ConRx™ Allergy Sinus
Sneezing - Runny Nose - Congestion

- Acetaminophen Headache
- Phenylephrine HCl Sinus Pressure/Nasal Congestion
- Chlorpheniramine Maleate Watery Eyes/Runny Nose

▲
TO OPEN
PUSH IN TAB AND PULL OUT

Compare to the Active Ingredients in
Tylenol® Allergy*
Multi-Symptom

50 Pouches of 2 Caplets Each



50 Pouches of 2 Caplets Each

- Acetaminophen Headache
- Phenylephrine HCl Sinus Pressure/Nasal Congestion
- Chlorpheniramine Maleate Watery Eyes/Runny Nose

ConRx™ Allergy Sinus
Multi-Symptom
Tylenol® Allergy*
Sneezing - Runny Nose - Congestion

Compare to the Active Ingredients in
See New Warnings Information & Directions

Drug Facts

Active ingredients (in each tablet)

Active ingredients (in each tablet)	Purpose
Acetaminophen 325 mg	Pain reliever
Chlorpheniramine maleate 2 mg	Antihistamine
Phenylephrine HCl 5 mg	Nasal decongestant

Uses ■ temporarily relieves these symptoms of hay fever or other upper respiratory allergies:
 ■ headache ■ sinus congestion and pressure ■ nasal congestion ■ runny nose and sneezing ■ minor aches and pains ■ itching of the nose or throat
 ■ itchy, watery eyes
 ■ helps decongest sinus openings and passages

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
 ■ more than 8 tablets in 24 hours, which is the maximum daily amount.

See New Warnings Information & Directions
Compare to the Active Ingredients in

Tylenol® Allergy*
Multi-Symptom

ConRx™ Allergy Sinus

<ul style="list-style-type: none"> with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product 					
<p>Do not use</p> <ul style="list-style-type: none"> with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. if you have ever had an allergic reaction to this product or any of its ingredients 					
<p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> liver disease heart disease high blood pressure thyroid disease diabetes trouble urinating due to an enlarged prostate gland a breathing problem such as emphysema or chronic bronchitis glaucoma 					
<p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none"> taking the blood thinning drug warfarin taking sedatives or tranquilizers 					
<p>When using this product</p> <ul style="list-style-type: none"> do not exceed recommended dosage excitability may occur, especially in children drowsiness may occur alcohol, sedatives and tranquilizers may increase drowsiness avoid alcoholic drinks be careful when driving a motor vehicle or operating machinery 					
<p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> nervousness, dizziness, or sleeplessness occur pain or nasal congestion gets worse or lasts more than 7 days fever gets worse or lasts more than 3 days redness or swelling is present new symptoms occur <p>These could be signs of a serious condition.</p>					
<p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.</p> <p>Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p>					
<p>Directions ■ do not take more than directed (see overdose warning)</p> <table border="1"> <tr> <td>adults and children 12 years and over</td> <td> <ul style="list-style-type: none"> take 2 tablets every 4 hours do not take more than 8 tablets in 24 hours </td> </tr> <tr> <td>children under 12 years</td> <td>consult a doctor</td> </tr> </table>		adults and children 12 years and over	<ul style="list-style-type: none"> take 2 tablets every 4 hours do not take more than 8 tablets in 24 hours 	children under 12 years	consult a doctor
adults and children 12 years and over	<ul style="list-style-type: none"> take 2 tablets every 4 hours do not take more than 8 tablets in 24 hours 				
children under 12 years	consult a doctor				
<p>Other information</p> <ul style="list-style-type: none"> avoid excessive heat and humidity store between 15-30°C (59-86°F) do not use if pouch is torn or open see side panel for lot number and expiration date 					
<p>Inactive ingredients colloidal silicon dioxide, sodium lauryl sulphate, edetate disodium, dioctyl sodium sulphosuccinate, polyvinylpyrrolidone, gelatin, corn starch, magnesium stearate, microcrystalline cellulose, polyethylene glycol 6000, sodium benzoate, sodium starch glycolate, stearic acid, purified talc, titanium dioxide, yellow iron oxide.</p>					
<p>Questions or comments? 1-800-570-8650</p>					

Sneezing = Runny Nose = Congestion

- Acetaminophen
- Phenylephrine HCl
- Chlorpheniramine Maleate

Headache
Sinus Pressure/Nasal Congestion
Watery Eyes/Runny Nose

DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN

*This product is not manufactured or distributed by: McNeil Consumer Speciality Pharmaceuticals Division of McNeil PPC, Inc. Eagle Distributors, Inc./ConRx™ does not own the Tylenol® Allergy Multi-Symptom trademark.

Product manufactured for:
Eagle Distributors, Inc.
Los Angeles, CA 90011

CONRX ALLERGY SINUS

acetaminophen, chlorpheniramine maleate, and phenylephrine hydrochloride tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68737-231
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Acetaminophen (UNII: 362O9ITL9D) (Acetaminophen - UNII:362O9ITL9D)	Acetaminophen	325 mg
Chlorpheniramine maleate (UNII: V1Q0O9OJ9Z) (Chlorpheniramine - UNII:3U6IO1965U)	Chlorpheniramine maleate	2 mg
Phenylephrine Hydrochloride (UNII: 04JA59TNSJ) (Phenylephrine - UNII:1WS297W6MV)	Phenylephrine Hydrochloride	5 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
DOCUSATE SODIUM (UNII: F05Q2T2JA0)	
POVIDONE K90 (UNII: RDH86HJV5Z)	
GELATIN (UNII: 2G86QN327L)	
STARCH, CORN (UNII: O8232NY3SJ)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	

POLYETHYLENE GLYCOL 6000 (UNII: 30IQX730WE)

SODIUM BENZOATE (UNII: OJ245FE5EU)

SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)

STEARIC ACID (UNII: 4ELV7Z65AP)

TALC (UNII: 7SEV7J4R1U)

TITANIUM DIOXIDE (UNII: 15FIX9V2JP)

FERRIC OXIDE YELLOW (UNII: EX438O2MRT)

Product Characteristics

Color	YELLOW	Score	2 pieces
Shape	OVAL	Size	18mm
Flavor		Imprint Code	CRX
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68737-231-15	50 in 1 BOX		
1		2 in 1 POUCH		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part341	02/15/2013	

Labeler - Eagle Distributors,Inc (929837425)

Revised: 2/2013

Eagle Distributors,Inc