

MEDIQUE ASPIRIN- medique aspirin tablet, film coated
HF Acquisition Co LLC, DBA HealthFirst

Aspirin (NSAID*) 325mg

*nonsteroidal anti-inflammatory drug

Pain Reliever / Fever Reducer

USES

Temporarily relieves minor aches and pains associated with: headache; muscular aches; minor arthritis pain; backache; common cold; toothache; menstrual cramps; temporarily reduces fever

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy Alert: Aspirin may cause a severe allergic reaction which may include: hives, skin reddening, facial swelling, rash, asthma (wheezing), blisters, shock. If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This contains an NSAID, which may cause severe stomach bleeding. The change is higher if you: are age 60 or older; have had stomach ulcers or bleeding problems; take a blood thinner (anticoagulant) or steroid drug; take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others), have 3 or more alcoholic drinks everyday while using this product, take more for a longer time than directed

DO NOT USE- ASPIRIN

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer; right before or after heart surgery; if you are taking a prescription drug for gout, diabetes or arthritis.

Ask a doctor before use if: stomach bleeding warning applies to you; you have a history of stomach problems such as heartburn; you have high blood pressure, heart disease, liver cirrhosis or kidney disease; you are taking a diuretic.

Ask a doctor or pharmacist before use if you are: under a doctor's care for any serious condition; taking any other drug.

When using this product: Take with food or milk if stomach upset occurs.

STOP USE AND ASK DOCTOR- ASPIRIN

Stop Use and ask a Doctor if:

You experience any of the following signs of stomach bleeding, you feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better, pain

gets worse or lasts more than 10 days, fever gets worse or lasts more than 3 days, you have difficulty swallowing, if ringing in the ears or loss of hearing occurs, redness or swelling is present in painful areas, or any new symptoms appear.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of Children. In case of overdose, get medical help or contact a poison control center right away.

DIRECTIONS

Do not use more than directed - the smallest effective dose should be used.

Drink a full glass of water with each dose.

Do not take longer than 10 days, unless directed by a doctor.

Adults and Children (12 years and older): take 1 or 2 tablets with water every 4 hours as needed. Do not take more than 12 tablets in 24 hours, or as directed by a doctor.

Children under 12 years: Do not give to children under 12 years of age.

Avoid excessive heat and humidity, do not use any open or torn packets.

Aspirin (NSAID*) 325mg

Do not use more than directed - the smallest effective dose should be used.

Drink a full glass of water with each dose.

Do not take longer than 10 days, unless directed by a doctor.

Adults and Children (12 years and older): take 1 or 2 tablets with water every 4 hours as needed. Do not take more than 12 tablets in 24 hours, or as directed by a doctor.

Children under 12 years: Do not give to children under 12 years of age.

Avoid excessive heat and humidity, do not use any open or torn packets.

Do not use more than directed - the smallest effective dose should be used.

Drink a full glass of water with each dose.

Do not take longer than 10 days, unless directed by a doctor.

Adults and Children (12 years and older): take 1 or 2 tablets with water every 4 hours as needed. Do not take more than 12 tablets in 24 hours, or as directed by a doctor.

Children under 12 years: Do not give to children under 12 years of age.

Avoid excessive heat and humidity, do not use any open or torn packets.

KEEP OUT OF REACH OF CHILDREN

**MINERAL OIL
CELLULOSE, MICROCRYSTALLINE
CROSCARMELLOSE SODIUM**

HYPROMELLOSES
TITANIUM DIOXIDE
STARCH, CORN



Información del medicamento

Ingredientes activos por cada comprimido		Peso
Aspirina (Ácido Acetilsalicílico)		325 mg
Medicamentos adyuvantes inactivos		Ver el prospecto

Efecto:

- Alivia los síntomas por inflamación y fiebre que se producen.
- Alivia el dolor.
- Reduce la fiebre.

Indicaciones:

- Alivia el dolor de cabeza, de dientes, de nervios, de resaca, de dolor de espalda, de dolor de estómago, de dolor de articulaciones, de dolor de cuello, de dolor de hombros, de dolor de manos, de dolor de pies, de dolor de espalda baja, de dolor de espalda alta, de dolor de cuello, de dolor de hombros, de dolor de manos, de dolor de pies, de dolor de espalda baja, de dolor de espalda alta.

Contraindicaciones:

- Alergia a la aspirina o a cualquier otro fármaco que contenga ácido acetilsalicílico.
- Alergia a los salicilatos.
- Alergia a otros antiinflamatorios no esteroideos (AINE).
- Alergia a cualquier otro fármaco que contenga ácido acetilsalicílico o salicilatos.
- Alergia a cualquier otro fármaco que contenga ácido acetilsalicílico o salicilatos.

Precauciones:

- Evitar el uso de alcohol durante el tratamiento.
- Evitar el uso de otros medicamentos que contengan ácido acetilsalicílico o salicilatos.
- Evitar el uso de otros medicamentos que contengan ácido acetilsalicílico o salicilatos.
- Evitar el uso de otros medicamentos que contengan ácido acetilsalicílico o salicilatos.

Advertencias:

- El uso de este medicamento puede causar problemas de salud graves, incluso la muerte.
- El uso de este medicamento puede causar problemas de salud graves, incluso la muerte.
- El uso de este medicamento puede causar problemas de salud graves, incluso la muerte.

Uso:

- Leer cuidadosamente el prospecto que acompaña a este medicamento.
- Tomar el medicamento exactamente como se indica en el prospecto.
- Tomar el medicamento exactamente como se indica en el prospecto.
- Tomar el medicamento exactamente como se indica en el prospecto.

(01)00351662163313
(17)230919
(21)351662141212
(10)HF1234TESTING



SEE MANUFACTURE'S INSERT
DISTRIBUTED BY HF ACQUISITIONS CO., LLC
MUKILTEO, WA 98275

MEDIQUE ASPIRIN

medique aspirin tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:51662-1633(NDC:47682-097)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients

Ingredient Name	Strength
MINERAL OIL (UNII: T5L8T28FGP)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	TCL;011
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51662-1633-1	2 in 1 PACKET; Type 0: Not a Combination Product	12/30/2008	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M004	12/30/2008	

Labeler - HF Acquisition Co LLC, DBA HealthFirst (045657305)

Registrant - HF Acquisition Co LLC, DBA HealthFirst (045657305)

Establishment

Name	Address	ID/FEI	Business Operations
HF Acquisition Co LLC, DBA HealthFirst		045657305	repack(51662-1633)

Revised: 3/2024

HF Acquisition Co LLC, DBA HealthFirst