

**MUCINEX CHILDRENS MIGHTY CHEWS COUGH- dextromethorphan hydrobromide tablet, chewable**  
**RB Health (US) LLC**

-----  
**Mucinex Children's Mighty Chews Cough**

**Drug Facts**

**Dextromethorphan HBr 10 mg**

Dextromethorphan HBr 10 mg.....Cough suppressant

**Uses**

**Uses**

■ temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold

**Warnings**

**Do not use**

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- a sodium-restricted diet
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

**When using this product**

- do not exceed recommended dosage

**Stop use and ask a doctor if**

■ cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache. These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away

**Directions**

**Directions**

- take every 4 hours or as directed by a doctor
  - chew thoroughly before swallowing
-

adults and children 12 years of age and over  
 children 6 to under 12 years of age  
 children under 6 years of age

chewable tablets every 4 hours, not to exceed 12 chewable tablets in any 24-hour period  
 1 chewable tablet every 4 hours, not to exceed 6 chewable tablets in any 24-hour period  
 do not use

### Other information

- each chewable tablet contains: potassium 5 mg and sodium 15 mg
- store in a cool dry place at 20-25°C (68-77°F)

### Inactive ingredients

anhydrous citric acid, corn syrup, FD&C red no. 40, flavors, glycerin, malic acid, maltodextrin, pectin, potassium sodium tartrate, purified water, sodium chloride, trisodium citrate (anhydrous), sodium polymetaphosphate, sucralose, sucrose

Questions? 1-866-MUCINEX (1-866-682-4639)

NDC 72854-061-16

Children's  
 Mucinex®

Cough Suppressant

Dextromethorphan HBr 5 mg

Age 6+

Mixed Berry Flavor

Chewable Tablets

**KEEP OUT OF REACH OF CHILDREN  
 USE UNDER ADULT SUPERVISION**

Patents: www.reckitt.com/patents  
 121423  
 3287750  
 ©2024 RB Health  
 Dist. by: RB Health (US)  
 Parsippany, NJ 07054-0224  
 Made in India

©2024 RB Health  
 Dist. by: RB Health (US)  
 Parsippany, NJ 07054-0224  
 Made in India

\*product shown not actual size

**PARENTS:**  
 Learn about how medicines abuse  
 www.StopMedicineAbuse.org

6382499788

**Mucinex®**  
 Children's  
**MIGHTY CHEWS**  
**COUGH**

**6+**  
 NDC 72854-081-16

**Drug Facts**

Active ingredient (in each chewable tablet)	Purpose
Dextromethorphan HBr 10 mg	Cough suppressant

**Uses**

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold

**Warnings**

**Do not use**

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug.

Tamper evident: Protected with a tamper evident seal. **PEEL CORNER TO READ COMPLETE DRUG FACTS AND INFORMATION**

Do not use if seal under cap is broken or missing.

**16 CHEWABLE TABLETS - MIXED BERRY FLAVOR**

<p><b>Drug Facts (continued)</b></p> <p>If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p> <p><b>Ask a doctor before use if you have</b></p> <ul style="list-style-type: none"> <li>a sodium-restricted diet</li> <li>persistent or chronic cough such as occurs with smoking, asthma, or emphysema</li> <li>cough that occurs with too much phlegm (mucus)</li> </ul> <p><b>When using this product</b></p> <ul style="list-style-type: none"> <li>do not exceed recommended dosage</li> </ul> <p><b>Stop use and ask a doctor if</b></p> <ul style="list-style-type: none"> <li>cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache. These could be signs of a serious condition.</li> </ul> <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p><b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away</p>	<p><b>Drug Facts (continued)</b></p> <p><b>Directions</b></p> <ul style="list-style-type: none"> <li>take every 4 hours or as directed by a doctor</li> <li>chew thoroughly before swallowing</li> </ul> <table border="1"> <thead> <tr> <th>adults and children 12 years of age and over</th> <th>2 chewable tablets every 4 hours, not to exceed 12 chewable tablets in any 24-hour period</th> </tr> </thead> <tbody> <tr> <td>children 6 to under 12 years of age</td> <td>1 chewable tablet every 4 hours, not to exceed 6 chewable tablets in any 24-hour period</td> </tr> <tr> <td>children under 6 years of age</td> <td>do not use</td> </tr> </tbody> </table> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>each chewable tablet contains: potassium 5 mg and sodium 15 mg</li> <li>store in a cool dry place at 20-25°C (68-77°F)</li> </ul> <p><b>Inactive ingredients</b></p> <p>anhydrous citric acid, corn syrup, FD&amp;C red no. 40, flavors, glycerin, malic acid, maltodextrin, pectin, potassium sodium tartrate, purified water, sodium</p>	adults and children 12 years of age and over	2 chewable tablets every 4 hours, not to exceed 12 chewable tablets in any 24-hour period	children 6 to under 12 years of age	1 chewable tablet every 4 hours, not to exceed 6 chewable tablets in any 24-hour period	children under 6 years of age	do not use	<p><b>Drug Facts (continued)</b></p> <p>chloride, trisodium citrate (anhydrous), sodium polymetaphosphate, sucralose, sucrose</p> <p><b>Questions? 1-866-MUCINEX (1-866-682-4639)</b></p>
adults and children 12 years of age and over	2 chewable tablets every 4 hours, not to exceed 12 chewable tablets in any 24-hour period							
children 6 to under 12 years of age	1 chewable tablet every 4 hours, not to exceed 6 chewable tablets in any 24-hour period							
children under 6 years of age	do not use							

# MUCINEX CHILDRENS MIGHTY CHEWS COUGH

dextromethorphan hydrobromide tablet, chewable

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:72854-081
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>CORN SYRUP</b> (UNII: 9G5L16BK6N)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>PECTIN</b> (UNII: 89NA02M4RX)	
<b>POTASSIUM SODIUM TARTRATE</b> (UNII: QH257BPV3J)	
<b>SUCROSE</b> (UNII: C151H8M554)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>ANHYDROUS TRISODIUM CITRATE</b> (UNII: RS7A450LGA)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>MALIC ACID</b> (UNII: 817L1N4CKP)	
<b>MALTODEXTRIN</b> (UNII: 7CVR7L4A2D)	
<b>SODIUM CHLORIDE</b> (UNII: 451W47IQ8X)	
<b>SODIUM POLYMETAPHOSPHATE</b> (UNII: P1BM4ZH95L)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	

## Product Characteristics

<b>Color</b>	red	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	234mm
<b>Flavor</b>	BERRY (Mixed Berry)	<b>Imprint Code</b>	M
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72854-081-16	16 in 1 BOTTLE; Type 0: Not a Combination Product	06/03/2024	

## Marketing Information

Marketing	Application Number or Monograph	Marketing Start	Marketing End
-----------	---------------------------------	-----------------	---------------

Category	Citation	Date	Date
OTC Monograph Drug	M012	06/03/2024	

**Labeler** - RB Health (US) LLC (081049410)

Revised: 4/2024

RB Health (US) LLC