

RUGBY ASPIRIN- aspirin tablet, chewable

Rugby Laboratories

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Rugby Laboratories Aspirin Drug Facts

Active ingredient (in each chewable tablet)

Aspirin 81 mg (NSAID*)

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

for the temporary relief of minor aches and pains

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - pain gets worse or lasts for more than 10 days
 - fever gets worse or lasts for more than 3 days
 - redness or swelling is present
 - new symptoms occur
 - ringing in the ears or loss of hearing occurs

These could be signs of a serious condition

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 chewable tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor

- children under 12 years: consult a doctor

Other information

- store at 20-25°C (68-77°F)

Inactive ingredients

corn starch, dextrose excipient, FD&C yellow no. 6 aluminum lake, flavors, saccharin sodium

Questions or comments?

1-800-645-2158

Principal Display Panel

Rugby®

COMPARE TO ACTIVE INGREDIENT IN BAYER® CHEWABLE LOW DOSE ASPIRIN

Chewable Orange Flavor Aspirin 81 mg

Pain Reliever (NSAID)

actual size

Low Dose Aspirin Regimen

Talk to your doctor or other healthcare provider before using this product for your heart.

36 TABLETS

SAVE CARTON FOR COMPLETE DRUG FACTS

SEE NEW WARNING

46768 8T C3



DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts

Active ingredient (in each chewable tablet)
Aspirin 81 mg (NSAID)¹,..... Pain reliever
¹nonsteroidal anti-inflammatory drug

Purpose
Pain reliever

Uses for the temporary relief of minor aches and pains

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:
 ■ hives ■ facial swelling
 ■ asthma (wheezing) ■ shock
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you
 ■ are age 60 or older
 ■ have had stomach ulcers or bleeding problems
 ■ take a blood thinning (anticoagulant) or steroid drug

Drug Facts (continued)

■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 ■ have 3 or more alcoholic drinks every day while using this product
 ■ take more or for a longer time than directed

Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer
 ■ for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor

Ask a doctor before use if
 ■ stomach bleeding warning applies to you
 ■ you have a history of stomach problems, such as heartburn
 ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
 ■ you are taking a diuretic ■ you have asthma

Ask a doctor or pharmacist before use if
 ■ you are taking a prescription drug for
 ■ gout ■ diabetes ■ arthritis
Stop use and ask a doctor if
 ■ an allergic reaction occurs. Seek medical help right away.
 ■ you experience any of the following signs of stomach bleeding:
 ■ feel faint ■ vomit blood
 ■ have bloody or black stools
 ■ have stomach pain that does not get better

Drug Facts (continued)

■ pain gets worse or lasts more than 10 days
 ■ fever gets worse or lasts more than 3 days
 ■ redness or swelling is present
 ■ new symptoms occur
 ■ ringing in the ears or a loss of hearing occurs
 ■ These could be signs of a serious condition

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions
 ■ drink a full glass of water with each dose
 ■ adults and children 12 years and over: take 4 to 8 chewable tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
 ■ children under 12 years: consult a doctor

Other information
 ■ store at 20-25°C (68-77°F)

Drug Facts (continue d
Inactive ingredients corn starch, dextrose excipient, FD&C yellow no. 6 aluminum lake, flavors, saccharin sodium
Questions or comments?
 1-800-645-2158
 *This product is not manufactured or distributed by Bayer Aktiengesellschaft, owner of the registered trademark Bayer®.
 Distributed by: RUGBY® LABORATORIES
 Indianapolis, IN 46268
 www.rugbylaboratories.com
 REV. 02/22 R-05

Rugby® NDC 0536-1008-36

Chewable Orange Flavor Aspirin

81 mg Pain Reliever (NSAID)

Low Dose Aspirin Regimen
 Talk to your doctor or other healthcare provider before using this product for your heart.



36 TABLETS

SAVE CARTON FOR COMPLETE DRUG FACTS

SEE NEW WARNING

RUGBY ASPIRIN
 aspirin tablet, chewable

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0536-1008
---------------------	----------------	---------------------------	---------------

Route of Administration	ORAL			
Active Ingredient/Active Moiety				
	Ingredient Name	Basis of Strength	Strength	
	ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg	
Inactive Ingredients				
	Ingredient Name	Strength		
	SACCHARIN SODIUM (UNII: SB8ZUX40TY)			
	STARCH, CORN (UNII: O8232NY3S)			
	FD&C YELLOW NO. 6 (UNII: H77VEI93A8)			
Product Characteristics				
Color	ORANGE	Score	no score	
Shape	ROUND	Size	8mm	
Flavor	ORANGE	Imprint Code	L467	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0536-1008-36	1 in 1 CARTON	03/21/2014	
1		36 in 1 BOTTLE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph not final	part343	03/21/2014		

Labeler - Rugby Laboratories (079246066)

Revised: 2/2023

Rugby Laboratories