

**CHILDRENS PLUS FLU- acetaminophen, chlorpheniramine maleate, dextromethorphan hydrobromide, phenylephrine hci suspension
Dolgenercorp, Inc. (DOLLAR GENERAL & REXALL)**

Drug Facts

Active ingredients (in each 5 mL)

Acetaminophen 160 mg

Chlorpheniramine maleate 1 mg

Dextromethorphan HBr 5 mg

Phenylephrine HCl 2.5 mg

Purposes

Pain reliever/fever reducer

Antihistamine

Cough suppressant

Nasal decongestant

Uses

- temporarily relieves the following cold/flu symptoms:
 - minor aches and pains
 - sore throat
 - cough
 - sneezing and runny nose
 - headache
 - stuffy nose
- temporarily reduces fever

Warnings

Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes:

- more than 5 doses in 24 hours, which is the maximum daily amount

- with other drugs containing acetaminophen

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- to make a child sleepy
- with any other drug containing acetaminophen (Prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if your child has

- liver disease
- glaucoma
- thyroid disease
- diabetes
- heart disease
- high blood pressure
- a breathing problem such as chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with asthma

Ask a doctor or pharmacist before use if the child is taking

- sedatives or tranquilizers
- the blood thinning drug warfarin

When using this product

- **do not exceed recommended dose (see overdose warning)**
- excitability may occur, especially in children
- marked drowsiness may occur
- sedatives and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- new symptoms occur
- redness or swelling is present
- pain, nasal congestion or cough gets worse or last more than 5 days
- fever gets worse or last more than 3 days
- nervousness, dizziness or sleeplessness occurs
- cough comes back or occurs with rash or headache that lasts,

These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) could cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- **this product does not contain directions or complete warnings for adult use.**
- mL = milliliter
- shake well before using
- measure only with dosing cup provided
- do not use with any other dosing device
- keep dosing cup with product
- do not give more than 5 times in 24 hours
- if needed, repeat dose every 4 hours while symptoms last
- find the right dose on chart below. If possible, use weight to dose: otherwise us

Weight (lb)	Age (yr)	Dose (mL)
48-95	6-11	10 mL
36-47	4-5	do not use unless directed by a doctor
under 36	under 4	do not use

Other Information

- store at 20-25°C (68-77F) .Do not refrigerate

Inactive ingredients

carboxymethylcellulose sodium, citric acid, D&C red #33, FD&C red #40, flavors, glycerin, microcrystalline cellulose, purified water, sodium benzoate, sorbitol, sucrose xanthan gum

Questions or comments?

Call **1-888-309-9030**

Principal Display Panel

For ages 6 to 11 years

Children's

Plus Flu

Acetaminophen / Pain reliever

Chlorpheniramine maleate

Dextromethorphan HBr

Phenylephrine HCl

Oral Suspension

Pain reliever/fever reducer

Antihistamine

Cough suppressant

Nasal decongestant

Relieves coughs, runny nose, sneezing, fever, sore throat, stuffy nose

Grape Flavor

FL OZ (mL)

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

DISTRIBUTED BY OLD EAST MAIN CO.

100 MISSION RIDGE,

GOODLETTSVILLE, TN 37072

Package Label

Drug Facts (continued)

- nervousness, dizziness or sleeplessness occurs
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use
- mL = milliliter
- shake well before using
- do not give more than 5 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- if needed, repeat dose every 4 hours while symptoms last
- find the right dose on chart below. If possible, use weight to dose; otherwise, use age.

Weight (lb)	Age (yr)	Dose (mL)
48-95	6-11	10 mL
36-47	4-5	do not use unless directed by a doctor
under 36	under 4	do not use

Other information

- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients carboxymethylcellulose sodium, citric acid, D&C red #33, FD&C blue #1, FD&C red #40, flavor, glycerin, microcrystalline cellulose, purified water, sodium benzoate, sorbitol, sucrose, xanthan gum

Questions or comments?
Call 1-888-309-9030

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

DISTRIBUTED BY OLD EAST MAIN CO.
100 MISSION RIDGE,
GOODLETTSVILLE, TN 37072

100%
Satisfaction
Guaranteed!
(888) 309-9030

PARENTS:
Learn about how medicine works
www.StopMedicineAbuse.org

DG™ health

For Ages 6 to 11 Years
Children's Plus Flu

Acetaminophen
Chlorpheniramine maleate
Dextromethorphan HBr
Phenylephrine HCl

Oral Suspension
Pain Reliever/Fever Reducer
Antihistamine
Cough Suppressant
Nasal Decongestant



4 FL OZ (118 mL)

Grape Flavor

DG™ health

For Ages 6 to 11 Years
Children's Plus Flu

Acetaminophen
Chlorpheniramine maleate
Dextromethorphan HBr
Phenylephrine HCl

Oral Suspension
Pain Reliever/Fever Reducer
Antihistamine
Cough Suppressant
Nasal Decongestant



4 FL OZ (118 mL)

Grape Flavor

Drug Facts

Active ingredients (in each 5 mL)

Active ingredients (in each 5 mL)	Purposes
Acetaminophen 160 mg	Pain reliever/fever reducer
Chlorpheniramine maleate 1 mg	Antihistamine
Dextromethorphan HBr 5 mg	Cough suppressant
Phenylephrine HCl 2.5 mg	Nasal decongestant

Uses

- temporarily relieves the following cold/flu symptoms
- minor aches and pains
- sore throat
- cough
- sneezing and runny nose
- headache
- nasal congestion
- stuffy nose
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes: more than 5 doses (10 mL each) in 24 hours, which is the maximum daily amount

- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening, blisters, rash. If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- to make a child sleep
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if your child has

- liver disease
- glaucoma
- thyroid disease
- diabetes
- heart disease
- high blood pressure
- a breathing problem such as chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with asthma

Ask a doctor or pharmacist before use if your child is taking

- the blood thinning drug warfarin
- sedatives or tranquilizers

When using this product

- do not exceed recommended dose (see Overdose warning)
- excitability may occur, especially in children
- marked drowsiness may occur
- sedatives and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- new symptoms occur
- redness or swelling is present
- pain, nasal congestion or cough gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days

A0140



PLD-J250C FC007481

Lot No.:

Exp. Date:

DOLLAR GENERAL HEALTH Children's Plus Flu

CHILDRENS PLUS FLU

acetaminophen, chlorpheniramine maleate, dextromethorphan hydrobromide, phenylephrine hci suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55910-031
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	1 mg in 5 mL

DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	5 mg in 5 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	2.5 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
GLYCERIN (UNII: PDC6A3C0OX)	
WATER (UNII: 059QF0KO0R)	
D&C RED NO. 33 (UNII: 9DBA0SBB0L)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
XANTHAN GUM (UNII: TTV12P4NEE)	
SUCROSE (UNII: C151H8M554)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55910-031-04	1 in 1 BOX	05/31/2015	
1		118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	05/31/2015	

Labeler - Dolgencorp, Inc. (DOLLAR GENERAL & REXALL) (068331990)