

MUCINEX NASAL FULL FORCE- oxymetazoline hydrochloride solution
Reckitt Benckiser, Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Mucinex®
NASAL SPRAY
FULL FORCE

Drug Facts

Active ingredient

Oxymetazoline hydrochloride 0.05%

Purpose

Nasal decongestant

Uses

- temporarily relieves nasal congestion due to:
 - a cold
 - hay fever
 - upper respiratory allergies
- promotes nasal and sinus drainage
- temporarily relieves sinus congestion and pressure
- helps clear nasal passages; shrinks swollen membranes

Warnings

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland

When using this product

- **do not use more than directed**
- do not use for more than 3 days. Use only as directed. Frequent or prolonged use may cause nasal congestion to recur or worsen.
- temporary discomfort such as burning, stinging, sneezing or an increase in nasal discharge may occur
- use of this container by more than one person may spread infection

Stop use and ask a doctor if symptoms persist

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- adults and children 6 to under 12 years of age (with adult supervision): 2 or 3 sprays in each nostril not more often than every 10 to 12 hours. Do not exceed 2 doses in any 24-hour period.
- children under 6 years of age: ask a doctor

Shake well before use. Before using the first time, remove the protective cap from the tip and prime metered pump by depressing firmly several times. To spray, hold bottle with thumb at base and nozzle between first and second fingers. Without tilting head, insert nozzle into nostril. Fully depress pump all the way down with a firm even stroke and sniff deeply. Wipe nozzle clean after use.

Other information

- store between 20-25°C (68-77°F)
- retain carton for future reference on full labeling

Inactive ingredients

benzalkonium chloride solution, camphor, edetate disodium, eucalyptol, glycine, menthol, polyethylene glycol, polysorbate 80, propylene glycol, purified water, sodium chloride, sodium hydroxide

Questions?

1-866-MUCINEX (1-866-682-4639)

You may also report side effects to this phone number.

Dist. by: Reckitt Benckiser Inc.
Parsippany, NJ 07054-0224

PRINCIPAL DISPLAY PANEL - 22 mL Bottle Carton

NDC 63824-123-75

Mucinex®

NASAL SPRAY

Oxymetazoline HCl 0.05%

Nasal Decongestant

FULL

FORCE™

CONCENTRATED VAPOR

TECHNOLOGY®

- **Powerful
mucus-moving
sensation**
- **Fast Relief
of sinus pressure
& nasal congestion**

12
HOUR

3/4 FL OZ (22 mL)

DO NOT USE IF PRINTED SEAL OVER CAP IS TORN OR MISSING

Mucinex[®]
NASAL SPRAY
Oxymetazoline HCl 0.05% / Nasal Decongestant
FULL FORCE[™]
CONCENTRATED VAPOR TECHNOLOGY[®]

NDC 63824-123-75

Mucinex[®]
NASAL SPRAY
Oxymetazoline HCl 0.05%
Nasal Decongestant
FULL FORCE[™]
CONCENTRATED VAPOR TECHNOLOGY[®]

Also Available:
Mucinex[®]
Moisture Smart³
with
MEMBRANE HYDRATION TECHNOLOGY[®]
soothes and provides fast, 12 HOUR relief!

For 12 HOUR congestion relief in a tablet, choose from the Mucinex[®] line of products to treat your symptoms:

Mucinex[®]
CHEST CONGESTION

Mucinex[®] DM
COUGH & CHEST CONGESTION

Mucinex[®] D
NASAL/SINUS CONGESTION BREAKS UP MUCUS

Mucinex[®] tablets use a patented, bi-layer technology to thin and loosen mucus for 12 hours.

Please visit our web site:
www.mucinex.com

Drug Facts
Active Ingredient Purpose
Oxymetazoline hydrochloride 0.05%...Nasal decongestant

Uses
temporarily relieves nasal congestion due to:
a cold hay fever upper respiratory allergies
promotes nasal and sinus drainage
temporarily relieves sinus congestion and pressure
helps clear nasal passages; shrinks swollen membranes

Warnings
Ask a doctor before use if you have
heart disease high blood pressure
thyroid disease diabetes
trouble urinating due to an enlarged prostate gland

When using this product
do not use more than directed
do not use for more than 3 days. Use only as directed. Frequent or prolonged use may cause nasal congestion to recur or worsen.
temporary discomfort such as burning, stinging, sneezing or an increase in nasal discharge may occur
use of this container by more than one person may spread infection

Stop use and ask a doctor if symptoms persist
if pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions
adults and children 6 to under 12 years of age (with adult supervision): 2 or 3 sprays in each nostril not more often than every 10 to 12 hours. Do not exceed 2 doses in any 24-hour period.
children under 6 years of age: ask a doctor
Shake well before use. Before using the first time, remove the protective cap from the tip and prime metered pump by depressing firmly several times. To spray, hold bottle with thumb at base and nozzle between first and second fingers. Without tilting head, insert nozzle into nostril. Fully depress pump all the way down with a firm even stroke and sniff deeply. Wipe nozzle clean after use.

Other Information
store between 20-25°C (68-77°F)
retain carton for future reference on full labeling

Drug Facts (continued)
Inactive ingredients
benzalkonium chloride solution, camphor, edetate disodium, eucalyptol, glycine, menthol, polyethylene glycol, polysorbate 80, propylene glycol, purified water, sodium chloride, sodium hydroxide

Questions?
1-866-MUCINEX (1-866-682-4639)
You may also report side effects to this phone number.

✓ Powerful mucus-moving sensation
✓ Fast Relief of sinus pressure & nasal congestion

12 HOUR

Reckitt Benckiser
Dist. by: Reckitt Benckiser Inc., Parsippany, NJ 07054-0224
©RBI 2010 062910 0282234

3/4 FL OZ (22 mL)

3-63824-83436-8

CM-5020C

MUCINEX NASAL FULL FORCE

oxymetazoline hydrochloride solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63824-123
Route of Administration	NASAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Oxymetazoline Hydrochloride (UNII: K89MJ0S5VY) (Oxymetazoline - UNII:8VLN5B44ZY)	Oxymetazoline Hydrochloride	0.5 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
benzalkonium chloride (UNII: F5UM2KM3W7)	
camphor (synthetic) (UNII: 5TJD82A1ET)	
edetate disodium (UNII: 7FLD91C86K)	
eucalyptol (UNII: RV6J6604TK)	
glycine (UNII: TE7660XO1C)	
menthol (UNII: L7T10EIP3A)	
polyethylene glycol (UNII: 3WJQ0SDW1A)	
polysorbate 80 (UNII: 6OZP39ZG8H)	
propylene glycol (UNII: 6DC9Q167V3)	
water (UNII: 059QF0KO0R)	
sodium chloride (UNII: 451W47IQ8X)	
sodium hydroxide (UNII: 55X04QC32I)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63824-123-75	1 in 1 CARTON		
1		22 mL in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part341	06/30/2010	

Labeler - Reckitt Benckiser, Inc. (094405024)