

**DAYTIME NIGHTTIME COLD AND FLU RELIEF MULTI-SYMP TOM-
acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine
hcl
P & L Development, LLC**

Drug Facts

Active ingredients in Daytime (in each softgel)

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Phenylephrine HCl 5 mg

Active ingredients in Nighttime (in each softgel)

Acetaminophen 325 mg

Dextromethorphan HBr 15 mg

Doxylamine succinate 6.25 mg

Purpose for Daytime

Pain reliever/fever reducer

Cough suppressant

Nasal decongestant

Purpose for Nighttime

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Uses

DAYTIME

- temporarily relieves common cold and flu symptoms
 - cough due to minor throat and bronchial irritation
 - nasal condition
 - headache
 - minor aches and pains
 - fever
 - sore throat

NIGHTTIME

- temporarily relieves common cold and flu symptoms
 - cough due to minor throat and bronchial irritation
 - sore throat
 - headache
 - minor aches and pains
 - fever
 - runny nose & sneezing

Warnings

DAYTIME NIGHTTIME

Liver warning: These products contain acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

DAYTIME NIGHTTIME

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

DAYTIME

- liver disease
- heart disease
- diabetes
- thyroid disease
- high blood pressure
- cough that occurs with too much phlegm (mucus)

- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occur with smoking, asthma, or emphysema

NIGHTTIME

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are

DAYTIME

- taking the blood thinning drug warfarin.

NIGHTTIME

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product,

DAYTIME

do not use more than directed.

NIGHTTIME

- **do not exceed recommended dosage**
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

DAYTIME

- nervousness, dizziness, or sleeplessness occur
- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- redness or swelling is present
- new symptoms occur
- fever gets worse or lasts more than 3 days
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

NIGHTTIME

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

DAYTIME NIGHTTIME

Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

DAYTIME

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in 24 hours
- swallow whole: do not crush, chew, or dissolve
- adults and children 12 years and over: take 2 softgels with water every 4 hours
- children under 12 years: do not use

NIGHT TIME

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in 24 hours
- swallow whole: do not crush, chew, or dissolve
- adult and children 12 years and over: take 2 softgels with water every 6 hours
- children under 12 years: do not use

Other information

- store between 15°-30°C (59°-86°F)
- avoid excessive heat

Inactive ingredients

Daytime FD&C red #40, FD&C yellow #6, gelatin, glycerin, lecithin, light mineral oil, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol white ink

Nighttime D&C yellow #10, FD&C blue #1, gelatin, glycerin, lecithin, light mineral oil, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink

Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal Display Panel

when using daytime and nighttime products, carefully read the labeling to ensure correct dosing.

DAYTIME

non-drowsy

multi-symptom

day time

cold & flu relief

Acetaminophen 325 mg

dextromethorphan HBr 10 mg

phenylephrine HCl 5 mg

pain reliever/fever reducer

cough suppressant

nasal decongestant

alcohol-free

antihistamine-free

softgels**

(**liquid-filled capsules)

NIGHTTIME

Compare to the active ingredients in Vicks® DayQuil® and NyQuil® Cold & Flu LiquiCap®†

multi-symptom

night time

cold & flu relief

Acetaminophen 325 mg

dextromethorphan HBr 15 mg

doxylamine succinate 6.25 mg

pain reliever/fever reducer

cough suppressant

antihistamine

alcohol-free

softgels

(**liquid-filled capsules)

*This product is not manufactured or distributed by The Procter & Gamble Company, Vicks® DayQuil® and NyQuil® LiquiCaps® are registered trademarks of The Procter and Gamble Company.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

Distributed by:

PL Developments

200 Hicks Street

Westbury, NY 11590

Product Label



when using daytime and nighttime products, carefully read the labeling to ensure correct dosing.

non-drowsy multi-symptom

day time cold & flu relief

Acetaminophen 325 mg
dextromethorphan HBr 10 mg
phenylephrine HCl 5 mg
 pain reliever/fever reducer
 cough suppressant
 nasal decongestant

alcohol-free
 antihistamine-free



actual size

16 softgels** (**liquid-filled capsules)

total 24 softgels



Compare to the active ingredients in Vicks® DayQuil® and NyQuil® Cold & Flu LiquiCaps®¹
 NDC 59726-901-24

multi-symptom

night time cold & flu relief

Acetaminophen 325 mg
dextromethorphan HBr 15 mg
doxylamine succinate 6.25 mg
 pain reliever/fever reducer
 cough suppressant
 antihistamine

alcohol-free



actual size

8 softgels** (**liquid-filled capsules)

PLD-A586B FC00766Z



3 59756 90124 7

Distributed by:
PL Developments
 200 Hicks Street
 Westbury, NY 11590



TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BULSTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

PARENTS:
 Learn about our medicine here
www.3p24hicksonline.org

Lot No.:
 Exp. Date:

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 ■ If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
Ask a doctor before use if you have ■ liver disease ■ heart disease ■ diabetes ■ thyroid disease ■ high blood pressure ■ cough that occurs with too much phlegm (mucus) ■ trouble urinating due to an enlarged prostate gland ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema
Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.
When using this product, do not exceed recommended dosage.
Stop use and ask a doctor if ■ nervousness, dizziness, or sleeplessness occur ■ pain, cough, or nasal congestion gets worse or lasts more than 7 days ■ redness or swelling is present ■ new symptoms occur ■ fever gets worse or lasts more than 3 days ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children. Overdose warning:** Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick notice any signs or symptoms.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 ■ If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
Ask a doctor before use if you have ■ liver disease ■ glaucoma ■ cough that occurs with too much phlegm (mucus) ■ a breathing problem or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis, or emphysema ■ trouble urinating due to an enlarged prostate gland ■ taking the blood thinning drug warfarin ■ taking sedatives or tranquilizers
Ask a doctor or pharmacist before use if you are ■ taking the blood thinning drug warfarin ■ taking sedatives or tranquilizers ■ alcohol, sedatives, and tranquilizers may increase drowsiness ■ marked drowsiness may occur ■ avoid alcoholic drinks ■ be careful when driving a motor vehicle or operating machinery
Stop use and ask a doctor if ■ pain or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children. Overdose warning:** Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick notice any signs or symptoms.

Drug Facts (continued)
Daytime Cold & Flu Softgel
Purposes Acetaminophen 325 mg Pain reliever/fever reducer
 Dextromethorphan HBr 10 mg Cough suppressant
 Phenylephrine HCl 5 mg Nasal decongestant
Uses ■ temporarily relieves common cold and flu symptoms: ■ cough due to minor throat and bronchial irritation ■ nasal congestion ■ headache ■ minor aches and pains ■ fever ■ sore throat
Warnings This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Drug Facts (continued)
Nighttime Cold & Flu Softgel
Purposes Acetaminophen 325 mg Pain reliever/fever reducer
 Doxylamine succinate 6.25 mg Cough suppressant
 Antihistamine
Uses ■ temporarily relieves common cold and flu symptoms: ■ cough due to minor throat and bronchial irritation ■ sore throat ■ headache ■ minor aches and pains ■ fever ■ runny nose & sneezing
Warnings This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

This product is not manufactured or distributed by The Procter & Gamble Company. Vicks®, DayQuil®, NyQuil®, and LiquiCaps® are registered trademarks of The Procter and Gamble Company.

Daytime Cold & Flu Softgel	Nighttime Cold & Flu Softgel
Drug Facts (continued)	Drug Facts (continued)
Directions <ul style="list-style-type: none"> ■ do not take more than directed (see Overdose warning) ■ do not take more than 4 doses in 24 hours ■ swallow whole; do not crush, chew, or dissolve ■ adults and children 12 years and over: take 2 softgels with water every 6 hours ■ children under 12 years: do not use 	Directions <ul style="list-style-type: none"> ■ do not take more than directed (see Overdose warning) ■ do not take more than 4 doses in 24 hours ■ swallow whole; do not crush, chew, or dissolve ■ adults and children 12 years and over: take 2 softgels with water every 6 hours ■ children under 12 years: do not use
Other information <ul style="list-style-type: none"> ■ store between 15-30°C (59-86°F) ■ avoid excessive heat 	Other information <ul style="list-style-type: none"> ■ store between 15-30°C (59-86°F) ■ avoid excessive heat
Inactive ingredients FD&C yellow #6, FD&C red #40, gelatin, glycerin, glycol, light mineral oil, mannitol, polyethylene glycol, polyethylene glycol, purified water, sorbitol, sorbitol, white ink	Inactive ingredients D&C yellow #10, FD&C blue #1, gelatin, glycerin, lactin, light mineral oil, mannitol, polyethylene glycol, polyethylene glycol, purified water, sorbitol, sorbitol, white ink
Questions or comments? Call 1-877-753-3935 Monday-Friday 9AM-5PM EST	Questions or comments? Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

READYinCASE Day Time Night Time Cold and Flu Relief

DAYTIME NIGHTTIME COLD AND FLU RELIEF MULTI-SYMPTOM

acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:59726-901
---------------------	----------------	---------------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:59726-901-24	1 in 1 KIT; Type 0: Not a Combination Product	03/26/2021	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 BLISTER PACK	8
Part 2	1 BLISTER PACK	16

Part 1 of 2

NIGHT TIME COLD AND FLU RELIEF MULTI SYMPTOM

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate capsule, liquid filled

Product Information

Item Code (Source)	NDC:59726-899
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
-----------------	-------------------	----------

ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
DOXYLAMINE SUCCINATE (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

Inactive Ingredients

Ingredient Name	Strength
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITAN (UNII: 6O92ICV9RU)	
SORBITOL (UNII: 506T60A25R)	
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)	
LIGHT MINERAL OIL (UNII: N6K5787QVP)	
MANNITOL (UNII: 3OWL53L36A)	

Product Characteristics

Color	green	Score	no score
Shape	OVAL	Size	21mm
Flavor		Imprint Code	PC10
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	03/26/2021	

Part 2 of 2

DAYTIME COLD AND FLU NON DROWSY

acetaminophen, dextromethorphan hbr, phenylephrine hcl capsule, liquid filled

Product Information

Item Code (Source)	NDC:59726-898
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)	
LIGHT MINERAL OIL (UNII: N6K5787QVP)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GELATIN (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITAN (UNII: 6O92ICV9RU)	
SORBITOL (UNII: 506T60A25R)	
MANNITOL (UNII: 3OWL53L36A)	

Product Characteristics

Color	orange	Score	no score
Shape	CAPSULE	Size	20mm
Flavor		Imprint Code	PC9
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		16 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing	Application Number or Monograph	Marketing Start	Marketing End
-----------	---------------------------------	-----------------	---------------

Category	Citation	Date	Date
OTC Monograph Drug	M012	03/26/2021	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	03/26/2021	

Labeler - P & L Development, LLC (800014821)

Revised: 4/2024

P & L Development, LLC