MUCINEX CHILDRENS MIGHTY CHEWS COUGH DAYTIME AND NIGHTTIME COMBO PACK- dextromethorphan hbr and doxylamine succinate
Reckitt Benckiser LLC

Mucinex Childrens Mighty Chews Cough Daytime and Nighttime Value Pack

Drug Facts

Mucinex Children's Mighty Chews Cough Daytime:

Active ingredient (in each chewable tablet)
Dextromethorphan HBr 10 mg

Mucinex Children's Mighty Chews Cough Nighttime:

Active ingredients (in each chewable tablet)
Dextromethorphan HBr 10 mg
Doxylamine succinate 6.25 mg

Dextromethorphan HBr 10 mg..... .....Cough suppressant
Doxylamine succinate 6.25 mg.....Antihistamine

Uses

Mucinex Children's Mighty Chews Cough Daytime:

temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold

Mucinex Children's Mighty Chews Cough Nighttime:

■ temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
■ controls cough to help you get to sleep
■ temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
■ runny nose
■ sneezing
■ itching of the nose or throat
■ itchy, watery eyes

Mucinex Children's Mighty Chews Cough Daytime:

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson’s disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have
■ a sodium-restricted diet

■ cough that occurs with too much phlegm (mucus)
■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema

When using this product do not exceed recommended dosage

Stop use and ask a doctor if cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache. These could be signs of a serious condition.

Mucinex Children's Mighty Chews Cough Nighttime:

Warnings

Do not use
■ to make a child sleepy
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson’s disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have
■ a sodium-restricted diet

■ glaucoma
■ trouble urinating due to an enlarged prostate gland
■ a breathing problem such as emphysema or chronic bronchitis
■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema
■ cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product
■ do not exceed recommended dosage

■ excitability may occur, especially in chidren

■ marked drowsiness may occur
■ alcohol, sedatives, and tranquilizers may increase drowsiness
■ avoid alcoholic drinks
■ use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use

.
Keep out of reach of children. In case of overdose, get medical help or

contact a Poison Control Center right away

Directions

Mucinex Children's Mighty Chews Cough Daytime+Mucinex Children's Mighty Chews Cough Nighttime::

■ take every 4 hours or as directed by a doctor
■ chew thoroughly before swallowing

■ do not swallow whole

adults and children 12 years of age and over	2 chewable tablets every 4 hours, not to exceed 12 chewable tablets in any 24-hour period
children 6 to under 12 years of age	1 chewable tablet every 4 hours, not to exceed 6 chewable tablets in any 24-hour period
children under 6 years of age	do not use

Other information
■ each chewable tablet contains: **potassium 5 mg and sodium 15 mg**
■ store in a cool dry place at 20-25°C (68-77°F)

Mucinex Children's Mighty Chews Cough Daytime:

Inactive ingredients anhydrous citric acid, corn syrup, FD&C red no. 40, flavor, glycerin, malic acid, maltodextrin, pectin, potassium sodium tartrate, purified water, sodium chloride, sodium polymetaphosphate,

sucralose, sucrose, trisodium citrate (anhydrous)

Mucinex Children's Mighty Chews Cough Nighttime:

Inactive ingredients anhydrous citric acid, corn syrup, FD&C blue no. 1, FD&C red no. 40, flavor, glycerin, malic acid, maltodextrin, pectin, potassium sodium tartrate, purified water, sodium chloride, sodium

polymetaphosphate, sucralose, sucrose, trisodium citrate (anhydrous)

Questions? 1-866-MUCINEX (1-866-682-4639)

Mucinex Children's Mighty Chews Cough Nighttime

Drug Facts

Active ingredient (in each chewable tablet) — **Purposes**

Dextromethorphan HBr 10 mg ..... Cough suppressant
Doxylamine succinate 6.25 mg ..... Antihistamine

Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- controls cough to help you get to sleep
- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
  - runny nose
  - sneezing
  - itching of the nose or throat
  - itchy, watery eyes

Warnings

Do not use

- to make a child sleepy
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- a sodium-restricted diet
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product

- do not exceed recommended dosage
- excitability may occur, especially in children

Mucinex Children's Mighty Chews Cough Daytime

Drug Facts

Active ingredient (in each chewable tablet) — **Purpose**

Dextromethorphan HBr 10 mg ..... Cough suppressant

Uses temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- a sodium-restricted diet
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

When using this product do not exceed recommended dosage

Stop use and ask a doctor if cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away

Directions

- take every 4 hours or as directed by a doctor
- chew thoroughly before swallowing
- do not swallow whole

adults and children 12 years of age and over	2 chewable tablets every 4 hours, not to exceed 12 chewable tablets in any 24-hour period
children 6 to under 12 years of age	1 chewable tablet every 4 hours, not to exceed 6 chewable tablets in any 24-hour period
children under 6 years of age	do not use

MUCINEX CHILDRENS MIGHTY CHEWS COUGH DAYTIME AND NIGHTTIME COMBO PACK

dextromethorphan hbr and doxylamine succinate kit

Product Information			
Product Type	HUMAN OTC DRUG	**Item Code (Source)**	NDC:72854-083

Packaging				
#	**Item Code**	**Package Description**	**Marketing Start Date**	**Marketing End Date**
1	NDC:72854-083-32	1 in 1 CARTON	06/03/2024	

Quantity of Parts		
Part #	**Package Quantity**	**Total Product Quantity**
Part 1	1 BOTTLE	16
Part 2	1 BOTTLE	16

Part 1 of 2

MUCINEX CHILDRENS MIGHTY CHEWS COUGH

dextromethorphan hydrobromide tablet, chewable

Product Information	
Item Code (Source)	NDC:72854-081
Route of Administration	ORAL

Active Ingredient/Active Moiety		
Ingredient Name	**Basis of Strength**	**Strength**
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg

Inactive Ingredients	
Ingredient Name	**Strength**
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SUCROSE (UNII: C151H8M554)	
SODIUM POLYMETAPHOSPHATE (UNII: P1BM4ZH95L)	
WATER (UNII: 059QF0KO0R)	
ANHYDROUS TRISODIUM CITRATE (UNII: RS7A450LGA)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	

MALTODEXTRIN (UNII: 7CVR7L4A2D)	
POTASSIUM SODIUM TARTRATE (UNII: QH257BPV3J)	
PECTIN (UNII: 89NA02M4RX)	
CORN SYRUP (UNII: 9G5L16BK6N)	
GLYCERIN (UNII: PDC6A3C0OX)	
MALIC ACID (UNII: 817L1N4CKP)	

Product Characteristics

Color	red	**Score**	no score
Shape	ROUND	**Size**	23mm
Flavor	BERRY (Mixed Berry)	**Imprint Code**	M
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72854-081-16	16 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/03/2024	

Part 2 of 2

MUCINEX CHILDRENS MIGHTY CHEWS COUGH NIGHTTIME

dextromethorphan hbr and doxylamine succinate tablet, chewable

Product Information

Item Code (Source)	NDC:72854-082
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C RED NO. 40 (UNII: WZB9127XOA)	
PECTIN (UNII: 89NA02M4RX)	
SODIUM POLYMETAPHOSPHATE (UNII: P1BM4ZH95L)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
WATER (UNII: 059QF0KO0R)	
POTASSIUM SODIUM TARTRATE (UNII: QH257BPV3J)	
ANHYDROUS TRISODIUM CITRATE (UNII: RS7A450LGA)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SUCROSE (UNII: C151H8M554)	
CORN SYRUP (UNII: 9G5L16BK6N)	
GLYCERIN (UNII: PDC6A3C0OX)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
MALIC ACID (UNII: 817L1N4CKP)	

Product Characteristics

Color	purple	**Score**	no score
Shape	ROUND	**Size**	23mm
Flavor	BERRY (Mixed Berry)	**Imprint Code**	M
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72854-082-16	16 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/03/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/03/2024	

Labeler - Reckitt Benckiser LLC (094405024)

Revised: 5/2026 Reckitt Benckiser LLC