

PHYSICIANS EZ USE B-12 COMPLIANCE - physicians ez use b-12 compliance
Asclemed USA, Inc.

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Drug Facts

Active ingredient

Isopropyl Alcohol, 70% by volume

Antiseptic

Use

For preparation of the skin prior to injection

Warnings

For external use only. Flammable, keep away from fire or flame.

Do not use with electrocautery procedures, or in/near eyes.

Stop use if irritation or redness develops.

If irritating condition persists for more than 72 hours, consult a physician.

Keep out of reach of children. If swallowed, seek medical attention and/or contact a Poison Control Center immediately.

Directions

Prepare site by wiping vigorously

Inactive ingredient

Purified water


DESCRIPTION

Cyanocobalamin Injection, USP is a sterile solution of cyanocobalamin for intramuscular or subcutaneous use.

Each mL contains 1000 mcg cyanocobalamin; sodium chloride 0.9%; benzyl alcohol 1.5%; Water for Injection q.s. Hydrochloric acid and/or sodium hydroxide for pH adjustment if necessary (4.5-7.0).

Cyanocobalamin appears as dark, red crystals or as an amorphous or crystalline, red powder. It is very hygroscopic in the anhydrous form, and sparingly soluble in water (1:80). It is stable to autoclaving for short periods at 121°C. The Vitamin B₁₂ coenzymes are very unstable in light.

The chemical name is 5,6-dimethyl-benzimidazolyl cyanocobamide. The cobalt content is 4.34%. The structural formula is represented below:



$C_{63}H_{88}CoN_{14}O_{14}P$

1355.38

CLINICAL PHARMACOLOGY

Vitamin B₁₂ is essential to growth, cell reproduction, hematopoiesis, nucleoprotein and myelin synthesis.

Cyanocobalamin is quantitatively and rapidly absorbed from intramuscular and subcutaneous sites of injection; the plasma level of the compound reaches its peak within one hour after intramuscular injection. Absorbed Vitamin B₁₂ is transported via specific B₁₂ binding proteins, transcobalamin I and II to the various tissues. The liver is the main organ for Vitamin B₁₂ storage.

Within 48 hours after injection of 100 or 1000 mcg of Vitamin B₁₂, 50 to 98% of the injected dose may appear in the urine. The major portion is excreted within the first eight hours. Intravenous administration results in even more rapid excretion with little opportunity for liver storage.

Gastrointestinal absorption of Vitamin B₁₂ depends on the presence of sufficient intrinsic factor and calcium ions. Intrinsic factor deficiency causes pernicious anemia, which may be associated with subacute combined degeneration of the spinal cord. Prompt parenteral administration of Vitamin B₁₂ prevents progression of neurologic damage.

The average diet supplies about 5 to 15 mcg/day of Vitamin B₁₂ in a protein-bound form that is available for absorption after normal digestion. Vitamin B₁₂ is not present in foods of plant origin, but is abundant in foods of animal origin. In people with normal absorption, deficiencies have been reported only in strict vegetarians who consume no products of animal origin (including no milk products or eggs).

Vitamin B₁₂ is bound to intrinsic factor during transit through the stomach; separation occurs in the terminal ileum in the presence of calcium, and Vitamin B₁₂ enters the mucosal cell for absorption. It is then transported by the transcobalamin binding proteins. A small amount (approximately 1% of the total amount ingested) is absorbed by simple diffusion, but this mechanism is adequate only with very large doses. Oral absorption is considered too undependable to rely on in patients with pernicious anemia or other conditions resulting in malabsorption of Vitamin B₁₂.

Cyanocobalamin is the most widely used form of Vitamin B₁₂, and has hematopoietic activity apparently identical to that of the antianemia factor in purified liver extract. Hydroxocobalamin is equally as effective as cyanocobalamin, and they share the cobalamin molecular structure.

INDICATIONS AND USAGE

Cyanocobalamin is indicated for Vitamin B₁₂ deficiencies due to malabsorption which may be associated with the following conditions:

Addisonian (pernicious) anemia

Gastrointestinal pathology, dysfunction, or surgery, including gluten enteropathy or sprue, small bowel bacterial overgrowth, total or partial gastrectomy

Fish tapeworm infestation

Malignancy of pancreas or bowel

Folic acid deficiency

It may be possible to treat the underlying disease by surgical correction of anatomic lesions leading to small bowel bacterial overgrowth, expulsion of fish tapeworm, discontinuation of drugs leading to vitamin malabsorption (see ***Drug/Laboratory Test Interactions***), use of a gluten-free diet in nontropical sprue, or administration of antibiotics in tropical sprue. Such measures remove the need for long-term administration of cyanocobalamin.

Requirements of Vitamin B₁₂ in excess of normal (due to pregnancy, thyrotoxicosis, hemolytic anemia, hemorrhage, malignancy, hepatic and renal disease) can usually be met with oral supplementation.

Cyanocobalamin injection is also suitable for the Vitamin B₁₂ absorption test (Schilling test).

CONTRAINDICATIONS

Sensitivity to cobalt and/or Vitamin B₁₂ is a contraindication.

WARNINGS

WARNING: This product contains aluminum that may be toxic. Aluminum may reach toxic levels with prolonged parenteral administration if kidney function is impaired. Premature neonates are particularly at risk because their kidneys are immature, and they require large amounts of calcium and phosphate solutions, which contain aluminum.

Research indicates that patients with impaired kidney function, including premature neonates, who receive parenteral levels of aluminum at greater than 4 to 5 mcg/kg/day accumulate aluminum at levels associated with central nervous system and bone toxicity. Tissue loading may occur at even lower rates of administration.

Patients with early Leber's disease (hereditary optic nerve atrophy) who were treated with cyanocobalamin suffered severe and swift optic atrophy.

Hypokalemia and sudden death may occur in severe megaloblastic anemia which is treated intensely.

Anaphylactic shock and death have been reported after parenteral Vitamin B₁₂ administration. An intradermal test dose is recommended before cyanocobalamin injection is administered to patients suspected of being sensitive to this drug.

This product contains benzyl alcohol. Benzyl alcohol has been reported to be associated with a fatal "Gaspig Syndrome" in premature infants.

PRECAUTIONS

General

Vitamin B₁₂ deficiency that is allowed to progress for longer than three months may produce permanent degenerative lesions of the spinal cord. Doses of folic acid greater than 0.1 mg/day may result in hematologic remission in patients with Vitamin B₁₂ deficiency. Neurologic manifestations will not be prevented with folic acid, and if not treated with Vitamin B₁₂, irreversible damage will result.

Doses of cyanocobalamin exceeding 10 mcg daily may produce hematologic response in patients with

folate deficiency. Indiscriminate administration may mask the true diagnosis.

Information for Patients

Patients with pernicious anemia should be instructed that they will require monthly injections of Vitamin B₁₂ for the remainder of their lives. Failure to do so will result in return of the anemia and in development of incapacitating and irreversible damage to the nerves of the spinal cord. Also, patients should be warned about the danger of taking folic acid in place of Vitamin B₁₂, because the former may prevent anemia but allow progression of subacute combined degeneration.

A vegetarian diet which contains no animal products (including milk products or eggs) does not supply any Vitamin B₁₂. Patients following such a diet should be advised to take oral Vitamin B₁₂ regularly. The need for Vitamin B₁₂ is increased by pregnancy and lactation. Deficiency has been recognized in infants of vegetarian mothers who were breast fed, even though the mothers had no symptoms of deficiency at the time.

Laboratory Tests

During the initial treatment of patients with pernicious anemia, serum potassium must be observed closely the first 48 hours and potassium replaced if necessary.

Hematocrit, reticulocyte count, Vitamin B₁₂, folate and iron levels should be obtained prior to treatment. Hematocrit and reticulocyte counts should be repeated daily from the 5th to 7th days of therapy and then frequently until the hematocrit is normal. If folate levels are low, folic acid should also be administered. If reticulocytes have not increased after treatment or if reticulocyte counts do not continue at least twice normal as long as the hematocrit is less than 35%, diagnosis or treatment should be reevaluated. Repeat determinations of iron and folic acid may reveal a complicating illness that might inhibit the response of the marrow.

Patients with pernicious anemia have about three times the incidence of carcinoma of the stomach as the general population, so appropriate tests for this condition should be carried out when indicated.

Drug/Laboratory Test Interactions

Persons taking most antibiotics, methotrexate and pyrimethamine invalidate folic acid and Vitamin B₁₂ diagnostic blood assays.

Colchicine, para-aminosalicylic acid and heavy alcohol intake for longer than two weeks may produce malabsorption of Vitamin B₁₂.

Carcinogenesis, Mutagenesis

Long-term studies in animals to evaluate carcinogenic potential have not been done. There is no evidence from long-term use in patients with pernicious anemia that cyanocobalamin is carcinogenic.

Pernicious anemia is associated with an increased incidence of carcinoma of the stomach, but this is believed to be related to the underlying pathology and not to treatment with cyanocobalamin.

Pregnancy

Pregnancy Category C—Adequate and well-controlled studies have not been done in pregnant women.

However, Vitamin B₁₂ is an essential vitamin and requirements are increased during pregnancy.

Amounts of Vitamin B₁₂ that are recommended by the Food and Nutrition Board, National Academy of Science-National Research Council for pregnant women (4 mcg daily) should be consumed during pregnancy.

Nursing Mothers

Vitamin B₁₂ is known to be excreted in human milk. Amounts of Vitamin B₁₂ that are recommended by the Food and Nutrition Board, National Academy of Science-National Research Council for lactating

women (4 mcg daily) should be consumed during lactation.

Pediatric Use

Intake in children should be in the amount (0.5 to 3 mcg daily) recommended by the Food and Nutrition Board, National Academy of Science-National Research Council.

ADVERSE REACTIONS

Generalized

Anaphylactic shock and death have been reported with administration of parenteral Vitamin B₁₂ (see **WARNINGS**).

Cardiovascular

Pulmonary edema and congestive heart failure early in treatment; peripheral vascular thrombosis.

Hematological

Polycythemia vera.

Gastrointestinal

Mild transient diarrhea.

Dermatological

Itching; transitory exanthema.

Miscellaneous

Feeling of swelling of entire body.

OVERDOSAGE

No overdosage has been reported with this drug.

DOSAGE AND ADMINISTRATION

Avoid using the intravenous route. Use of this product intravenously will result in almost all of the vitamin being lost in the urine.

Pernicious Anemia

Parenteral Vitamin B₁₂ is the recommended treatment and will be required for the remainder of the patient's life. The oral form is not dependable. A dose of 100 mcg daily for six or seven days should be administered by intramuscular or deep subcutaneous injection. If there is clinical improvement and if a reticulocyte response is observed, the same amount may be given on alternate days for seven doses, then every three to four days for another two to three weeks. By this time hematologic values should have become normal. This regimen should be followed by 100 mcg monthly for life. Folic acid should be administered concomitantly if needed.

Patients With Normal Intestinal Absorption

Where the oral route is not deemed adequate, initial treatment similar to that for patients with pernicious anemia may be indicated depending on the severity of the deficiency. Chronic treatment should be with an oral B₁₂ preparation. If other vitamin deficiencies are present, they should be treated.

Schilling test

The flushing dose is 1000 mcg.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit.

HOW SUPPLIED

Product Number	NDC Number	Cyanocobalamin mcg/mL	Volume
4401	63323-044-01	1,000	1 mL in a 2 mL vial

1 mL vials are multiple dose vials, packaged 25 vials per tray.

Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].

PROTECT FROM LIGHT.

Use only if solution is clear and seal intact.



45813E

Revised: April 2008

Principal Display Panel - Swab Label

Easy•Touch®

Alcohol Prep Pads

Gamma-Sterilized

Isopropyl Alcohol, 70% by Volume

For External Antiseptic Use Only

Sterilized with Gamma Radiation

CONTAINS ONE PAD

Easy•Touch[®]

Alcohol Prep Pads

Gamma-Sterilized

Isopropyl Alcohol, 70% by Volume
For External Antiseptic Use Only
Sterilized with Gamma Radiation

CONTAINS ONE PAD

Principal Display Panel - 1 mL Vial Label

NDC 63323-044-01

4401

Cyanocobalamin Injection, USP

1,000 mcg/mL

For IM or SC Use

1 mL Multiple Dose Vial

Protect from Light.

Rx only



NDC 63323-044-01

4401

CYANOCOBALAMIN *INJECTION, USP*



1,000 mcg/mL

For IM or SC Use

1 mL Multiple Dose Vial

PROTECT FROM LIGHT.

Rx only

NDC: 76420-527-01 RX-Only

Physicians EZ Use B-12 Compliance Injection Kit™

Kit Contains

1 Cyanocobalamin Injection, USP 1,000 mcg/mL (1mL)

1 Isopropyl Alcohol 70% Prep Pad

1 Pair Nitrile Powder Free Sterile Gloves (M)

1 Drape

1 Adhesive Bandage

5 Non Sterile 4x4 Gauze

Needles and Syringes Not Included

1 Dose

Single Use Only

Distributed by:

Enovachem™

PHARMACEUTICALS

Torrance, CA 90501

NDC: 76420-527-01

Rx Only

Physicians EZ Use B-12 Compliance Injection Kit™

Kit Contains

- 1 Cyanocobalamin Injection, USP 1,000 mcg/mL (1mL)
 - 1 Isopropyl Alcohol 70% Prep Pad
 - 1 Pair Nitrile Powder Free Sterile Gloves (M)
 - 1 Drape
 - 1 Adhesive Bandage
 - 5 Non Sterile 4x4 Gauze
- Needles and Syringes Not Included
- 1 Dose
Single Use Only

Distributed by



PHYSICIANS EZ USE B-12 COMPLIANCE

physicians ez use b-12 compliance kit

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:76420-527
---------------------	-------------------------	---------------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:76420-527-01	1 in 1 CARTON; Type 1: Convenience Kit of Co-Package	05/23/2016	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 VIAL, MULTI-DOSE	1 mL
Part 2	1 POUCH	5 mL

Part 1 of 2

CYANOCOBALAMIN

cyanocobalamin solution

Product Information

Route of Administration INTRAMUSCULAR, SUBCUTANEOUS

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Cyanocobalamin (UNII: P6YC3EG204) (Cyanocobalamin - UNII:P6YC3EG204)	Cyanocobalamin	1000 ug in 1 mL

Inactive Ingredients

Ingredient Name	Strength
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
BENZYL ALCOHOL (UNII: LKG8494WBH)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		1 mL in 1 VIAL, MULTI-DOSE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA080557	10/18/2000	

Part 2 of 2

EASYTOUCH ALCOHOL PREP PADS STERILE

isopropyl alcohol swab

Product Information

Route of Administration TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Isopropyl Alcohol (UNII: ND2M416302) (Isopropyl Alcohol - UNII:ND2M416302)	Isopropyl Alcohol	70 mL in 100 mL

Inactive Ingredients

Ingredient Name		Strength		
WATER (UNII: 059QF0KO0R)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		5 mL in 1 POUCH; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph not final	part333A	05/01/2012		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
Unapproved drug other		10/18/2000		

Labeler - Asclemed USA, Inc. (059888437)

Establishment

Name	Address	ID/FEI	Business Operations
Asclemed USA, Inc.		059888437	REPACK(76420-527)

Revised: 6/2017

Asclemed USA, Inc.