

UP AND UP LANSOPRAZOLE- lansoprazole capsule, delayed release
Target Corporation

Target Corporation Lansoprazole Drug Facts

Active ingredient (in each capsule)

Lansoprazole 15 mg

Purpose

Acid Reducer

Use

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to lansoprazole

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- liver disease
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are

- taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding,

ask a health care professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**
- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F)
- keep product out of high heat and humidity
- protect product from moisture
- close cap tightly after use

Inactive ingredients

D&C red no. 28, D&C yellow no. 10, FD&C blue no. 1, FD&C red no. 40, gelatin, hypromellose, low substituted hydroxypropyl cellulose, mannitol, meglumine, methacrylic acid copolymer, pharmaceutical ink, polyethylene glycol, polysorbate 80, sodium lauryl sulfate, sugar spheres, talc, titanium dioxide

Questions or comments?

1-888-547-7400

Package/Label Principal Display Panel

Compare to active ingredient in Prevacid[®] 24 HR

lansoprazole delayed release capsules, 15 mg

acid reducer

24-hour

may take 1 to 4 days for full effect

sodium free

treats frequent heartburn

up & up[™]

3 bottles inside

ACTUAL SIZE

42 CAPSULES

42 CAPSULES

THREE 14-DAY COURSES OF TREATMENT



lansoprazole
 delayed release
 capsules, 15 mg
 acid reducer **24-hour**

Compare to active ingredient in
 Prevacid® 24 HR*

NDC 11673-280-03

lansoprazole
 delayed release
 capsules, 15 mg

acid reducer **24-hour**

may take 1 to 4 days for full effect
 sodium free
 treats frequent heartburn



3 bottles inside



ACTUAL SIZE

42
 CAPSULES

42 CAPSULES
 THREE 14-DAY COURSES OF TREATMENT

3T3D7 UW C11

GLUTEN FREE
 245 05 0677 R00 C-002016-01-009
 Distributed by Target Corporation
 Minneapolis, MN 55403
 TM & ©2023 Target Brands, Inc.



Drug Facts

Active ingredient (in each capsule) **Purpose**
 Lansoprazole 15 mg..... Acid Reducer

Use ■ treats frequent heartburn (occurs 2 or more days a week)
 ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to lansoprazole

Do not use

■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have ■ liver disease

■ had heartburn over 3 months. This may be a sign of a more serious condition.
 ■ heartburn with lightheadedness, sweating or dizziness

■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

■ frequent chest pain ■ frequent wheezing, particularly with heartburn
 ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain

Ask a doctor or pharmacist before use if you are

■ taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

■ your heartburn continues or worsens ■ you need to take this product for more than 14 days
 ■ you need to take more than 1 course of treatment every 4 months
 ■ you get diarrhea ■ you develop a rash or joint pain

If pregnant or breast-feeding, ask a health care professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

■ adults 18 years of age and older
 ■ this product is to be used once a day (every 24 hours), every day for 14 days
 ■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

■ swallow 1 capsule with a glass of water before eating in the morning
 ■ take every day for 14 days ■ do not take more than 1 capsule a day
 ■ swallow whole. Do not crush or chew capsules.
 ■ do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)

■ you may repeat a 14-day course every 4 months
 ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor

■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Tips for Managing Heartburn

- avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- eat slowly and do not eat big meals.
- do not eat late at night or just before bedtime.
- do not lie flat or bend over soon after eating.
- raise the head of your bed.
- wear loose-fitting clothing around your stomach.
- if you are over weight, lose weight.
- if you smoke, quit smoking.

KEEP THE CARTON. IT CONTAINS IMPORTANT INFORMATION.

DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING OR BLACK BAND AROUND THE CENTER OF EACH CAPSULE IS MISSING OR BROKEN.

*This product is not manufactured or distributed by Takeda Pharmaceutical U.S.A., Inc., owner of the registered trademark Prevacid®.

Drug Facts (continued)

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F) ■ keep product out of high heat and humidity
- protect product from moisture ■ close cap tightly after use

Inactive ingredients D&C red no. 28, D&C yellow no. 10, FD&C blue no. 1, FD&C red no. 40, gelatin, hypromellose, low substituted hydroxypropyl cellulose, mannitol, methylcellulose, methacrylic acid copolymer, pharmaceutical ink, polyethylene glycol, polyorbate 80, sodium lauryl sulfate, sugar spheres, bic, titanium dioxide

Questions or comments? 1-888-547-7400

UP AND UP LANSOPRAZOLE

lansoprazole capsule, delayed release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11673-280
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LANSOPRAZOLE (UNII: 0K5C5T2QPG) (LANSOPRAZOLE - UNII:0K5C5T2QPG)	LANSOPRAZOLE	15 mg

Inactive Ingredients

Ingredient Name	Strength
D&C RED NO. 28 (UNII: 767IP0Y5NH)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
LOW-SUBSTITUTED HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 2165RE0K14)	
MANNITOL (UNII: 3OWL53L36A)	
MEGLUMINE (UNII: 6HG8UB2MUY)	
METHACRYLIC ACID - ETHYL ACRYLATE COPOLYMER (1:1) TYPE A (UNII: NX76LV5T8J)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	PINK, GREEN	Score	no score
Shape	CAPSULE	Size	15mm
Flavor		Imprint Code	24HR
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11673-280-01	1 in 1 CARTON	06/18/2021	
1		14 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:11673-280-03	3 in 1 CARTON	06/18/2021	
2		14 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA202319	06/18/2021	

Labeler - Target Corporation (006961700)