SCRUB CARE EXIDINE -4 CHG- chlorhexidine gluconate liquid Xttrium Laboratories, Inc.

Drug Facts CareFusion 4%

Active Ingredients

Chlorhexidine Gluconate 4% Solution

Purpose

Antiseptic

Uses

surgical hand scrub: significantly reduces the number of microorganisms on the hands and forearms prior to surgery or patient care

healthcare personnel handwash: helps reduce bacteria that potentially can cause disease **patient preoperatve skin preparation:** for the preparation of the patient's skin prior to surgery **skin wound and general skin cleansing**

Warnings

For external use only

Allergy alert:

This product may cause a severe allergic reaction. Symptoms may include:

- wheezing/difficulty breathing
- shock
- facial swelling
- hives
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use

- if you or the patient is allergic to chlorhexidine gluconate or any other ingredient in this product
- in contact with meninges
- in the genital area
- as a preoperative skin preparation of the head or face

When using this product

- keep out of eyes, ears, and mouth. May cause serious and permanent eye injury if placed or kept in the eye during surgical procedures or may cause deafness when instilled in the middle ear through perforated eardrums
- if solution should contact these areas, rinse out promptly and thoroughly with water
- wounds which involve more than the superficial layers of the skin should not be routinely treated
- repeated general skin cleansing of large body areas should not be done except when advised by a health care provider

Stop use and ask a doctor if

irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.

Surgical hand scrub:

- wet hands and forearms with water
- scrub for 3 minutes with about 5 mL of product and a wet brush, paying close attention to the nails, cuticles, and interdigital spaces
- a separate nail cleaner may be used
- rinse thoroughly
- wash for an additional 3 minutes with 5 mL of product and rinse under running water
- dry thoroughly

Healthcare personnel handwash:

- wet hands with water
- dispense about 5 mL of product into cupped hands and wash in vigorous manner for about 15 seconds
- rinse and dry thoroughly

Patient preoperative skin preparation:

- apply product liberally to surgical site and swab for at least 2 minutes and dry with a sterile towel
- repeat procedure for an additional 2 minutes and dry with a sterile towel

Skin wound and general skin cleansing:

- thoroughly rinse the area to be cleaned with water
- apply the minimum amount of product necessary to cover the skin or wound area and wash gently
- rinse again thoroughly

Other information

- store at 20°-25° C (68°-77° F)
- avoid excessive heat above 40° C (104° F)

Inactive ingredients

cocamide DEA, fragrance, glucono-delta-lactone, hydroxyethylcellulose, isopropyl alcohol, lauramine oxide, PEG-75 lanolin, purified water, tridecyl alcohol

Questions or comments?

Call **1-800-587-3721**

Monday through Friday 8:00 AM to 4:30 PM CST

Package label and principal display panel

NDC 0116-2994-04

REF 29900-404

BD Scrub CareTM

ExidineTM-4 CHG 4%

Chlorhexidine Gluconate 4% Solution

Antiseptic

Non-sterile Solution

Contans:

4% Chlorhexidine Gluconate

For External Use Only

For Single Use Only

Distributed By

Becton, Dickinson and Company

Vernon Hills, IL 60061 USA

51-10096 Rev. A

4CFB04BTLBLE

Net Contents: 4 fl oz (118 mL)



Drug Facts

Active ingredient

Purpose

Chlorhexidine gluconate

4% solution Antiseptic

Uses

- surgical hand scrub: significantly reduces the number of microorganisms on the hands and forearms prior to surgery or patient care
- healthcare personnel handwash: helps reduce bacteria that potentially can cause disease
- patient preoperative skin preparation: for the preparation of the patient's skin prior to surgery
- skin wound and general skin cleansing



Drug Facts (continued)

Warnings

For external use only

Allergy alert:

This product may cause a severe allergic reaction. Symptoms may include:

- wheezing/difficulty breathing
- shock
- facial swelling
- hives
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use

- If you or the patient is allergic to chlorhexidine gluconate or any other ingredient in this product
- In contact with meninges
- In the genital area
- as a preoperative skin preparation of the head or face

When using this product

- keep out of the eyes, ears and mouth. May cause serious and permanent eye injury if placed or kept in the eye during surgical procedures or may cause deafness when instilled in the middle ear through perforated eardrums
- If solution should contact these areas, rinse out promptly and thoroughly with water
- wounds which involve more than the superficial layers of the skin should not be routinely treated
- repeated general skin cleansing of large body areas should not be done except when advised by a health care provider

¥

Drug Facts (continued)

Stop use and ask doctor if Irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

 use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.

Surgical hand scrub:

- wet hands and forearms with water
- scrub for 3 minutes with about 5 mL of product and a wet brush paying close attention to the nails, cuticles, and interdigital spaces
- a separate nail cleaner may be used
- rinse thoroughly
- wash for an additional 3 minutes with 5 mL of product and rinse under running water
- dry thoroughly

Healthcare personnel handwash:

- wet hands with water
- dispense about 5 mL of product into cupped hands and wash in a vigorous manner for 15 seconds
- rinse and dry thoroughly

Patient preoperative skin preparation:

 apply product liberally to surgical site and swab for at least 2 minutes and dry with a sterile towel

Drug Facts (continued)

 repeat procedure for an additional 2 minutes and dry with a sterile towel

Skin wound and general skin cleaning:

- thoroughly rinse the area to be cleaned with water
- apply the minimum amount of product necessary to cover the skin or wound area and wash gently
- rinse again thoroughly

Other information

- store at 20-25 °C (68-77 °F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients cocamide DEA,

fragrance, glucono-delta-lactone, hydroxyethylcellulose, isopropyl alcohol, lauramine

oxide, PEG-75 Ianolin, purified water, tridecyl alcohol

Questions or comments?

call 1-800-587-3721

Monday through Friday 8:00 AM to 4:30 PM CST

Laundering/Cleaning Instructions:

Chlorhexidine gluconate skin cleansers will cause stains if used with chlorine releasing products, such as chlorine bleach. Rinse completely and use only non-chlorine detergents. NDC 0116-2994-05

BD Scrub CareTM

ExidineTM-4 CHG 4%

Chlorhexidine Gluconate 4% Solution

Antiseptic

Non-sterile Solution

Contains:

4% Chlorhxidine Gluconate

For External Use Only

For Single Use Only

Distributed By


Becton, Dickinson and Company

Vernon Hills, IL 60061 USA

51-10081 Rev. A

ETOBTLLBLE

Net Contents: 4 fl oz (118 mL)



NDC 0116-2994-08

REF 29900-408

BD Scrub CareTM

ExidineTM-4 CHG 4%

Chlorhexidine Gluconate

4% Solution

Antiseptic

Contains: 4% Chlorhexidine Gluconate

For External Use Only

Distributed by


Becton, Dickinson and Company

Vernon Hills, IL 60061 USA

51-10097 Rev. A

4CF08BTLBLE

Net Contents: 8 fl oz (237 mL)



NDC 0116-2994-16

REF 29900-416

BD Scrub Care™

ExidineTM-4 CHG 4%

Chlorhexidine Gluconate

4% Solution

Antiseptic

Contains:

4% Chlorhexidine Gluconate

For External Use Only

Distributed By

Becton, Dickinson and Company

Vernon Hills, IL 60061 USA

51-10098 Rev. A

4CFB16BTLLBLC

Net Contents: 16 fl oz (1 pt) (473 mL)



NDC 0116-2994-17

REF 29900-416P

BD Scrub CareTM

ExidineTM-4 CHG 4%

Chlorhexidine Gluconate

4% Solution

Antiseptic

Contains:

4% Chlorhexidine Gluconate

For External Use Only

Distributed By

Becton, Dickinson and Company

Vernon Hills, IL 60061 USA

51-10099 Rev. A

4CFB16BTLPLBLC

Net Contents: 16 fl oz (1 pt) (473 mL)



NDC 0116-2994-32

REF 29900-432

BD Scrub CareTM

ExidineTM-4 CHG 4%

Chlorhexidine Gluconate

4% Solution

Antiseptic

Contains: 4% Chlorhexidine Gluconate

For External Use Only

Distributed by

Becton, Dickinson and Company

Vernon Hills, IL 60061 USA

51-10100 Rev. A

4CFB32BTLLBLC

Net Contents: 32 fl oz (1 qt) (946 mL)



NDC 0116-2994-32 REF 29900-432

Chlorhexidine Gluconate 4% Solution Antiseptic

Contains: 4% Chlorhexidine Gluconate For External Use Only

Distributed by Becton, Dickinson and Company Vernon Hills, IL 60061 USA

51-10100 Rev. A 4CFB32BTLLBLC



Net Contents: 32 fl oz (1 qt) (946 mL)

Drug Facts

Active ingredient Chlorhexidine gluconate 4% solution

Warnings For external use only

shock
facial swelling
hives

cause disease skin wound and general skin deansing

surgical hand scrub: significantly reduces the number of microorganisms on the hands and forearms prior to surgery or patient care.

I healthcare personnel handwash: fidps reduce bacteria that potentially can cause disease.

Allergy alert: This product may cause a severe allergic reaction. Symptoms may include: wheeling difficulty breathing

If an allergic reaction occurs, stop use and seek medical help right away

To not use a superior to the s

I more sentations that When using this product.

I keep out of eyes, ear, and mouth, May cause serious and permanent eye injuly if permitted to enter and remain in the eye or may cause deaffness when institled in the middle ear through perforated eardinums.

I solution should cortact these areas, rines out promptly and thoroughly with

water

wounds which involve more than the superficial layers of the skin should not be routinely tracted

a repeated general skin cleansing of large body areas should not be done except when advised by a health care provider

Stop use and ask a doctor if in fraction, sensitiation, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Central Center right away.

Purpos

Drug Facts (continued)

Directions

usewith care in premature infants or infants under 2 months of age. These products may cause initiation or chamical burns.

Surgical hand saruk

**with hands and foreams withwater

**surb for a limites with about 5 mL of product and a wet brush paying close attention to the naist, cuticles, and interdigital spaces

**a separate and closurer may be used

**intes thoroughly

**wash for an additional 3 minutes with 5 mL of product and rinse under

Healthcare personnel handwash:

**wat hands with water

**deprese about 5 ml. of product into cupped hands and wash in a vigorous manner for about 15 seconds

**einse and by thoroughly

**inse and by thoroughly

Skin wound and general skin cleaning:

throughly rinse the area to be cleaned with water

apply the minimum amount of product necessary to cover the skin or wound
area and wash gently

■ rinse again thoroughly

Other information
store at 20-25°C (68-77°F)
avoid excessive heat above 40°C (104°F)

Inactive ingredients cocomide DEA, fragrance, glucono-deita-lactone, hydracysthylcelulosa, topropyl dicohol, lauramine oxide, PEG-75 lanolin, purified water, tridecyl alcohol

Questions or comments? call 1-800-587-3721 Monday through Friday 8:00 AM to 4:30 PM CST

NDC 0116-2994-33

REF 29900-432P

BD Scrub CareTM

ExidineTM-4 CHG 4%

Chlorhexidine Gluconate

4% Solution

Antiseptic

Contains: 4% Chlorhexidine Gluconate

For External Use Only

Distributed by


Becton, Dickinson and Company

Vernon Hills, IL 60061 USA

51-10101 Rev. A

4CF32BTLPLBLC

Net Contents: 32 fl oz (1 qt) (946 mL)



NDC 0116-2994-33 REF 29900-432P

Chlorhexidine Gluconate 4% Solution Antiseptic

Contains: 4% Chlorhexidine Gluconate For External Use Only

Distributed by Becton, Dickinson and Company Vernon Hills, IL 60061 USA

51-10101 Rev. A 4CF32BTLPLBLC



Net Contents: 32 fl oz (1 qt) (946 mL)

Drug Facts

Warnings For external use only

Active ingredient Chlorhexidine gluconate 4% solution

cause disease skin wound and general skin deansing

= surgical hand scrub: significantly reduces the number of microorganisms on the hands and forearms prior to surgery or patient care.

• health care personnal handwash: helps reduce bacteria that potentially can cause disease.

Allergy alert
This product may cause a severe allergic reaction. Symptoms may include:
Whese Ing-difficulty breathing
Shock
Facual swelling
Hives
Teach
If an allergic reaction occurs, stop use and seek medical help right away.

To anothers

If an along's recomm occurs, stop use and seek measure map right away.

To not use

If you or the patient is alongly to chlorhexidine gluconate or any other
ingredient in this product

in contact with meningss

in the gen

In the gential area
When using this product
Leap out of eyes, ears, and mouth. May cause serious and permanent eye injury
If permitted to arter and remain in this eye or may cause deafness when
Institled in the middle ear of through perforated eardiums.
If solution should contact these areas, rinse out promptly and thoroughly with

water

would which involve more than the superficial layers of the skin should not be routinely tracted

respected general skin cleansing of large body areas should not be done accept when advised by a health care provider

Stop use and ask a doctor if in fraction, senatitation, or allergic reaction occurs. These may be signs of a serious condition.

Respected reach of children.

If swallowed, get medical help or contact a Poison Central Center right away.

Purpose

Directions
use with care in premature infants or infants under 2 months of age. These products may cause initiation or chemical burns.

Drug Facts (continued)

Surgical hand scruts

Wat hands and foreams withwater

State for 3 minutes with about 5 mL of product and awet brush paying close attantion to the nails, cuticles, and interdigital spaces

a separate nail cleaner may be used

insectionoughly

■ rinse thoroughly ■ wash for an additional 3 minutes with 5 mL of product and rinse under running water ■ dry thoroughly

Healthcare personnel handwash:

**wat hands with water

**wat hands with water

**desperse about 5 mil. of product into cupped hands and wash in a vigorous

manner for about 5 mil. of product into cupped hands and wash in a vigorous

manner for about 15 seconds

**inse and dry thoroughly

Slain wound and general slain dearning:

**Thoroughly insis the area to be a cleaned with water

**apply the minimum amount of product necessary to cover the skin or woun

area and wash gently

**inse again theroughly

Other information
store at 20-25°C (68-77°F)
avoid excessive heat above 40°C (104°F)

Inactive ingredients cocamide DEA, fragrance, glucono-deito-lactone hydroxysthylcallulose, isopropyl alcohol, lauramine oxide, PEG-75 lanolin, purified water, tridecyl alcohol

Questions or comments? call 1-800-587-3721 Monday through Friday 8:00 AM to 4:30 PM CST

NDC 0116-2994-01

REF 29900-401

BD Scrub CareTM

ExidineTM-4 CHG 4%

Chlorhexidine Gluconate

4% Solution

Antiseptic

Contains:

4% Chlorhexidine Gluconate

For External Use Only

Distributed by


Becton, Dickinson and Company

Vernon Hills, IL 60061 USA

51-10095 Rev. A

4CF1GBTLLBLD

Net Contents: 128 fl oz (1 gal) (3.8 L)



SCRUB CARE EXIDINE -4 CHG

chlorhexidine gluconate liquid

_				
D	MOD	nct	Into	rmation

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0116-2994
--------------	----------------	--------------------	---------------

Route of Administration TOPICAL

Active Ingredient/Active Moiety Ingredient Name Basis of Strength CHLORHEXIDINE GLUCONATE (UNII: MOR84MUD8E) (CHLORHEXIDINE UNII:R4KO0DY52L) CHLORHEXIDINE GLUCONATE in 100 mL

Inactive Ingredients				
Ingredient Name	Strength			
LAURAMINE O XIDE (UNII: 4F6 FC4MI8 W)				
COCO DIETHANOLAMIDE (UNII: 92005F972D)				
PEG-75 LANOLIN (UNII: 09179OX7TB)				
HYDROXYETHYL CELLULOSE (2000 CPS AT 1%) (UNII: S38J6RZN16)				
ISOPROPYL ALCOHOL (UNII: ND2M416302)				
WATER (UNII: 059QF0KO0R)				
TRIDECYL ALCOHOL (UNII: 819428 H868)				
GLUCONOLACTONE (UNII: WQ29KQ9POT)				

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:0116-2994- 04	118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	11/0 1/20 12		
2	NDC:0116-2994- 08	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	11/0 1/20 12		
3	NDC:0116-2994- 16	473 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	11/0 1/20 12		
4	NDC:0116-2994- 17	473 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	11/0 1/20 12		
5	NDC:0116-2994- 32	946 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	11/0 1/20 12		
6	NDC:0116-2994- 33	946 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	11/0 1/20 12		
7	NDC:0116-2994- 01	3786 mL in 1 JUG; Type 0: Not a Combination Product	11/0 1/20 12		

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
NDA	NDA0 19 125	11/0 1/20 12		

Labeler - Xttrium Laboratories, Inc. (007470579)

Registrant - Xttrium Laboratories, Inc. (007470579)

Establishment					
Name	Address	ID/FEI	Business Operations		
Xttrium Laboratories, Inc.		007470579	manufacture(0116-2994)		

Revised: 6/2020 Xttrium Laboratories, Inc.