ASPIRIN- aspirin tablet, film coated Chain Drug Consortium

Premier Value 44-157

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID)*
*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

temporarily relieves:

- minor pain of arthritis
- headache
- muscle pain
- menstrual pain
- toothache
- pain and fever of colds

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- facial swelling
- shock
- hives
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- have had stomach ulcers or bleeding problems
- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- you are taking a diuretic
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients

corn starch, hypromellose, polyethylene glycol, propylene glycol

Questions or comments?

1-800-426-9391

Principal display panel

†COMPARE TO THE ACTIVE INGREDIENT IN GENUINE BAYER® ASPIRIN

Premier Value®

Film-Coated **Aspirin**PAIN RELIEVER/FEVER REDUCER
(NSAID)

Aid for relief of headache, cold discomforts, muscular aches

100 Tablets 325 mg EACH

INDEPENDENTLY TESTED
PV
SATISFACTION GUARANTEED

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Distributed By: Pharmacy Value Alliance, LLC 407 East Lancaster Avenue, Wayne, PA 19087

If for any reason you are not satisfied with this product, please return it to the store where purchased for a full refund.

†This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Genuine

Bayer® Aspirin. 50844 REV0122C15712



Premier Value 44-157

an allergic reaction occurs.

, such as

pain that does not get better

STOP PEELING

Inactive Ingredients

■ use by expiration date on package excursions permitted between

years and over: take 1 or 2 tablets

each dose

ASPIRIN aspirin tablet, film coated **Product Information HUMAN OTC DRUG** NDC:68016-747 **Product Type** Item Code (Source) **ORAL Route of Administration Active Ingredient/Active Moiety Ingredient Name Basis of Strength** Strength ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E) **ASPIRIN** 325 mg

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	

Product Characteristics				
Color	white	Score	no score	
Shape	ROUND	Size	10mm	
Flavor		Imprint Code	44;157;AS PIRIN	
Contains				

P	Packaging					
#	Item Code	Package Description	Marketing Start Date	Marketing End Date		
1	NDC:68016-747- 10	100 in 1 BOTTLE; Type 0: Not a Combination Product	02/10/1996			
2	NDC:68016-747- 50	300 in 1 BOTTLE; Type 0: Not a Combination Product	02/10/1996	02/20/2023		
3	NDC:68016-747- 30	500 in 1 BOTTLE; Type 0: Not a Combination Product	02/10/1996	10/29/2021		

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M013	02/10/1996		

Labeler - Chain Drug Consortium (101668460)

Establishment				
Name	Address	ID/FEI	Business Operations	
LNK International, Inc.		038154464	pack(68016-747)	

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(68016-747)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(68016-747)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(68016-747)

Establishment				
Name	Address	ID/FEI	Business Operations	
LNK International, Inc.		967626305	pack(68016-747)	

Revised: 2/2024 Chain Drug Consortium