

CAREONE DIARRHEA CONTROL- loperamide hcl tablet, film coated
American Sales Company

American Sales Company Diarrhea Control Drug Facts

Active ingredient (in each caplet)

Loperamide HCl 2 mg

Purpose

Anti-diarrheal

Use

controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Heart alert: Taking more than directed can cause serious heart problems or death

Do not use

if you have bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product

tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

Other information

- store at 20°-25°C (68°-77°F)
- see end panel for lot number and expiration date

Inactive ingredients

anhydrous lactose, carnauba wax, D&C yellow no. 10 aluminum lake, FD&C blue no. 1 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch

Questions or comments?

1-800-719-9260

Principal Display Panel

Compare to the active ingredient in Imodium[®] A-D Caplets

DIARRHEA CONTROL

Loperamide Hydrochloride Tablets, 2 mg

Anti Diarrheal

Anti Diarrheal

Controls the symptoms of diarrhea

Gluten Free

See New Warnings

Actual Size

OUR PHARMACISTS RECOMMEND

24 CAPLETS*

*Capsule-Shaped Tablets

DO NOT USE IF BLISTER UNIT IS BROKEN OR TORN

**This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Imodium® A-D Caplets.

DISTRIBUTED BY: FOODHOLD U.S.A., LLC
LANDOVER, MD 20785 · 1-877-846-9949
©2019 Ahold Delhaize Licensing Sarl
Quality guaranteed or your money back.

CAREone®

NDC 41520-813-82

Compare to the active ingredient
in Imodium® A-D Caplets**

DIARRHEA CONTROL

Loperamide Hydrochloride Tablets, 2 mg
Anti Diarrheal

Anti Diarrheal
Controls the symptoms of diarrhea

Gluten Free

See New Warnings



Actual Size



24 CAPLETS*

*Capsule-Shaped Tablets

22462 OF 02

CODE AREA

Drug Facts

Active ingredient (in each caplet) Purpose
Loperamide HCl 2 mg Anti-diarrheal

Use Controls symptoms of diarrhea, including Traveler's Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl
Heart alert: Taking more than directed can cause serious heart problems or death

Do not use if you have bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool
- history of liver disease
- history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product tiredness, dizziness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
 - diarrhea lasts for more than 2 days
 - you get abdominal swelling or bloating.
- These may be signs of a serious condition.

If pregnant or breastfeeding, ask a health professional before use.

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact Poison Control Center right away (1-800-222-1222).


Directions

- drink plenty of clear fluids to help prevent dehydration caused by diarrhea
- find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (80-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (24 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

Other information

- store at 20°-25°C (68°-77°F)
- see end panel for lot number and expiration date



CAREONE DIARRHEA CONTROL

loperamide hcl tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41520-813
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	

Product Characteristics

Color	GREEN	Score	2 pieces
Shape	OVAL	Size	10mm
Flavor		Imprint Code	L2
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41520-813-53	12 in 1 CARTON	09/14/2016	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

2	NDC:41520-813-62	24 in 1 CARTON	09/14/2016	
2		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:41520-813-67	1 in 1 CARTON	09/14/2016	
3		48 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075232	09/14/2016	

Labeler - American Sales Company (809183973)

Revised: 12/2019

American Sales Company