ANTI-DIARRHEAL- loperamide hydrochloride capsule, liquid filled MEIJER, INC.

Drug Facts

Active ingredient (in each capsule)

Loperamide HCI 2 mg

Purpose

Anti-diarrheal

Use

controls symptoms of diarrhea, including Travelers' Diarrhea.

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCI.

Do not use

- if you have bloody or black stool
- in children under 12 years of age

Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease

Ask a doctor or pharmacist before use if you are

taking antibiotics.

When using this product

tiredness, drowsiness, or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- drink plenty of clear fluids to help prevent dehydration caused by diarrhea
- not for use in children under 12 years of age
- adults and children 12 years of age and over: 2 softgels after the first loose stool; 1 softgel after each subsequent loose stool; but no more than 4 softgels in 24 hours

Other information

- store at 20°-25°C (68°-77°F). Protect from light.
- avoid excessive heat above 40°C (104°F)
- do not use if carton or blister unit is open or torn
- see side panel for lot number and expiration date

Inactive ingredients

butylated hydroxyanisole, edible ink, FD&C Blue #1, gelatin, glycerin, glyceryl caprylate, polyoxyl 40 hydrogenated castor oil, purified water.

Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

Principal Display Panel

Compare to Imodium® A-D active ingredient

Loperamide HCI

Soft Gelatin Capsules 2 mg

Anti-Diarrheal

Controls the Symptoms of Diarrhea

Suitable for Adults and Children 12 Years and Over

SOFTGELS**

**each Liquid-Filled capsule contains 2 mg Loperamide HCI

*This product is not manufactured or distributed by McNeil Consumer & Specialty Pharmaceuticals, distributor of Imodium® A-D.

THIS PRODUCT IS PACKAGED IN A CHILD RESISTANT AND TAMPER EVIDENT PACKAGE. USE ONLY IF BLISTERS ARE INTACT.


KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

DIST. BY MEIJER DISTRIBUTION, INC.

GRAND RAPIDS, MI 49544

www.meijer.com

Product Label



Anti-Diarrheal/Loperamide HCl 2 mg

Softgels

meijer

Stop use and ask a doctor if symptoms get worse diarrhea lasts for more than 2 days	Questions or comments? Cuestions or comments?
When using this product thedness, drowsiness, or dizziness may occur. Be careful when driving or operating machinery.	Inactive ingredients butylated hydroxyanisole, edible ink, FD&C blue #1, gelatin, glycerin, glyceryl caprylate, polyoxyl 40 hydrogenated castor oil, purified water
Ask a doctor or pharmacist before use if you are taking antibiotics.	do not use if carton or blister unit is open or torn see side panel for lot number and expiration date
Ask a doctor before use if you have ■ fever ■ mucus in the stool ■ a history of liver disease	Other information ■ store at 20°-25°C (68°-77°F). Protect from light. ■ avoid excessive heat above 40°C (104°F) ■ do not use if carton of bister unit is page of tors
Do not use ■ if you have bloody or black stool ■ in children under 12 years of age	after the first loose stool; 1 softgel after each subsequent loose stool; but no more than 4 softgels in 24 hours
Warnings Milergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCI. Heart alert: Taking more than directed can cause serious heart problems or death.	Directions a drink plenty of clear fluids to help prevent dehydration caused by diarrhea not for use in children under 12 years of age adults and children 12 years of age and over: 2 softgels control over: 2 softgels con
USG Controls symptoms of diarrhea, including Travelers' Diarrhea.	Keep out of reach of children. In case of overdose, get Meep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
lsərhrsib-itnA	If pregnant or breast-feeding, ask a health professional before use.
esoqruq taiseare furpose ingrese ingre	you get abdominal swelling or bulging. These may be signs of a serious condition.
Drug Facts	Drug Facts (continued)

This product is not manufactured or distributed by McNeil Consumer & Specialty Pharmaceuticals, distributor of Imodium® A-D.



loperamide hydrochloride	e capsule, liquio	d filled				
Product Information	I					
Product Type	HUMAN OTO	C DRUG	Item Code (Source)		NDC:41250-520	
Route of Administration	ORAL					
Active Ingredient/Act	-					
	ngredient Nan			Basis of Stren		Strengt
LOPERAMIDE HYDROCHLOI UNII:6X9OC3H4II)	RIDE (UNII: 77TI35	393C) (LOPERA	RAMIDE - LOPERAMIDE HYDROCHLORIDE		_	2 mg
Inactive Ingredients	Ingred	lient Name			5	trength
						LICIULII
BUTYLATED HYDROXYANIS	OLE (UNII: REK496					uengui
						trengti
BUTYLATED HYDROXYANIS FD&C BLUE NO. 1 (UNII: H3 GELATIN (UNII: 2G86QN327L)	R47K3TBD)					uengui
FD&C BLUE NO. 1 (UNII: H3	R47K3TBD))					trengtin
FD&C BLUE NO. 1 (UNII: H3) GELATIN (UNII: 2G86QN327L) GLYCERIN (UNII: PDC6A3C00) GLYCERYL MONOCAPRYLA	R47K3TBD)) X) FE (UNII: TM2TZD4	50K2U) 1G4A)				uengui
FD&C BLUE NO. 1 (UNII: H3) GELATIN (UNII: 2G86QN327L) GLYCERIN (UNII: PDC6A3C00) GLYCERYL MONOCAPRYLA POLYOXYL 40 HYDROGENA	R47K3TBD)) X) FE (UNII: TM2TZD4	50K2U) 1G4A)	5GQ8F)			uengui
FD&C BLUE NO. 1 (UNII: H3) GELATIN (UNII: 2G86QN327L) GLYCERIN (UNII: PDC6A3C00) GLYCERYL MONOCAPRYLA	R47K3TBD)) X) FE (UNII: TM2TZD4	50K2U) 1G4A)	5GQ8F)			uengun
FD&C BLUE NO. 1 (UNII: H3) GELATIN (UNII: 2G86QN327L) GLYCERIN (UNII: PDC6A3C00) GLYCERYL MONOCAPRYLA POLYOXYL 40 HYDROGENA	R47K3TBD)) X) FE (UNII: TM2TZD4 TED CASTOR OII	50K2U) 1G4A)	5GQ8F)			uengun
FD&C BLUE NO. 1 (UNII: H3) GELATIN (UNII: 2G86QN327L) GLYCERIN (UNII: PDC6A3C00) GLYCERYL MONOCAPRYLA POLYOXYL 40 HYDROGENA WATER (UNII: 059QF0K00R)	R47K3TBD)) X) FE (UNII: TM2TZD4 TED CASTOR OII	50K2U) 1G4A)	5GQ8F)		no score	
FD&C BLUE NO. 1 (UNII: H3) GELATIN (UNII: 2G86QN327L) GLYCERIN (UNII: PDC6A3C00 GLYCERYL MONOCAPRYLA POLYOXYL 40 HYDROGENA WATER (UNII: 059QF0K00R) Product Characterist Color	R47K3TBD)))X) FE (UNII: TM2TZD4 TED CASTOR OII	50K2U) IG4A) L (UNII: 7YC686	5GQ8F)			
FD&C BLUE NO. 1 (UNII: H3) GELATIN (UNII: 2G86QN327L) GLYCERIN (UNII: PDC6A3C00 GLYCERYL MONOCAPRYLA POLYOXYL 40 HYDROGENA WATER (UNII: 059QF0K00R) Product Characterist	R47K3TBD)))X) FE (UNII: TM2TZ D4 TED CASTOR OII tiCS blue	50K2U) IG4A) L (UNII: 7YC686 Size	5GQ8F) • t Code		no score	

Pa	ckaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	NDC:41250-520- 24	24 in 1 CARTON	08/31/2016	04/26/2024
1		1 in 1 BLISTER PACK; Type 0: Not a Combination		
		Product		
		Product		
Μ	arketing	Information		
Μ	arketing Marketing Category		Marketing Start Date	Marketing End Date

Labeler - MEIJER, INC. (006959555)

Revised: 10/2021

MEIJER, INC.