CAREONE TUSSIN DM COUGH PLUS CHEST CONGESTION- dextromethorphan hbr, guaifenes in solution

American Sales Company

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

American Sales Company Tussin DM Drug Facts

Active ingredients (in each 10 mL)

Dextromethorphan HBr, USP 20 mg Guaifenesin, USP 200 mg

Purposes

Cough suppressant

Expectorant

Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

Warnings

Do not use

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

Stop use and ask a doctor if

cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- keep dosing cup with product
- mL = milliliter
- this adult product is not intended for use in children under 12 years of age

age	dose	
adults and children 12 years and over	10 mL every 4 hours	
children under 12 years	do not use	

Other information

- each 10 mL contains: sodium 6 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

anhydrous citric acid, FD&C red no. 40, flavor, glycerin, high fructose corn syrup, menthol, propylene glycol, purified water, sodium benzoate, sodium citrate, sucralose

Questions or comments?

1-800-719-9260

Principal Display Panel

Compare to the active ingredients in Robitussin® Cough + Chest Congestion DM

TUSSIN DM

COUGH + CHEST CONGESTION

Cough Suppressant-Dextromethorphan HBr

Expectorant-Guaifenesin

Relieves:

Cough

Mucus

Adult

For Ages 12 & Over

Non-Drowsy

Alcohol Free

Gluten Free

OUR PHARMACISTS RECOMMEND

Cherry Flavor

4 FL OZ (118mL)



IS BROKEN OR MISSING DO NOT USE IF PRINTED NECKBAND



Compare to the active ingredients in Robitussin® Cough + Chest Congestion DM



Compare to the active ingredients in Robitussin® Cough + Chest Congestion DM^o

COUGH + CHEST CONGESTION

Cough Suppressant-**Dextromethorphan HBr Expectorant-GualfenesIn**

Relieves:

- Cough
- Mucus

Adult

For Ages 12 & Over

Non-Drowsy Alcohol Free

Gluten Free



Cherry Flavor

4 FL OZ (118mL)

COUGH + CHEST CONGESTION

Cough Suppressant-Dextromethorphan HBr **Expectorant-Gualfenesin**

Relieves:

- Cough
- Mucus

Adult

For Ages 12 & Over

Non-Drowsy **Alcohol Free**

Gluten Free





4 FL OZ (118mL)

Drug Facts

Active ingredients (in each 10 mL) Dextromethorphan HBr,

..Cough suppressant USP 20 mg......Co Guaifenesin, USP 200 mg....

Purposes

Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur
- with a cold

 helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAO) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- Ask a doctor before use if you have a cough that occurs with too much phlegm (mucus) a cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Stop use and ask a doctor if cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

If pregnant or breast-feeding, ask a health ... program or preast-feeding, ask a healt professional before use. Keep out of reach of children. In case of overdose not medical.

overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- Directions

 do not take more than 6 doses in any
 24-hour period

 measure only with dosing cup provided

 keep dosing cup with product

 mt. milliliter

 this adult product is not intended for use in
 children under 12 years of age

Drug Facts (continued)

age	dose
adults and children 12 years and over	10 mL every 4 hours
children under 12 years	do not use

Other information

■ each 10 mL contains: sodium 6 mg ■ store at 20-25°C (68-77°F).

Inactive ingredients anhydrous citric acid, FD&C red no. 40, flavor, glycerin high fructose corn syrup, menthol, propylene glycol, purified water, sodium benzoate, sodium citrate, sucralose

Questions or comments? 1-800-719-9260

*This product is not manufactured or distributed by Pfizer, distributor of Robitussin® Cough + Chest Congestion DM.

DISTRIBUTED BY FOODHOLD U.S.A., LLC LANDOVER, MD 20785 · 1-877-846-9949 ©2017 S&S Brands, LLC

Quality guaranteed or your money back.



Learn about teen medicine abuse

www.StopMedicineAbuse.org





LOT NO.

FXP

: 35926 OF CL

CAREONE TUSSIN DM COUGH PLUS CHEST CONGESTION

dextromethorphan hbr, guaifenesin solution

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:41520-636

Route of Administration ORAL

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9 D2RTI9 KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 10 mL	
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg in 10 mL	

Inactive Ingredients			
Ingredient Name	Strength		
ANHYDRO US CITRIC ACID (UNII: XF417D3PSL)			
FD&C RED NO. 40 (UNII: WZB9127XOA)			
GLYCERIN (UNII: PDC6 A3C0 OX)			
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)			
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)			
WATER (UNII: 059QF0KO0R)			
SODIUM BENZOATE (UNII: OJ245FE5EU)			
MENTHOL (UNII: L7T10EIP3A)			
SODIUM CITRATE (UNII: 1Q73Q2JULR)			
SUCRALOSE (UNII: 96K6UQ3ZD4)			

Product Characteristics			
Color	RED (clear)	Score	
Shape		Size	
Flavor	CHERRY (Fruity)	Imprint Code	
Contains			

P	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41520-636-26	1 in 1 CARTON	08/05/2016	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:41520-636-34	1 in 1 CARTON	09/05/2017	
2		237 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	08/05/2016	

Labeler - American Sales Company (809183973)

Revised: 10/2017 American Sales Company