

PRECAINE B- benzocaine topical anesthetic gel
Pascal Company, Inc.

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Precaine B

Directoons

Application Directions:

For topical use only

For best results apply to previously dried oral mucosa with cotton swab or similar applicator for adequate control of pain

Removal of excess saliva during application minimizes dilution of the anesthetic and perimts maximum penetration

Not more than 1.2mg per Kg body weight per patient should be applied during a 24-hour period

Warnings

Warning:

Recommended dosage should not be exceeded due to possible side effects. Keep out of the reach of children. For professional use only. Do not use in the eyes. Avoid swallowing. Not for home or unsupervised consumer use. Not for use on children 2 and younger or pregnant or nursing women.

Contraindications

Contraindications:

Precaine is contraindicated in patients with known hypersensitivity to benzocaine or PABA.

Other Information

For product MSDS information, please go to www.pascaldental.com or contact Pascal directly

Store product between 60 degress F (16 degress C) and 86 degress F (30 degress C)

Rx only in USA

Made in USA

Precaine B

Precaine B

Topical Anesthetic Gel for oral use

Strawberry

Contain: Benzocaine 20% in a flavored aqueous base

Net Contents 30g

Pascal International

2929 NE Northup Way

Bellevue, WA 98004

425.827.4694

REF 15-350

SN72505/0912

Application Directions:

- For topical use only.
- For best results apply to previously dried oral mucosa with cotton swab or similar applicator, for adequate control of pain.
- Removal of excess saliva during application minimizes dilution of the anesthetic and permits maximum penetration.
- Not more than 1.2mg per Kg body weight per patient should be applied during a 24-hour period.



Warnings:

Recommended dosage should not be exceeded due to possible side effects. Keep out of the reach of children. For professional use only. Do not use in the eyes. Avoid swallowing. Not for home or unsupervised consumer use. Not for use on children 2 and younger, or pregnant or nursing women.

Contraindications:

Precaine is contraindicated in patients with known hypersensitivity to benzocaine or PABA.

For product MSDS information please go to www.pascal-intl.com or contact Pascal directly.

Pascal
International
2929 NE Northup Way
Bellevue, WA 98004 USA
425.827.4694

REF 15-350

88°F (30°C)

80°C Only (in USA)
Made in USA
SN72505/0912



PRECAINE B

benzocaine topical anesthetic gel

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:10866-0087
Route of Administration	DENTAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	221 mg in 1 g

Product Characteristics

Color	red	Score	
Shape		Size	
Flavor	STRAWBERRY	Imprint Code	

Contains**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:10866-0087-1	30 g in 1 JAR; Type 0: Not a Combination Product	04/27/2009	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		04/27/2009	

Labeler - Pascal Company, Inc. (009260217)**Establishment**

Name	Address	ID/FEI	Business Operations
Pascal Company, Inc.		009260217	manufacture(10866-0087)

Revised: 12/2022

Pascal Company, Inc.