

**TOPCARE NASAL DECONGESTANT MAXIMUM STRENGTH NON DROWSY-  
pseudoephedrine hcl tablet, film coated  
Topco Associates LLC**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Topco Associates LLC. Nasal Decongestant Drug Facts**

**Active ingredient (in each tablet)**

Pseudoephedrine HCl 30 mg

**Purpose**

Nasal decongestant

**Uses**

- temporarily relieves sinus congestion and pressure
- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies

**Warnings**

**Do not use**

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland

**When using this product**

**do not exceed recommended dosage**

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur

- symptoms do not improve within 7 days or occur with a fever

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

adults and children 12 years and over	<ul style="list-style-type: none"> <li>• take 2 tablets every 4 to 6 hours</li> <li>• do not take more than 8 tablets in 24 hours</li> </ul>
children ages 6 to 11 years	<ul style="list-style-type: none"> <li>• take 1 tablet every 4 to 6 hours</li> <li>• do not take more than 4 tablets in 24 hours</li> </ul>
children under 6 years	do not use this product in children under 6 years of age

**Other information**

- **each tablet contains:** calcium 20 mg
- store at 20° -25° C (68° -77° F)
- **do not use if blister unit is broken or torn**

**Inactive ingredients**

carnauba wax, dibasic calcium phosphate dihydrate, FD&C red no. 40 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polysorbate 80, silicon dioxide, titanium dioxide

**Questions or comments?**

**1-888-423-0139**

**Principal Display Panel**

TopCare® health

COMPARE TO SUDAFED® CONGESTION ACTIVE INGREDIENT

NON-DROWSY

Nasal Decongestant

PSEUDOEPHEDRINE HYDROCHLORIDE TABLETS, 30 mg

MAXIMUM STRENGTH

- Sinus Pressure + Congestion
- Pseudoephedrine HCl

24 TABLETS

actual size



This product is not manufactured or distributed by McNeil Consumer HealthCare, distributor of Sudafed® Congestion.



Scan here for more information or call 1-888-423-0139



Disbuted by Perrigo®  
Alligan, MI 49010 PBRAD21  
This TopCare® product is laboratory tested to guarantee its highest quality. Your total satisfaction is guaranteed.



**Drug Facts**  
Pseudoephedrine HCl 30 mg.....Nasal decongestant

**Active ingredient (in each tablet)**  
Pseudoephedrine HCl 30 mg.....Nasal decongestant

**Uses**  
■ temporarily relieves sinus congestion and pressure  
■ temporarily relieves nasal congestion due to the common cold,  
■ fever or other upper respiratory allergies

**Warnings**  
Do not use if you are now taking a prescription medication  
oxidase inhibitor (MAOI) (certain drugs for depression,  
psychiatric, or emotional conditions, or Parkinson's disease), or  
for 2 weeks after stopping the MAOI drug. If you do not know if  
your prescription drug contains an MAOI, ask a doctor or  
pharmacist before taking this product.

Ask a doctor before use if you have  
■ heart disease ■ high blood pressure  
■ thyroid disease ■ diabetes  
■ trouble urinating due to an enlarged prostate gland

When using this product do not exceed recommended dosage  
■ symptoms do not improve within 7 days or occur with a fever  
■ nervousness, dizziness, or sleeplessness occur

Stop use and ask a doctor if  
■ pregnant or breast-feeding, ask a health professional  
before use.  
Keep out of reach of children. In case of overdose, get medical  
help or contact a Poison Control Center right away  
(1-800-222-1222).

**Directions**  
adults and children  
■ take 2 tablets every 4 to 6 hours  
■ do not take more than 8 tablets  
in 24 hours

children ages  
6 to 11 years  
■ take 1 tablet every 4 to 6 hours  
■ do not take more than 4 tablets  
in 24 hours

children under  
6 years  
do not use this product in children  
under 6 years of age

**Other information**  
■ each tablet contains calcium 20 mg  
■ store at 20°-25°C (68°-77°F)  
■ do not use if blister unit is broken or torn

**Inactive ingredients** camauaba wax, dibasic calcium  
phosphate dihydrate, FD&C red no. 40 aluminum lake,  
hydroxyethylcellulose, magnesium stearate, microcrystalline cellulose,  
polyethylene glycol, polysorbate 80, silicon dioxide,  
titanium dioxide

**Questions or comments?** 1-888-423-0139



# Nasal Decongestant

PSEUDOEPHEDRINE HYDROCHLORIDE TABLETS, 30 mg

MAXIMUM STRENGTH



COMPARE TO SUDAFED® CONGESTION ACTIVE INGREDIENT\*

NON-DROWSY

# Nasal Decongestant

PSEUDOEPHEDRINE HYDROCHLORIDE TABLETS, 30 mg

MAXIMUM STRENGTH

- Sinus Pressure + Congestion
- Pseudoephedrine HCl

## 24 TABLETS

actual size



43262 88 C9

NDC 36800-432-62

# TOPCARE NASAL DECONGESTANT MAXIMUM STRENGTH NON DROWSY

pseudoephedrine hcl tablet, film coated

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:36800-432
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>PSEUDOEPHEDRINE HYDROCHLORIDE</b> (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)	PSEUDOEPHEDRINE HYDROCHLORIDE	30 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>CARNAUBA WAX</b> (UNII: R12CBM0EIZ)	
<b>DIBASIC CALCIUM PHOSPHATE DIHYDRATE</b> (UNII: O7TSZ97GEP)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	

## Product Characteristics

<b>Color</b>	RED	<b>Score</b>	no score
<b>Shape</b>	ROUND (convex)	<b>Size</b>	7mm
<b>Flavor</b>		<b>Imprint Code</b>	L432
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-432-62	24 in 1 CARTON	01/15/1988	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

2	NDC:36800-432-67	48 in 1 CARTON	02/15/1990	
2		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:36800-432-80	96 in 1 CARTON	12/26/2019	
3		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	01/15/1988	

**Labeler** - Topco Associates LLC (006935977)

Revised: 8/2022

Topco Associates LLC