

**TYLENOL EXTRA STRENGTH- acetaminophen tablet, coated**  
**Johnson & Johnson Consumer Inc., McNeil Consumer Healthcare Division**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**TYLENOL**

**Extra strength**

**Drug Facts**

**Active ingredient (in each gelcap)**

Acetaminophen 500 mg

**Purpose**

Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
  - headache
  - muscular aches
  - backache
  - minor pain of arthritis
  - the common cold
  - toothache
  - premenstrual and menstrual cramps
- temporarily reduces fever

**Warnings**

**Liver warning**

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

**Ask a doctor before use if you have** liver disease

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin

**Stop use and ask a doctor if**

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none"><li>▪ take 2 gelcaps every 6 hours while symptoms last</li><li>▪ do not take more than 6 gelcaps in 24 hours, unless directed by a doctor</li><li>▪ do not use for more than 10 days unless directed by a doctor</li></ul>
children under 12 years	ask a doctor

## **Other information**

- store between 20-25°C (68-77°F). Avoid high humidity.
- **do not use if carton is opened. Do not use if foil inner seal imprinted with "TYLENOL" is broken or missing**

## **Inactive ingredients**

benzyl alcohol, butylparaben, carboxymethylcellulose sodium, D&C yellow no. 10, edetate calcium disodium, FD&C blue no. 1, FD&C red no. 40, gelatin, hypromellose, iron oxide, magnesium stearate, methylparaben, modified starch, polyethylene glycol, polysorbate 80, powdered cellulose, pregelatinized starch, propylene glycol, propylparaben, red iron oxide, sodium lauryl sulfate, sodium propionate, sodium starch glycolate, titanium dioxide, yellow iron oxide

## **Questions or comments?**

call **1-877-895-3665** (toll-free) or **215-273-8755** (collect)

## **PRINCIPAL DISPLAY PANEL**

NDC 50580-488-24

Extra Strength  
TYLENOL<sup>®</sup>  
FOR ADULTS

Acetaminophen  
Pain Reliever  
Fever Reducer

RAPID RELEASE  
GELS

Actual Size

24 Gelcaps\*  
500 mg each

\*Gelatin-Coated Tablets

**TYLENOL**

Distributed by:  
**JOHNSON & JOHNSON**  
 McNeil Consumer  
 Healthcare Division  
 Fort Washington, PA 19034  
 USA  
 USD 500 849; USD 525 356;  
 US\$ 879 334; US\$ 067 029;  
 @J&JC12018  
**Contains No Aspirin**

OPEN HERE

30043733

How can we help?  
 1-877-895-3665

NDC 50580-488-24 **Extra Strength**  
**TYLENOL**<sup>®</sup>  
 FOR ADULTS

Acetaminophen Pain Reliever  
 Fever Reducer

**RAPID RELEASE**  
 GELS

\*Gelatin-Coated Tablets

Actual Size

24 Gelcaps\*  
 500 mg each

**Important:** Read all product information before using.  
 Keep this box for important information.

**Drug Facts**

**Active ingredient (in each gelcap)** Purpose  
 Acetaminophen 500 mg ..... Pain reliever and reducer

**Uses**

- temporarily relieves minor aches and pains due to:
  - headache
  - muscular aches
  - backache
  - minor pain of arthritis
  - the common cold
  - toothache
  - premenstrual and menstrual cramps
- temporarily reduces fever

**Warnings**

- Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take
  - more than 4,000 mg of acetaminophen in 24 hours
  - with other drugs containing acetaminophen
  - 3 or more alcoholic drinks every day while using this product
- Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:
  - skin redness
  - hives
  - rash
- If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen

**Drug Facts (continued)**

(prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
 If you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- your pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away; (1-800-322-0222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- do not take more than directed (see overdose warning)

**Drug Facts (continued)**

adults and children  
 take 2 gelcaps every 6 hours while symptoms last  
 do not take more than 6 gelcaps in 24 hours, unless directed by a doctor  
 do not use for more than 10 days unless directed by a doctor  
 children under 12 years  
 ask a doctor

**Other information**

- store between 20-25°C (68-77°F). Avoid high humidity.
- do not use if carton is opened. Do not use if inner seal imprinted with "TYLENOL" is broken or missing

**Inactive ingredients** benzyl alcohol, butylparaben, carboxymethylcellulose sodium, D&C yellow no. 10, croscarellin sodium, FD&C blue no. 1, FD&C red no. 40, gelatin, hypromellose, iron oxide, magnesium stearate, methylparaben, modified starch, polyethylene glycol, poly sorbate 80, powdered cellulose, pregelatinized starch, propylene glycol, propylparaben, red iron oxide, sodium lauryl sulfate, sodium propionate, sodium starch glycolate, titanium dioxide, yellow iron oxide

**Questions or comments?**

call 1-877-895-3665 (not free) or 215-273-8755 (collect)



30043733

How can we help?  
 1-877-895-3665

**TYLENOL EXTRA STRENGTH**

acetaminophen tablet, coated

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:50580-488
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
<b>BENZYL ALCOHOL</b> (UNII: LKG8494WBH)	
<b>BUTYLPARABEN</b> (UNII: 3QPI1U3FV8)	
<b>CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED FORM</b> (UNII: K679OBS311)	
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>EDETATE CALCIUM DISODIUM</b> (UNII: 25IH6R4SGF)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GELATIN, UNSPECIFIED</b> (UNII: 2G86QN327L)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>FERROSFERRIC OXIDE</b> (UNII: XM0M87F357)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>METHYLPARABEN</b> (UNII: A2I8C7HI9T)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)	
<b>POWDERED CELLULOSE</b> (UNII: SMD1X3XO9M)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>PROPYLPARABEN</b> (UNII: Z8IX2SC1OH)	
<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)	
<b>SODIUM LAURYL SULFATE</b> (UNII: 368GB5141J)	
<b>SODIUM PROPIONATE</b> (UNII: DK6Y9P42IN)	
<b>SODIUM STARCH GLYCOLATE TYPE A</b> (UNII: H8AV0SQX4D)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>FERRIC OXIDE YELLOW</b> (UNII: EX438O2MRT)	

**Product Characteristics**

<b>Color</b>	blue, gray, red	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	21mm
<b>Flavor</b>		<b>Imprint Code</b>	TY;500
<b>Contains</b>			

**Packaging**

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:50580-488-10	1 in 1 CARTON	01/16/2017	
1		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:50580-488-24	1 in 1 CARTON	01/16/2017	
2		24 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

3	NDC:50580-488-25	1 in 1 CARTON	01/16/2017	
3		225 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
4	NDC:50580-488-28	290 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	01/16/2017	
5	NDC:50580-488-01	50 in 1 TRAY	04/17/2017	12/31/2018
5		2 in 1 POUCH; Type 0: Not a Combination Product		
6	NDC:50580-488-52	1 in 1 CARTON	06/17/2019	
6		50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	01/16/2017	

**Labeler** - Johnson & Johnson Consumer Inc., McNeil Consumer Healthcare Division (878046358)

Revised: 5/2022

Johnson & Johnson Consumer Inc., McNeil Consumer Healthcare Division