

**CAPCOF- codeine phosphate, phenylephrine hydrochloride, chlorpheniramine maleate liquid**

**Capital Pharmaceutical, LLC**

-----

**CapCof**

**Drug Facts**

**Active ingredients**

***(in each 5 mL teaspoonful)***

Chlorpheniramine Maleate 2 mg

Codeine Phosphate\* 10 mg

\*(WARNING: May be habit forming)

Phenylephrine Hydrochloride 5 mg

**Purpose**

Antihistamine

Antitussive

Nasal Decongestant

**Uses**

Temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:

- runny nose
- sneezing
- itching of the nose or throat
- itchy, watery eyes
- cough due to minor throat and bronchial irritation
- nasal congestion
- reduces swelling of nasal passages

**Warnings**

**Do not exceed recommended dosage.**

**Do not use this product**

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a cough that lasts or is chronic such as occurs with smoking, asthma or emphysema
- a cough that occurs with too much phlegm (mucus)
- chronic pulmonary disease or shortness of breath, or children who are taking other drugs
- heart disease
- high blood pressure
- thyroid disease
- diabetes

**Ask a doctor or pharmacist before use if you are**

taking sedatives or tranquilizers.

**When using this product**

- excitability may occur, especially in children
- may cause marked drowsiness
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase the drowsiness effect
- be careful when driving a motor vehicle or operating machinery
- may cause or aggravate constipation

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- cough or nasal congestion persists for more than 1week, tends to recur, or is accompanied by a fever, rash or persistent headache. A persistent cough may be a sign of a serious condition.
- new symptoms occur

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

**Do not exceed recommended dosage.**

Adults and children 12 years of age and over:	2 teaspoonfuls (10 mL) every 4 hours, not to exceed 12 teaspoonfuls in 24 hours.
Children 6 to under 12 years of age:	1 teaspoonful (5 mL) every 4 hours, not to exceed 6 teaspoonfuls in 24 hours.
Children under 6 years of age:	Consult a doctor

A special measuring device should be used to give an accurate dose of this product to children under 6 years of age. Giving a higher dose than recommended by a doctor could result in serious side effects for your child.

### **Other information**

Store at 59° - 86°F (15° - 30°C)

### **Inactive ingredients**

Citric Acid, Cotton Candy Flavor, Glycerin, Propylene Glycol, Purified Water, Sodium Citrate, Sodium Saccharin, Sorbitol.

### **Questions? Comments?**

Call 1-614-638-4622.

CV

### **CapCof 16 fl oz. (473 mL) Bottle Label**



NDC 29978-420-16

# CapCof

Antihistamine • Antitussive  
• Nasal Decongestant

Each 5 mL (one teaspoonful) for oral administration contains:

- Chlorpheniramine Maleate ..... 2 mg
- Codeine Phosphate\* ..... 10 mg
- \*(WARNING-May be habit forming)
- Phenylephrine Hydrochloride ... 5 mg

Alcohol Free • Gluten Free  
Sugar Free • Dye Free  
Cotton Candy Flavor



**CAPITAL**  
pharmaceutical

16 fl oz. (473 mL)

Tamper evident by foil seal under cap. Do not use if foil seal is broken or missing.  
 Dispense in a tight, light-resistant container with a child-resistant cap.  
**THIS BOTTLE IS NOT TO BE DISPENSED TO THE CONSUMER.**  
 Manufactured for: Capital Pharmaceutical, LLC, Powell, OH 43065  
 Rev. 05/17

## Drug Facts Lift Here

### Active ingredients Purpose (in each 5 mL teaspoonful)

Chlorpheniramine Maleate 2 mg .....	Antihistamine
Codeine Phosphate* 10 mg .....	Antitussive
*(WARNING-May be habit forming)	
Phenylephrine Hydrochloride 5 mg ...	Nasal Decongestant

### Uses

Temporarily relieves these symptoms due to the common cold, hay fever (allergic rhinitis) or other upper respiratory allergies:

- runny nose ■ sneezing ■ itching of the nose or throat ■ itchy, watery eyes
- cough due to minor throat and bronchial irritation ■ nasal congestion
- reduces swelling of nasal passages

### Warnings

Do not exceed recommended dosage.

#### Do not use this product

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

#### Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma

**Drug Facts** (continued)

- trouble urinating due to an enlarged prostate gland
- a cough that lasts or is chronic such as occurs with smoking, asthma or emphysema
- a cough that occurs with too much phlegm (mucus)
- chronic pulmonary disease or shortness of breath, or children who are taking other drugs
- heart disease ■ high blood pressure
- thyroid disease ■ diabetes

**Ask a doctor or pharmacist before use if you are** taking sedatives or tranquilizers.

**When using this product**

- excitability may occur, especially in children
- may cause marked drowsiness
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase the drowsiness effect
- be careful when driving a motor vehicle or operating machinery
- may cause or aggravate constipation

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- cough or nasal congestion persists for more than 1 week, tends to recur, or is accompanied by a fever, rash or persistent headache. A persistent cough may be a sign of a serious condition.
- new symptoms occur

**If pregnant or breast-feeding,** ask a health professional before use.

**Drug Facts** (continued)

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

**Do not exceed recommended dosage.**

Adults and children 12 years of age and over:	2 teaspoonfuls (10 mL) every 4 hours, not to exceed 12 teaspoonfuls in 24 hours.
Children 6 to under 12 years of age:	1 teaspoonful (5 mL) every 4 hours, not to exceed 6 teaspoonfuls in 24 hours.
Children under 6 years of age:	Consult a doctor.

A special measuring device should be used to give an accurate dose of this product to children under 6 years of age. Giving a higher dose than recommended by a doctor could result in serious side effects for your child.

**Other information**

Store at 59° - 86° F (15° - 30° C)

**Inactive ingredients**

Citric Acid, Cotton Candy Flavor, Glycerin, Propylene Glycol, Purified Water, Sodium Citrate, Sodium Saccharin, Sorbitol.

**Drug Facts** (continued)**Questions? Comments?**

Call 1-614-638-4622

**CAPCOF**

codeine phosphate, phenylephrine hydrochloride, chlorpheniramine maleate liquid

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:29978-420
<b>Route of Administration</b>	ORAL	<b>DEA Schedule</b>	CV

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>CODEINE PHOSPHATE</b> (UNII: GSL05Y1MN6) (CODEINE ANHYDROUS - UNII:UX6OWY2V7J)	CODEINE PHOSPHATE	10 mg in 5 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 5 mL
<b>CHLORPHENIRAMINE MALEATE</b> (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg in 5 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>CITRIC ACID MONOHYDRATE</b> (UNII: 2968PHW8QP)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SODIUM CITRATE</b> (UNII: 1Q73Q2JULR)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SORBITOL</b> (UNII: 506T60A25R)	

## Product Characteristics

<b>Color</b>		<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	COTTON CANDY	<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:29978-420-16	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	10/08/2010	
2	NDC:29978-420-15	15 mL in 1 BOTTLE; Type 0: Not a Combination Product	10/08/2010	01/01/2021

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	10/08/2010	

**Labeler** - Capital Pharmaceutical, LLC (831545541)

Revised: 1/2024

Capital Pharmaceutical, LLC