

ACID REDUCER - omeprazole tablet, delayed release
Wal-Mart Stores, Inc.

Tips for Managing Heartburn

- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

Drug Facts

Active ingredient (in each tablet)

Omeprazole delayed-release tablet 20 mg (equivalent to 20.6 mg omeprazole magnesium USP)

Purpose

Acid reducer

Use

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert:

Do not use if you are allergic to omeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are taking:

- warfarin, clopidogrel or cilostazol (blood-thinning medicines)
- prescription antifungal or anti-yeast medicines
- diazepam (anxiety medicine)
- digoxin (heart medicine)
- tacrolimus or mycophenolate mofetil (immune system medicines)
- prescription antiretrovirals (medicines for HIV infection)
- methotrexate (arthritis medicine)

Stop use and ask a doctor if:

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- for adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours

14-Day Course of Treatment

- swallow 1 tablet with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 tablet a day
- do not use for more than 14 days unless directed by your doctor

- swallow whole. Do not chew or crush tablets.

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**
- children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20° to 25°C (68° to 77° F) and protect from moisture

Inactive ingredients

crospovidone, glyceryl monostearate, hydroxypropyl cellulose, hypromellose, magnesium stearate, methacrylic acid copolymer dispersion, microcrystalline cellulose, polyethylene glycol, polysorbate 80, red iron oxide, silicified microcrystalline cellulose, sodium hydroxide, sodium stearyl fumarate, sugar spheres [which contains liquid glucose, starch (maize) and sucrose], talc, titanium dioxide, triethyl citrate and yellow iron oxide.

Questions?

Call **1-855-274-4122**

**DISTRIBUTED BY: Walmart Inc.,
Bentonville, AR 72716**
PRODUCT OF INDIA
TS/DRUGS/22/2009

PACKAGE LABEL-PRINCIPAL DISPLAY PANEL - 20 mg (14 Tablet Bottle)

***equate*TM**
NDC 49035-983-05
Omeprazole
Delayed-Release
Tablets 20 mg
24 HR

ACID REDUCER

Treats **Frequent** Heartburn!

14 TABLETS

One 14-day course of treatment

- May take 1 to 4 days for full effect

Top Ply

Active ingredient (in each tablet)

Omeprazole delayed-release tablet 20 mg (equivalent to 20.6 mg omeprazole magnesium USP) **Purpose** Acid reducer

Use ■ treats frequent heartburn (occurs **2 or more** days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings Allergy alert: Do not use if you are allergic to omeprazole **Do not use if you have:** ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with **lightheadedness, sweating or dizziness** ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent **chest pain** These may be signs of a serious condition. See your doctor. **Ask a doctor before use if you have:** ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain **Ask a doctor or pharmacist before use if you are taking:** ■ warfarin, clopidogrel or cilostazol (blood-thinning medicines) ■ prescription antifungal or anti-yeast medicines ■ diazepam (anxiety medicine) ■ digoxin (heart medicine) ■ tacrolimus or mycophenolate mofetil (immune system medicines) ■ prescription antiretrovirals (medicines for HIV infection) ■ methotrexate (arthritis medicine) **Stop use and ask a doctor if:** ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days

Back of Top Ply

Bottom Ply

■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain **If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. **Directions** ■ for adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours **14-Day Course of Treatment** ■ swallow 1 tablet with a glass of water before eating in the morning ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor ■ swallow whole. Do not chew or crush tablets. **Repeated 14-Day Courses (if needed)** ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor ■ children under 18 years of age: ask a doctor, Heartburn in children may sometimes be caused by a serious condition. **Other information** ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20° to 25°C (68° to 77°F) and protect from moisture **Inactive ingredients** crosopovidone, glyceryl monostearate, hydroxypropyl cellulose, hypromellose, magnesium stearate, methacrylic acid copolymer dispersion, microcrystalline cellulose, polyethylene glycol, polysorbate 80, red iron oxide, silicified microcrystalline cellulose, sodium hydroxide, sodium stearyl fumarate, sugar spheres [which contains liquid glucose, starch (maize) and sucrose], talc, titanium dioxide, triethyl citrate and yellow iron oxide. **Questions?** Call 1-888-287-1915

PACKAGE LABEL-PRINCIPAL DISPLAY PANEL - 20 mg Blister Carton Label

3 PACK
equate™

NDC 49035-983-62

Compare to Prilosec OTC®

active ingredient*

Omeprazole

**Delayed-Release
Tablets 20 mg
24 HR**

ACID REDUCER

Treats **Frequent** Heartburn!

•May take 1 to 4 days for full effect

Actual Size

42

TABLETS

Three 14-day courses of treatment

Other information
 ■ read the directions and warnings before use ■ keep the carton, it contains important information.
 ■ store at 20° to 25°C (68° to 77°F) and protect from moisture.

Directions
 ■ for adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ it may take 1 to 4 days for full effect; some people get complete relief of symptoms within 24 hours
 14-Day Course of Treatment
 ■ swallow 1 tablet with a glass of water before eating in the morning ■ take every day for 14 days ■ do not take more than 1 tablet a day ■ do not use for more than 14 days unless directed by your doctor ■ swallow whole. Do not chew or crush tablets.
 Repeated 14-Day Courses (if needed)
 ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor ■ children under 18 years of age: ask a doctor. Heartburn in children may sometimes be caused by a serious condition.

Warnings
 Allergy alert: Do not use if you are allergic to omeprazole

Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain. These may be signs of a serious condition. See your doctor.
 Ask a doctor before use if you have: ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain
 Ask a doctor or pharmacist before use if you are taking:
 ■ warfarin, clopidogrel or clostazolol (blood-thinning medicines) ■ prescription antifungal or anti-yeast medicines ■ diazepam (anxiety medicine) ■ digoxin (heart medicine) ■ tacrolimus or cyclosporine (immunosuppressant medicines) ■ prescription antiretrovirals (medicines for HIV infection) ■ methotrexate (arthritis medicine)

Stop use and ask a doctor if:
 ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea ■ you develop a rash or joint pain

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Use
 ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Active ingredient (in each tablet)
 Omeprazole delayed-release tablet 20 mg (equivalent to 20.6 mg omeprazole magnesium USP).....Acid reducer

Purpose

Drug Facts

- Tips for Managing Heartburn**
- Do not lie flat or bend over after eating
 - Do not wear tight-fitting clothing around the stomach
 - Do not eat before bedtime
 - Raise the head of your bed
 - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol and certain fruits and vegetables
 - Eat slowly and avoid big meals
 - If overweight, lose weight
 - Quit smoking

TAMPER EVIDENT - Do not use if seal over bottle opening is broken or missing.

Satisfaction guaranteed - Or we'll replace it or give you your money back. For questions or comments or to report an undesired reaction or side effect, please call 1-888-287-1915.

DISTRIBUTED BY: Walmart Inc., Bentonville, AR 72716
 PRODUCT OF INDIA
 *This product is not manufactured or distributed by ASTRAZENCA PHARMACEUTICALS LP, owner of the registered trademark of Prilosec OTC®
 IS/DRUGS/22/2009



LM-3472

Lot: _____
 Expiry: _____

3 PACK
 THREE 14-DAY COURSES OF TREATMENT

equate™

Omeprazole
 Delayed-Release Tablets 20 mg

NDC 49035-983-62

Compare to Prilosec OTC® active ingredient*

Omeprazole
 Delayed-Release Tablets



ACID REDUCER

omeprazole tablet, delayed release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:49035-983
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OMEPRAZOLE MAGNESIUM (UNII: 426QFE7XLK) (OMEPRAZOLE - UNII:KG60484QX9)	OMEPRAZOLE	20 mg

Inactive Ingredients

Ingredient Name	Strength
CROSPVIDONE (12 MPA.S AT 5%) (UNII: 40UAA97IT9)	
GLYCERYL MONOSTEARATE (UNII: 230OU9XXE4)	
HYDROXYPROPYL CELLULOSE (90000 WAMW) (UNII: UKE75GEA7F)	
HYPROMELLOSE 2910 (5 MPA.S) (UNII: R75537T0T4)	

MAGNESIUM STEARATE (UNII: 70097M6I30)	
METHACRYLIC ACID - ETHYL ACRYLATE COPOLYMER (1:1) TYPE A (UNII: NX76LV5T8J)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL 4000 (UNII: 4R4HFI6D95)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
SODIUM STEARYL FUMARATE (UNII: 7CV7WJK4UI)	
DEXTROSE, UNSPECIFIED FORM (UNII: IY9XDZ35W2)	
STARCH, CORN (UNII: O8232NY3SJ)	
SUCROSE (UNII: C151H8M554)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
HYDROXYPROPYL CELLULOSE (45000 WAMW) (UNII: 8VAB711C5E)	
HYPROMELLOSE 2910 (6 MPA.S) (UNII: 0WZ8WG20P6)	

Product Characteristics

Color	PINK	Score	no score
Shape	RECTANGLE (Oblong)	Size	14mm
Flavor		Imprint Code	Z;69
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49035-983-62	3 in 1 CARTON	06/06/2018	
1	NDC:49035-983-05	14 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA206877	06/06/2018	

Labeler - Wal-Mart Stores, Inc. (051957769)

Registrant - Aurohealth LLC (078728447)

Establishment

Name	Address	ID/FEI	Business Operations
Aurobindo Pharma Limited		650381903	ANALYSIS(49035-983) , MANUFACTURE(49035-983)

Revised: 6/2024

Wal-Mart Stores, Inc.