

SEVERE SINUS CONGESTION AND PAIN- acetaminophen, guaifenesin, and phenylephrine hydrochloride tablet, coated

Chain Drug Marketing Association

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

1119 - QCH - 2014-1028

Drug Facts

<i>Active ingredients (in each caplet)</i>	<i>Purpose</i>
Acetaminophen 325 mg	Pain reliever/fever reducer
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- for the temporary relief of:
 - sinus congestion and pressure
 - headache
 - nasal congestion
 - minor aches and pains
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive
- temporarily relieves nasal congestion due to the common cold, and hay fever or other upper respiratory allergies
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure

- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dosage

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none"> ▪ take 2 caplets every 4 hours ▪ swallow whole – do not crush, chew, or dissolve ▪ do not take more than 10 caplets in 24 hours
children under 12 years	<ul style="list-style-type: none"> ▪ ask a doctor

Other information

- store between 20-25°C (68-77°F) in a dry place
- retain carton for complete product information

Inactive ingredients

acesulfame potassium, colloidal silicon dioxide, corn starch, croscarmellose sodium, crospovidone, flavor, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, propylene glycol, stearic acid, talc, titanium dioxide

PRINCIPAL DISPLAY PANEL

NDC 63868-071-24

QUALITY CHOICE

†Compare to TYLENOL® Sinus Congestion & Pain Severe active ingredients

Daytime, Non-Drowsy

Severe Sinus Congestion & Pain

For Adults

Pain Reliever / Fever Reducer, Nasal Decongestant, Expectorant

Acetaminophen, Phenylephrine HCl, Guaifenesin

For Relief of:

Sinus Headache / Sinus Pressure

Nasal Congestion

Clears Mucus, Chest Congestion

24 Cool Taste Caplets

SEE NEW WARNINGS INFORMATION & DIRECTIONS

Non-Drowsy Daytime

NDC

NDC

NDC

Drug Facts (continued)

- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours.

- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- high blood pressure
- liver disease
- thyroid disease
- trouble urinating due to an enlarged prostate gland
- chronic bronchitis, or emphysema

persistent or chronic cough such as occurs with smoking, asthma.

Directions

- adults and children 12 years and over
 - take 2 caplets every 4 hours
 - swallow whole – do not crush, chew, or dissolve
 - do not take more than 10 caplets in 24 hours
- children under 12 years
 - ask a doctor

do not take more than directed (see overdose warning)

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Keep out of reach of children.

If pregnant or breast-feeding, ask a health professional before use. These could be signs of a serious condition.

- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts more than 7 days
- fever gets worse or lasts more than 3 days
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- nervousness, dizziness, or sleeplessness occur

Stop use and ask a doctor if

When using this product do not exceed recommended dosage

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Drug Facts (continued)

Drug Facts (continued)

Drug Facts

Active ingredients (in each caplet) Purpose

- Acetaminophen 325 mg: Pain reliever/fever reducer
- Guaifenesin 200 mg: Expectorant
- Phenylephrine HCl 5 mg: Nasal decongestant

Uses for the temporary relief of:

- sinus congestion and pressure
- headache
- nasal congestion
- minor aches and pains

Drug Facts (continued)

- temporarily relieves nasal congestion due to the common cold, and may fever or other upper respiratory allergies

6 35515 95990 3

NDC 63868-071-24

QC QUALITY CHOICE

*Compare to **TYLENOL**[®] Sinus Congestion & Pain Severe active ingredients

Daytime, Non-Drowsy Severe Sinus Congestion & Pain For Adults

Pain Reliever | Fever Reducer, Nasal Decongestant, Expectorant

Acetaminophen, Phenylephrine HCl, Guaifenesin

For Relief of:

- Sinus Headache | Sinus Pressure
- Nasal Congestion
- Clears Mucus, Chest Congestion

24 Cool Taste Caplets

SEE NEW WARNINGS INFORMATION & DIRECTIONS

QC QUALITY CHOICE

SALES

Distributed by C.D.M.A., Inc.
43157 W. Nine Mile
Novi, MI 48376-0995
www.qualitychoice.com
Questions: 248-449-9300

F111905QCH_R0

†This product is not manufactured or distributed by McKel Consumer Healthcare, distributor of Tylenol® Sinus Congestion and Pain Severe.

DO NOT USE IF BLISTER UNITS ARE TORN OR BROKEN

Drug Facts (continued)

Other Information

- store between 20-25°C (68-77°F) in a dry place
- retain carton for complete product information

Drug Facts (continued)

Inactive ingredients

- silicon dioxide, corn starch, croscarmellose sodium, crospovidone, acsulfame potassium, colloidal
- flavor, magnesium stearate, malto-dextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, propylene glycol, stearic acid, talc, titanium dioxide

acetaminophen, guaifenesin, and phenylephrine hydrochloride tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63868-071
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
CROSPVIDONE (UNII: 68401960MK)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL (UNII: 532B59J990)	
POVIDONE (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3SJ)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	white	Score	no score
Shape	OVAL	Size	19mm
Flavor	MINT	Imprint Code	AAA;1119
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63868-071-24	2 in 1 CARTON	11/13/2009	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	11/13/2009	

Labeler - Chain Drug Marketing Association (011920774)

Revised: 12/2018

Chain Drug Marketing Association