

**LORATADINE AND PSEUDOEPHEDRINE SULFATE- loratadine and pseudoephedrine sulfate tablet, film coated, extended release**  
**Chain Drug Marketing Association Inc.**

-----  
**Loratadine and Pseudoephedrine Sulfate**

***Drug Facts***

<b><i>Active ingredients (in each tablet)</i></b>	<b><i>Purpose</i></b>
Loratadine, USP 10 mg	Antihistamine
Pseudoephedrine sulfate, USP 240 mg	Nasal decongestant

**Uses**

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
  - sneezing
  - itchy, watery eyes
  - runny nose
  - itching of the nose or throat
- reduces swelling of nasal passages
- temporarily relieves sinus congestion and pressure
- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- temporarily restores freer breathing through the nose

**Warnings**

**Do not use**

- if you have ever had an allergic reaction to this product or any of its ingredients
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- heart disease
- thyroid disease
- high blood pressure
- diabetes
- trouble urinating due to an enlarged prostate gland
- liver or kidney disease. Your doctor should determine if you need a different dose.

**When using this product do not take more than directed.** Taking more than directed may cause drowsiness.

**Stop use and ask a doctor if**

- an allergic reactions to this product occurs. Seek medical help right away.
- symptoms do not improve within 7 days or are accompanied by a fever
- nervousness, dizziness or sleeplessness occurs

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### Directions

- do not divide, crush, chew or dissolve the tablet

adults and children 12 years and over	1 tablet daily with a full glass of water; not more than 1 tablet in 24 hours
children under 12 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

### Other information

- sodium:** contains 10 mg/tablet
- calcium:** contains 25 mg/tablet
- TAMPER EVIDENT: DO NOT USE IF BLISTER UNITS ARE TORN, BROKEN OR SHOW ANY SIGNS OF TAMPERING.**
- store between 20° C to 25° C (68° F to 77° F)
- protect from light and store in a dry place

### Inactive ingredients

calcium carbonate, colloidal silicon dioxide, hydroxypropyl cellulose, hypromellose, iron oxide black, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, propylene glycol, shellac glaze, sodium alginate, sodium citrate, talc and titanium dioxide

### Questions?

call **1-800-406-7984**

Distributed by C.D.M.A., Inc.©  
43157 W. Nine Mile  
Novi, MI 48376-0995

### PRINCIPAL DISPLAY PANEL - 10 Tablet Blister Pack Carton

QUALITY®  
CHOICE

NDC 63868-154-10

†Compare to  
Active Ingredients in  
CLARITIN-D® 24 Hour

Original Prescription Strength  
Loratadine-D  
Loratadine, USP 10 mg | Antihistamine

Pseudoephedrine Sulfate, USP 240 mg  
Nasal Decongestant

Allergy Relief and Nasal Decongestant

Indoor & Outdoor Allergies

Relief of:

Nasal and Sinus Congestion Due to Colds or Allergies

Sneezing | Runny Nose | Itchy, Watery Eyes

Itchy Throat or Nose Due to Allergies

24 Hour Formula | Non-Drowsy\*

10 Extended-Release Tablets

\*When taken as directed.

See Drug Facts Panel.

**Drug Facts (continued)**

- reduces swelling of nasal passages
- temporarily relieves sinus congestion and pressure
- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- temporarily restores free breathing through the nose

**Warnings**

- Do not use
  - if you have ever had an allergic reaction to this product or any of its ingredients
  - if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or for 2 weeks after stopping Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- heart disease
- thyroid disease
- high blood pressure
- diabetes
- trouble urinating due to an enlarged prostate gland
- liver or kidney disease. Your doctor should determine if you need a different dose.

**When using this product do not take more than directed.** Taking more than directed may cause drowsiness.

**Drug Facts (continued)**

**Stop use and ask a doctor if**

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve within 7 days or are accompanied by a fever
- neurosis, dizziness or sleeplessness occurs

**If pregnant or breast-feeding, ask a health professional before use.**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222).

**Directions**

- do not divide, crush, chew or dissolve the tablet
- adults and children: 1 tablet daily with a full glass of water; not more than 1 tablet
- children under 12 years of age: ask a doctor
- consumers with liver or kidney disease: ask a doctor

**Other information**

- sodium: contains 10 mg/tablet
- calcium: contains 25 mg/tablet
- TAMPER EVIDENT: DO NOT USE IF BLISTER UNITS ARE TORN, BROKEN OR SHOW ANY SIGNS OF TAMPERING.
- store between 20° C to 25° C (68° F to 77° F)
- protect from light and store in a dry place

**Drug Facts**

**Active ingredients (in each tablet)**

Loratadine, USP 10 mg  
Pseudoephedrine sulfate, USP 240 mg

**Purpose**

Antihistamine  
Nasal decongestant

**Uses**

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: sneezing, itchy, watery eyes, runny nose, itching of the nose or throat



**QC**  
QUALITY CHOICE

NDC 93868-154-10

Compare to  
**Active Ingredients in  
CLARITIN-D<sup>®</sup> 24 Hour**

**Original Prescription Strength**

# Loratadine-D

**Loratadine, USP 10 mg | Antihistamine  
Pseudoephedrine Sulfate, USP 240 mg  
Nasal Decongestant**

**Allergy Relief and Nasal Decongestant**

**Indoor & Outdoor Allergies**

**Relief of:**

Nasal and Sinus Congestion Due to Colds or Allergies  
Sneezing | Runny Nose | Itchy, Watery Eyes  
Itchy Throat or Nose Due to Allergies

**24 Hour Formula | Non-Drowsy\***



**10 Extended-Release Tablets**

\*When taken as directed. See Drug Facts Panel.


**Drug Facts (continued)**

**Inactive ingredients** calcium carbonate, colloidal silicon dioxide, hydroxypropyl cellulose, hypromellose, iron oxide black, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, propylene glycol, shellac glaze, sodium alginate, sodium citrate, talc and titanium dioxide

**Questions?** call 1-800-406-7984

**Keep the carton. It contains important information.**  
See end panel for expiration date.

R0114



100% SATISFACTION  
GUARANTEED

Distributed by C.D.M.A., Inc.<sup>®</sup>  
43157 W. Nine Mile  
Novi, MI 48376-0995  
www.qualitychoices.com  
Questions: 248-449-9300

3 5 5 1 5 9 5 3 7 3

Expiration Date:  
Batch No.  
**Non Varnish Area**



# LORATADINE AND PSEUDOEPHEDRINE SULFATE

loratadine and pseudoephedrine sulfate tablet, film coated, extended release

## Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63868-154
Route of Administration	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LORATADINE (UNII: 7AJO3BO7QN) (LORATADINE - UNII:7AJO3BO7QN)	LORATADINE	10 mg
PSEUDOEPHEDRINE SULFATE (UNII: Y9DL7QPE6B) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)	PSEUDOEPHEDRINE SULFATE	240 mg

## Inactive Ingredients

Ingredient Name	Strength
CALCIUM CARBONATE (UNII: H0G9379FGK)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
HYDROXYPROPYL CELLULOSE (120000 MW) (UNII: RFW2ET671P)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
FERROSO FERRIC OXIDE (UNII: XM0M87F357)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3SJ)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SHELLAC (UNII: 46N107B71O)	
SODIUM ALGINATE (UNII: C269C4G2ZQ)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

## Product Characteristics

Color	white (White to Off-White)	Score	no score
Shape	CAPSULE	Size	17mm
Flavor		Imprint Code	RX724
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
---	-----------	---------------------	----------------------	--------------------

1	NDC:63868-154-10	10 in 1 BLISTER PACK; Type 0: Not a Combination Product	11/17/2004	
<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>		<b>Marketing Start Date</b>	<b>Marketing End Date</b>
ANDA	ANDA076557		11/17/2004	

**Labeler** - Chain Drug Marketing Association Inc. (011920774)

**Registrant** - Ranbaxy Pharmaceuticals Inc. (937890044)

<b>Establishment</b>			
<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
Ohm Laboratories Inc.		051565745	manufacture(63868-154)

Revised: 8/2018

Chain Drug Marketing Association Inc.