

GENTLE LAXATIVE- bisacodyl tablet, delayed release
United Natural Foods, Inc. dba UNFI

Equaline 44-607

Active ingredient (in each tablet)

Bisacodyl USP, 5 mg

Purpose

Stimulant laxative

Uses

- for relief of occasional constipation and irregularity
- this product generally produces bowel movement in 6 to 12 hours

Warnings

Do not use

if you cannot swallow without chewing.

Ask a doctor before use if you have

- stomach pain, nausea or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

When using this product

- do not use within 1 hour after taking an antacid or milk
- do not chew or crush tablet(s)
- you may have stomach discomfort, faintness and cramps

Stop use and ask a doctor if

- you have rectal bleeding or fail to have a bowel movement after use of a laxative. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- take with a glass of water

adults and children 12 years and over	take 1 to 3 tablets in a single daily dose
children 6 to under 12 years	take 1 tablet in a single daily dose
children under 6 years	ask a doctor

Other information

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- avoid excessive humidity
- see end flap for expiration date and lot number

Inactive ingredients

acacia, ammonium hydroxide, calcium carbonate, carnauba wax, colloidal anhydrous silica, corn starch, D&C red #27 aluminum lake, FD&C blue #2 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, iron oxide black, lactose anhydrous, magnesium stearate, methylparaben, polydextrose, polyethylene glycol, polyvinyl acetate phthalate, povidone, propylene glycol, propylparaben, shellac glaze, simethicone, sodium alginate, sodium benzoate, sodium bicarbonate, stearic acid, sucrose, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments?

1-855-423-2630

Principal Display Panel

EQUALINE®

**compare to
Dulcolax Pink®**
active ingredient*

NDC 41163-607-56

**gentle
laxative**

bisacodyl USP 5 mg
(stimulant laxative)

- *gentle, dependable
constipation relief*
- *enteric coated*

25 tablets

actual size

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS
OPENED OR IF BLISTER UNIT IS TORN, BROKEN
OR SHOWS ANY SIGNS OF TAMPERING**

*This product is not manufactured or
distributed by Sanofi-Aventis Deutschland
GMBH, owner of the registered trademark
Dulcolax Pink®.

50844 REV0119D60756

**DISTRIBUTED BY UNFI
PROVIDENCE, RI 02908 USA
855-423-2630**

**100% *Quality*
GUARANTEED**

gentle, dependable constipation relief

EQUALINE[®]

compare to
Dulcolax Pink[®]
active ingredient*

gentle laxative

EQUALINE[®]

compare to
Dulcolax Pink[®]
active ingredient*
NDC 41163-607-56

gentle
laxative
bisacodyl USP 5 mg
(stimulant laxative)

- gentle, dependable constipation relief
- enteric coated

25 tablets

actual size

LEBG29615

50844 REV0119D60756

*This product is not manufactured or distributed by Sanofi-Aventis Deutschland GmbH, owner of the registered trademark Dulcolax Pink[®].

DISTRIBUTED BY UNFI
PROVIDENCE, RI 02908 USA
855-423-2630

100% Quality GUARANTEED

Drug Facts (continued)

Questions or comments? 1-855-423-2630

B-0019E-607-56-R
REV0119D60756

gentle lax

EQUALINE

Inactive ingredients acacia, ammonium hydroxide, calcium carbonate, carnauba wax, colloidal anhydrous silica, corn starch, D & C red #27 aluminum lake, FD & C blue #2 aluminum lake, FD & C yellow #6 aluminum lake, hypromellose, iron oxide black, lactose, anhydrous, magnesium stearate, methylparaben, polydextrose, polyethylene glycol, polyvinyl acetate phthalate, povidone, propylene glycol, propylparaben, shellac glaze, simethicone, sodium alginate, sodium benzoate, sodium bicarbonate, stearic acid, sucrose, talc, titanium dioxide, triacetin, triethyl citrate

or milk ■ do not chew or crush tablet(s) ■ you may have stomach discomfort, flatness and cramps
Stop use and ask a doctor if
■ you have rectal bleeding or fail to have a bowel movement after use of a laxative. These could be signs of a serious condition.
■ you need to use a laxative for more than 1 week
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.



kative

compare to
Dulcolax Pink®
active ingredient*

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

<p style="text-align: center; font-weight: bold; color: red;">Drug Facts (continued)</p> <p>Directions ■ take with a glass of water</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 50%; padding: 2px;">adults and children take 1 to 3 tablets in a single daily dose</td> <td style="width: 50%; padding: 2px;">children 6 to under 12 years take 1 tablet in a single daily dose</td> </tr> <tr> <td colspan="2" style="padding: 2px;">ask a doctor</td> </tr> </table>	adults and children take 1 to 3 tablets in a single daily dose	children 6 to under 12 years take 1 tablet in a single daily dose	ask a doctor		<p style="text-align: center; font-weight: bold;">Other Information</p> <p>■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN</p> <p>■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)</p> <p>■ avoid excessive humidity</p> <p>■ see end flap for expiration date and lot number</p>	<p style="text-align: center; font-weight: bold;">Drug Facts</p> <p style="text-align: center; font-weight: bold;">Active ingredient (in each tablet)</p> <p>Bisacodyl USP, 5 mgStimulant laxative</p> <p style="text-align: center; font-weight: bold;">Purpose</p> <p>■ for relief of occasional constipation and irregularly this product generally produces bowel movement in 6 to 12 hours</p> <p style="text-align: center; font-weight: bold;">Uses</p> <p>■ do not use within 1 hour after taking an antacid</p> <p style="text-align: center; font-weight: bold;">Warnings</p> <p>Do not use if you cannot swallow without chewing.</p> <p>Ask a doctor before use if you have stomach pain, nausea or vomiting</p> <p>■ a sudden change in bowel habits that lasts more than 2 weeks</p> <p>When using this product</p>
adults and children take 1 to 3 tablets in a single daily dose	children 6 to under 12 years take 1 tablet in a single daily dose					
ask a doctor						

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

Equaline 44-607

GENTLE LAXATIVE

bisacodyl tablet, delayed release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41163-607
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISACODYL (UNII: 10X0709Y6I) (DEACETYLBISACODYL - UNII:R09078E41Y)	BISACODYL	5 mg

Inactive Ingredients

Ingredient Name	Strength
ACACIA (UNII: 5C5403N26O)	
AMMONIA (UNII: 5138Q19F1X)	
CALCIUM CARBONATE (UNII: H0G9379FGK)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
D&C RED NO. 27 ALUMINUM LAKE (UNII: ZK64F7XSTX)	
FD&C BLUE NO. 2 ALUMINUM LAKE (UNII: 4AQJ3LG584)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
FERROSO FERRIC OXIDE (UNII: XM0M87F357)	
ANHYDROUS LACTOSE (UNII: 3S5Y5LH9PMK)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
METHYL PARABEN (UNII: A2I8C7HI9T)	

POLYDEXTROSE (UNII: VH2XOU12IE)
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)
Polyvinyl Acetate Phthalate (UNII: 58QVG85GW3)
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)
PROPYLPARABEN (UNII: Z8IX2SC1OH)
SHELLAC (UNII: 46N107B710)
DIMETHICONE (UNII: 92RU3N3Y1O)
WATER (UNII: 059QF0KO0R)
SODIUM ALGINATE (UNII: C269C4G2ZQ)
SODIUM BENZOATE (UNII: OJ245FE5EU)
SODIUM BICARBONATE (UNII: 8MDF5V39QO)
STEARIC ACID (UNII: 4ELV7Z65AP)
SUCROSE (UNII: C151H8M554)
TALC (UNII: 7SEV7J4R1U)
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)
TRIACETIN (UNII: XHX3C3X673)
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)

Product Characteristics

Color	pink	Score	no score
Shape	ROUND	Size	6mm
Flavor		Imprint Code	5
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41163-607-56	1 in 1 CARTON	04/01/2013	
1		25 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:41163-607-52	5 in 1 CARTON	04/01/2013	
2		18 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	505G(a)(3)	04/01/2013	

Labeler - United Natural Foods, Inc. dba UNFI (943556183)

Establishment

Name	Address	ID/FEI	Business Operations
------	---------	--------	---------------------

LNK International, Inc.		038154464	pack(41163-607)
-------------------------	--	-----------	-----------------

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(41163-607) , pack(41163-607)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(41163-607)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(41163-607)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(41163-607)

Revised: 1/2024

United Natural Foods, Inc. dba UNFI