

CEFAZOLIN- cefazolin injection, powder, for solution
Hikma Pharmaceuticals USA Inc.

CEFAZOLIN FOR INJECTION, USP


Rx Only

To reduce the development of drug-resistant bacteria and maintain the effectiveness of Cefazolin for Injection and other antibacterial drugs, Cefazolin for Injection should be used only to treat or prevent infections that are proven or strongly suspected to be caused by bacteria.

DESCRIPTION

Cefazolin for Injection, USP is a semi-synthetic cephalosporin for parenteral administration. It is the sodium salt of 3-[[[5-methyl-1,3,4-thiadiazol-2-yl]thio]-methyl]-8-oxo-7-[2-(1H-tetrazol-1-yl) acetamido]-5-thia-1-azabicyclo [4.2.0]oct-2-ene-2-carboxylic acid. The molecular formula is $C_{14}H_{13}N_8NaO_4S_3$ and molecular weight is 476.49.

Structural Formula:



Each vial contains 48 mg of sodium/1 gram of cefazolin sodium.

Cefazolin for Injection, USP is white to off-white crystalline powder, supplied in vials equivalent to 500 mg or 1 gram of cefazolin.

CLINICAL PHARMACOLOGY

After intramuscular administration of Cefazolin for Injection to normal volunteers, the mean serum concentrations were 37 mcg/mL at 1 hour and 3 mcg/mL at 8 hours following a 500 mg dose, and 64 mcg/mL at 1 hour and 7 mcg/mL at 8 hours following a 1 gram dose.

Studies have shown that following intravenous administration of Cefazolin for Injection to normal volunteers, mean serum concentrations peaked at approximately 185 mcg/mL and were approximately 4 mcg/mL at 8 hours for a 1 gram dose.

The serum half-life for Cefazolin for Injection is approximately 1.8 hours following IV administration and approximately 2 hours following IM administration.

In a study (using normal volunteers) of constant intravenous infusion with dosages of 3.5 mg/kg for 1 hour (approximately 250 mg) and 1.5 mg/kg the next 2 hours (approximately 100 mg), Cefazolin for Injection produced a steady serum level at the third hour of approximately 28 mcg/mL.

Studies in patients hospitalized with infections indicate that Cefazolin for Injection produces mean peak serum levels approximately equivalent to those seen in normal volunteers.

Bile levels in patients without obstructive biliary disease can reach or exceed serum levels by up to five times; however, in patients with obstructive biliary disease, bile levels of Cefazolin for Injection are considerably lower than serum levels (< 1 mcg/mL).

In synovial fluid, the level of Cefazolin for Injection becomes comparable to that reached in serum at about 4 hours after drug administration. Studies of cord blood show prompt transfer of Cefazolin for Injection across the placenta. Cefazolin for Injection is present in very low concentrations in the milk of nursing mothers.

Cefazolin for Injection is excreted unchanged in the urine. In the first 6 hours approximately 60% of the drug is excreted in the urine and this increases to 70% to 80% within 24 hours. Cefazolin for Injection achieves peak urine concentrations of approximately 2,400 mcg/mL and 4,000 mcg/mL respectively following 500 mg and 1 gram intramuscular doses.

In patients undergoing peritoneal dialysis (2 L/hr.), Cefazolin for Injection produced mean serum levels of approximately 10 and 30 mcg/mL after 24 hours' instillation of a dialyzing solution containing 50 mg/L and 150 mg/L, respectively. Mean peak levels were 29 mcg/mL (range 13 to 44 mcg/mL) with 50 mg/L (3 patients), and 72 mcg/mL (range 26 to 142 mcg/mL) with 150 mg/L (6 patients). Intraperitoneal administration of Cefazolin for Injection is usually well tolerated.

Controlled studies on adult normal volunteers, receiving 1 gram 4 times a day for 10 days, monitoring CBC, SGOT, SGPT, bilirubin, alkaline phosphatase, BUN, creatinine and urinalysis, indicated no clinically significant changes attributed to Cefazolin for Injection.

Microbiology

Mechanism of Action

Cefazolin is a bactericidal agent that acts by inhibition of bacterial cell wall synthesis.

Resistance

Predominant mechanisms of bacterial resistance to cephalosporins include the presence of extended-spectrum beta-lactamases and enzymatic hydrolysis.

Antimicrobial Activity

Cefazolin has been shown to be active against most isolates of the following microorganisms, both *in vitro* and in clinical infections as described in the **INDICATIONS AND USAGE(1)** section:

Gram-Positive Bacteria:

Staphylococcus aureus

Staphylococcus epidermidis

Streptococcus agalactiae

Streptococcus pneumoniae

Streptococcus pyogenes

Methicillin-resistant staphylococci are uniformly resistant to cefazolin.

Gram-Negative Bacteria:

Escherichia coli

Proteus mirabilis

Most isolates of indole positive *Proteus* (*Proteus vulgaris*), *Enterobacter* spp., *Morganella morganii*, *Providencia rettgeri*, *Serratia* spp., and *Pseudomonas* spp. are resistant to cefazolin.

Susceptibility Test Methods

For specific information regarding susceptibility test interpretive criteria and associated test methods and quality control standards recognized by FDA for this drug, please see: <https://www.fda.gov/STIC>.

INDICATIONS AND USAGE

Cefazolin for Injection, USP is indicated in the treatment of the following serious infections due to susceptible organisms:

Respiratory Tract Infections: Due to *S. pneumoniae*, *Klebsiella* species, *H. influenzae*, *S. aureus* (penicillin-sensitive and penicillin-resistant), and group A beta-hemolytic streptococci. Injectable benzathine penicillin is considered to be the drug of choice in treatment and prevention of streptococcal infections, including the prophylaxis of rheumatic fever.

Cefazolin for Injection is effective in the eradication of streptococci from the nasopharynx; however, data establishing the efficacy of Cefazolin for Injection in the subsequent prevention of rheumatic fever are not available at present.

Urinary Tract Infections: Due to *E. coli*, *P. mirabilis*, *Klebsiella* species, and some strains of enterobacter and enterococci.

Skin and Skin Structure Infections: Due to *S. aureus* (penicillin-sensitive and penicillin-resistant), group A beta-hemolytic streptococci, and other strains of streptococci.

Biliary Tract Infections: Due to *E. coli*, various strains of streptococci, *P. mirabilis*, *Klebsiella* species, and *S. aureus*.

Bone and Joint Infections: Due to *S. aureus*.

Genital Infections: (i.e., prostatitis, epididymitis) due to *E. coli*, *P. mirabilis*, *Klebsiella* species, and some strains of enterococci.

Septicemia: Due to *S. pneumoniae*, *S. aureus* (penicillin-sensitive and penicillin-resistant), *P. mirabilis*, *E. coli* and *Klebsiella* species.

Endocarditis: Due to *S. aureus* (penicillin-sensitive and penicillin-resistant) and group A beta-hemolytic streptococci.

Perioperative Prophylaxis: The prophylactic administration of Cefazolin for Injection preoperatively, intraoperatively and postoperatively may reduce the incidence of certain postoperative infections in patients undergoing surgical procedures which are classified as contaminated or potentially contaminated (e.g., vaginal hysterectomy, and cholecystectomy in high-risk patients such as those older than 70 years, with acute cholecystitis, obstructive jaundice, or common duct bile stones).

The perioperative use of Cefazolin for Injection may also be effective in surgical patients in whom infection at the operative site would present a serious risk (e.g., during open-heart surgery and prosthetic arthroplasty).

The prophylactic administration of Cefazolin for Injection should usually be discontinued within a 24 hour period after the surgical procedure. In surgery where the occurrence of infection may be particularly devastating (e.g., open-heart surgery and prosthetic arthroplasty), the prophylactic administration of Cefazolin for Injection may be continued for 3 to 5 days following the completion of surgery.

If there are signs of infection, specimens for cultures should be obtained for the identification of the causative organism so that appropriate therapy may be instituted (see **DOSAGE AND ADMINISTRATION**).

To reduce the development of drug-resistant bacteria and maintain the effectiveness of Cefazolin for Injection, USP and other antibacterial drugs, Cefazolin for Injection, USP should be used only to treat or prevent infections that are proven or strongly suspected to be caused by susceptible bacteria. When culture and susceptibility information are available, they should be considered in selecting or modifying antibacterial therapy. In the absence of such data, local epidemiology and susceptibility patterns may contribute to the empiric selection of therapy.

CONTRAINDICATIONS

Cefazolin for Injection IS CONTRAINDICATED IN PATIENTS WITH KNOWN ALLERGY TO THE CEPHALOSPORIN GROUP OF ANTIBIOTICS.

WARNINGS

BEFORE THERAPY WITH CEFAZOLIN FOR INJECTION IS INSTITUTED, CAREFUL INQUIRY SHOULD BE MADE TO DETERMINE WHETHER THE PATIENT HAS HAD PREVIOUS HYPERSENSITIVITY REACTIONS TO CEFAZOLIN, CEPHALOSPORINS, PENICILLINS, OR OTHER DRUGS. IF THIS PRODUCT IS GIVEN TO PENICILLIN-SENSITIVE PATIENTS, CAUTION SHOULD BE EXERCISED BECAUSE CROSS-HYPERSENSITIVITY AMONG BETA-LACTAM ANTIBIOTICS HAS BEEN CLEARLY DOCUMENTED AND MAY OCCUR IN UP TO 10% OF PATIENTS WITH A HISTORY OF PENICILLIN ALLERGY. IF AN ALLERGIC REACTION TO CEFAZOLIN FOR INJECTION OCCURS, DISCONTINUE TREATMENT WITH THE DRUG. SERIOUS ACUTE HYPERSENSITIVITY REACTIONS MAY REQUIRE TREATMENT WITH EPINEPHRINE AND OTHER EMERGENCY MEASURES, INCLUDING OXYGEN, IV FLUIDS, IV ANTIHISTAMINES, CORTICOSTEROIDS, PRESSOR AMINES, AND AIRWAY MANAGEMENT, AS CLINICALLY INDICATED.

Pseudomembranous colitis has been reported with nearly all antibacterial agents, including cefazolin, and may range in severity from mild to life-threatening. Therefore, it is important to consider this diagnosis in patients who present with diarrhea subsequent to the administration of antibacterial agents.

Treatment with antibacterial agents alters the normal flora of the colon and may permit overgrowth of clostridia. Studies indicate that a toxin produced by *Clostridium difficile* is a primary cause of “antibiotic-associated colitis”.

After the diagnosis of pseudomembranous colitis has been established, therapeutic measures should be initiated. Mild cases of pseudomembranous colitis usually respond to drug discontinuation alone. In moderate to severe cases, consideration should be given to management with fluids and electrolytes, protein supplementation, and treatment with an oral antibacterial drug clinically effective against *C. difficile* colitis.

PRECAUTIONS

General

Prolonged use of Cefazolin for Injection may result in the overgrowth of nonsusceptible organisms. Careful clinical observation of the patient is essential.

When Cefazolin for Injection is administered to patients with low urinary output because of impaired renal function, lower daily dosage is required (see **DOSAGE AND ADMINISTRATION**).

As with other β -lactam antibiotics, seizures may occur if inappropriately high doses are administered to patients with impaired renal function (see **DOSAGE AND ADMINISTRATION**).

Cefazolin for Injection, as with all cephalosporins, should be prescribed with caution in individuals with a history of gastrointestinal disease, particularly colitis.

Cephalosporins may be associated with a fall in prothrombin activity. Those at risk include patients with renal or hepatic impairment or poor nutritional state, as well as patients receiving a protracted course of antimicrobial therapy, and patients previously stabilized on anticoagulant therapy. Prothrombin time should be monitored in patients at risk and exogenous vitamin K administered as indicated.

Prescribing Cefazolin for Injection in the absence of a proven or strongly suspected bacterial infection or a prophylactic indication is unlikely to provide benefit to the patient and increases the risk of the development of drug-resistant bacteria.

Drug Interactions

Probenecid may decrease renal tubular secretion of cephalosporins when used concurrently, resulting in increased and more prolonged cephalosporin blood levels.

Drug/Laboratory Test Interactions

A false positive reaction for glucose in the urine may occur with Benedict's solution, Fehling's solution or with CLINITEST[®] tablets, but not with enzyme-based tests such as CLINISTIX[®].

Positive direct and indirect antiglobulin (Coombs) tests have occurred; these may also occur in neonates whose mothers received cephalosporins before delivery.

Information for Patients

Patients should be counseled that antibacterial drugs including Cefazolin for Injection, should only be used to treat bacterial infections. They do not treat viral infections (e.g., the common cold). When Cefazolin for Injection is prescribed to treat a bacterial infection, patients should be told that although it is common to feel better early in the course of therapy, the medication should be taken exactly as directed. Skipping doses or not completing the full course of therapy may: (1) decrease the effectiveness of the immediate treatment, and (2) increase the likelihood that bacteria will develop resistance and will not be treatable by Cefazolin for Injection or other antibacterial drugs in the future.

Carcinogenesis/Mutagenesis

Mutagenicity studies and long-term studies in animals to determine the carcinogenic potential of Cefazolin for Injection have not been performed.

Pregnancy

Teratogenic Effects

Pregnancy Category B

Reproduction studies have been performed in rats, mice and rabbits at doses up to 25 times the human dose and have revealed no evidence of impaired fertility or harm to the fetus due to Cefazolin for Injection. There are, however, no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

Labor and Delivery

When cefazolin has been administered prior to caesarean section, drug levels in cord blood have been approximately one quarter to one third of maternal drug levels. The drug appears to have no adverse effect on the fetus.

Nursing Mothers

Cefazolin for Injection is present in very low concentrations in the milk of nursing mothers. Caution should be exercised when Cefazolin for Injection is administered to a nursing woman.

Pediatric Use

Safety and effectiveness for use in premature infants and neonates have not been established. See **DOSAGE AND ADMINISTRATION** for recommended dosage in pediatric patients older than 1 month.

Geriatric Use

Of the 920 subjects who received Cefazolin for Injection in clinical studies, 313 (34%)

were 65 years and over, while 138 (15%) were 75 years and over. No overall differences in safety or effectiveness were observed between these subjects and younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients, but greater sensitivity of some older individuals cannot be ruled out.

This drug is known to be substantially excreted by the kidney, and the risk of toxic reactions to this drug may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function (see **PRECAUTIONS, General** and **DOSAGE AND ADMINISTRATION**).

ADVERSE REACTIONS

The following reactions have been reported:

Gastrointestinal

Diarrhea, oral candidiasis (oral thrush), vomiting, nausea, stomach cramps, anorexia, and pseudomembranous colitis. Onset of pseudomembranous colitis symptoms may occur during or after antibiotic treatment (see **WARNINGS**). Nausea and vomiting have been reported rarely.

Allergic

Anaphylaxis, eosinophilia, itching, drug fever, skin rash, Stevens-Johnson syndrome.

Hematologic

Neutropenia, leukopenia, thrombocytopenia, thrombocythemia.

Hepatic

Transient rise in SGOT, SGPT, and alkaline phosphatase levels has been observed. As with other cephalosporins, reports of hepatitis have been received.

Renal

As with other cephalosporins, reports of increased BUN and creatinine levels, as well as renal failure, have been received.

Local Reactions

Rare instances of phlebitis have been reported at site of injection. Pain at the site of injection after intramuscular administration has occurred infrequently. Some induration has occurred.

Other Reactions

Genital and anal pruritus (including vulvar pruritus, genital moniliasis, and vaginitis).

To report SUSPECTED ADVERSE REACTIONS, contact Hikma Pharmaceuticals USA Inc. at 1-877-845-0689, or the FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DOSAGE AND ADMINISTRATION

Usual Adult Dosage

Type of Infection	Dose	Frequency
--------------------------	-------------	------------------

Moderate to severe infections	500 mg to 1 gram	every 6 to 8 hours
Mild infections caused by susceptible gram-positive cocci	250 mg to 500 mg	every 8 hours
Acute, uncomplicated urinary tract infections	1 gram	every 12 hours
Pneumococcal pneumonia	500 mg	every 12 hours
Severe, life-threatening infections (e.g., endocarditis, septicemia)*	1 gram to 1.5 grams	every 6 hours

* In rare instances, doses of up to 12 grams of Cefazolin for Injection per day have been used.

Perioperative Prophylactic Use

To prevent postoperative infection in contaminated or potentially contaminated surgery, recommended doses are:

- a. 1 gram IV or IM administered 1/2 hour to 1 hour prior to the start of surgery.
- b. For lengthy operative procedures (e.g., 2 hours or more), 500 mg to 1 gram IV or IM during surgery (administration modified depending on the duration of the operative procedure).
- c. 500 mg to 1 gram IV or IM every 6 to 8 hours for 24 hours postoperatively.

It is important that (1) the preoperative dose be given just (1/2 to 1 hour) prior to the start of surgery so that adequate antibiotic levels are present in the serum and tissues at the time of initial surgical incision; and (2) Cefazolin for Injection be administered, if necessary, at appropriate intervals during surgery to provide sufficient levels of the antibiotic at the anticipated moments of greatest exposure to infective organisms.

In surgery where the occurrence of infection may be particularly devastating (e.g., open-heart surgery and prosthetic arthroplasty), the prophylactic administration of Cefazolin for Injection may be continued for 3 to 5 days following the completion of surgery.

Dosage Adjustment for Patients with Reduced Renal Function

Cefazolin for Injection may be used in patients with reduced renal function with the following dosage adjustments: Patients with a creatinine clearance of 55 mL/min. or greater or a serum creatinine of 1.5 mg % or less can be given full doses. Patients with creatinine clearance rates of 35 to 54 mL/min. or serum creatinine of 1.6 to 3 mg % can also be given full doses but dosage should be restricted to at least 8 hour intervals. Patients with creatinine clearance rates of 11 to 34 mL/min. or serum creatinine of 3.1 to 4.5 mg % should be given 1/2 the usual dose every 12 hours. Patients with creatinine clearance rates of 10 mL/min. or less or serum creatinine of 4.6 mg % or greater should be given 1/2 the usual dose every 18 to 24 hours. All reduced dosage recommendations apply after an initial loading dose appropriate to the severity of the infection. Patients undergoing peritoneal dialysis: See **CLINICAL PHARMACOLOGY**.

Pediatric Dosage

In pediatric patients, a total daily dosage of 25 to 50 mg per kg (approximately 10 to 20 mg per pound) of body weight, divided into 3 or 4 equal doses, is effective for most mild to moderately severe infections. Total daily dosage may be increased to 100 mg per kg

(45 mg per pound) of body weight for severe infections. Since safety for use in premature infants and in neonates has not been established, the use of Cefazolin for Injection in these patients is not recommended.

Pediatric Dosage Guide

Weight		25 mg/kg/day Divided into 3 Doses		25 mg/kg/day Divided into 4 Doses	
Lbs	Kg	Approximate Single Dose mg/q8h	Vol. (mL) needed with dilution of 125 mg/mL	Approximate Single Dose mg/q6h	Vol. (mL) needed with dilution of 125 mg/mL
10	4.5	40 mg	0.35 mL	30 mg	0.25 mL
20	9	75 mg	0.6 mL	55 mg	0.45 mL
30	13.6	115 mg	0.9 mL	85 mg	0.7 mL
40	18.1	150 mg	1.2 mL	115 mg	0.9 mL
50	22.7	190 mg	1.5 mL	140 mg	1.1 mL

Weight		50 mg/kg/day Divided into 3 Doses		50 mg/kg/day Divided into 4 Doses	
Lbs	Kg	Approximate Single Dose mg/q8h	Vol. (mL) needed with dilution of 225 mg/mL	Approximate Single Dose mg/q6h	Vol. (mL) needed with dilution of 225 mg/mL
10	4.5	75 mg	0.35 mL	55 mg	0.25 mL
20	9	150 mg	0.7 mL	110 mg	0.5 mL
30	13.6	225 mg	1 mL	170 mg	0.75 mL
40	18.1	300 mg	1.35 mL	225 mg	1 mL
50	22.7	375 mg	1.7 mL	285 mg	1.25 mL

In pediatric patients with mild to moderate renal impairment (creatinine clearance of 70 to 40 mL/min.), 60 percent of the normal daily dose given in equally divided doses every 12 hours should be sufficient. In patients with moderate impairment (creatinine clearance of 40 to 20 mL/min.), 25 percent of the normal daily dose given in equally divided doses every 12 hours should be adequate. Pediatric patients with severe renal impairment (creatinine clearance of 20 to 5 mL/min.) may be given 10 percent of the normal daily dose every 24 hours. All dosage recommendations apply after an initial loading dose.

RECONSTITUTION

Preparation of Parenteral Solution

Parenteral drug products should be SHAKEN WELL when reconstituted, and inspected visually for particulate matter prior to administration. If particulate matter is evident in reconstituted fluids, the drug solutions should be discarded.

When reconstituted or diluted according to the instructions below, Cefazolin for Injection is stable for 24 hours at room temperature or for 10 days if stored under refrigeration (5°C or 41°F). Reconstituted solutions may range in color from pale yellow to yellow without a change in potency.

Single-Dose Vials

For IM injection, IV direct (bolus) injection or IV infusion, reconstitute with Sterile Water for Injection according to the following table. SHAKE WELL.

Vial Size	Amount of Diluent	Approximate Concentration	Approximate Available Volume
500 mg	2 mL	225 mg/mL	2.2 mL
1 gram	2.5 mL	330 mg/mL	3 mL

ADMINISTRATION

Intramuscular Administration

Reconstitute vials with Sterile Water for Injection according to the dilution table above. Shake well until dissolved. Cefazolin for Injection should be injected into a large muscle mass. Pain on injection is infrequent with Cefazolin for Injection.

Intravenous Administration

Direct (bolus) injection: Following reconstitution according to the above table, further dilute vials with approximately 5 mL Sterile Water for Injection. Inject the solution slowly over 3 to 5 minutes, directly or through tubing for patients receiving parenteral fluids (see list below).

Intermittent or continuous infusion: Dilute reconstituted Cefazolin for Injection in 50 to 100 mL of 1 of the following solutions:

- Sodium Chloride Injection, USP
- 5% or 10% Dextrose Injection, USP
- 5% Dextrose in Lactated Ringer's Injection, USP
- 5% Dextrose and 0.9% Sodium Chloride Injection, USP
- 5% Dextrose and 0.45% Sodium Chloride Injection, USP
- 5% Dextrose and 0.2% Sodium Chloride Injection, USP
- Lactated Ringer's Injection, USP
- Invert Sugar 5% or 10% in Sterile Water for Injection
- Ringer's Injection, USP
- 5% Sodium Bicarbonate Injection, USP

HOW SUPPLIED

Cefazolin for Injection, USP is supplied in vials containing cefazolin sodium equivalent to 500 mg or 1 gram cefazolin.

Cefazolin for Injection, USP	Vial Size	Packaged	NDC No.
500 mg	10 mL	Carton of 25 vials	0143-9923-90
1 gram	10 mL	Carton of 25 vials	0143-9924-90

Also available as Pharmacy Bulk Package, as follows:

Cefazolin for Injection, USP	Vial Size	Packaged	NDC No.
10 grams	100 mL	Carton of 10 vials	0143-9983-03
20 grams	100 mL	Carton of 10 vials	0143-9665-10

As with other cephalosporins, Cefazolin for Injection, USP tends to darken depending on storage conditions; within the stated recommendations, however, product potency is not adversely affected.

Before reconstitution protect from light and store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].

CLINITEST is a registered trademark of Miles, Inc.

CLINISTIX is a registered trademark of Bayer Corporation.

Manufactured by:

HIKMA FARMACÊUTICA (PORTUGAL) S.A.
Estrada do Rio da Mó, 8, 8A e 8B – Fervença
2705-906 Terrugem SNT PORTUGAL

Distributed by:

Hikma Pharmaceuticals USA Inc.
Berkeley Heights, NJ 07922

Revised: June 2020

PIN001-WES/8

PRINCIPAL DISPLAY PANEL

NDC 0143-**9923**-90 Rx only

Cefazolin

for Injection, USP

500 mg* per vial

For Intravenous or

Intramuscular use

NDC 0143-9923-90

Rx only

Cefazolin for Injection, USP

500 mg* per vial

For Intravenous or
Intramuscular use

*Each vial contains sterile cefazolin sodium equivalent to 500 mg cefazolin. The sodium content is 24 mg.

Usual Adult Dosage: 250 mg to 1 gram every 6 to 8 hours. For more information see package insert.

For Intramuscular Administration: Add 2.0 mL of Sterile Water for Injection, SHAKE WELL to dissolve. Withdraw entire contents. Provides an approximate volume of 2.2 mL (225 mg/mL).

For Intravenous Administration: See package insert. Reconstituted solution is stable for 24 hours at room temperature or for 10 days under refrigeration (5°C or 41°F). Reconstituted solutions may range in color from pale yellow to yellow without a change in potency.

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature], Protect from light.

Mfd. by HIKMA FARMACÊUTICA (PORTUGAL), S.A.
Dist. by Hikma, Berkeley Heights, NJ 07922



PLB136-WES/5

Lot:

Exp:

NDC 0143-9923-90 Rx only

Cefazolin

for Injection, USP

500 mg* per vial

For Intravenous or
Intramuscular use

25 x 500 mg Vials

Cefazolin for Injection, USP

500 mg* per vial

To open - Cut seal along dotted line.



NDC 0143-9923-90

PLB136-WES/2
Rx only

Cefazolin for Injection, USP

500 mg* per vial

For Intravenous or
Intramuscular use

25 x 500 mg Vials

*Each vial contains sterile cefazolin sodium equivalent to 500 mg cefazolin. The sodium content is 24 mg.

Usual Adult Dosage: 250 mg to 1 gram every 6 to 8 hours. For more information see package insert.

For Intramuscular Administration: Add 2.0 mL of Sterile Water for Injection, SHAKE WELL to dissolve. Withdraw entire contents. Provides an approximate volume of 2.2 mL (225 mg/mL).

For Intravenous Administration: See package insert. Reconstituted solution is stable for 24 hours at room temperature or for 10 days under refrigeration (5°C or 41°F). Reconstituted solutions may range in color from pale yellow to yellow without a change in potency.

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature], Protect from light.

Mfd. by HIKMA FARMACÊUTICA (PORTUGAL), S.A.
Dist. by Hikma, Berkeley Heights, NJ 07922

hikma.

PRINCIPAL DISPLAY PANEL

NDC 0143-9924-90 Rx only

Cefazolin

for Injection, USP
1 gram* per vial
For Intravenous or
Intramuscular use


NDC 0143-9924-90 Rx only

Cefazolin
for Injection, USP

1 gram* per vial

For Intravenous or
Intramuscular use

*Each vial contains sterile cefazolin sodium equivalent to 1 gram cefazolin. The sodium content is 48 mg.
Usual Adult Dosage: 250 mg to 1 gram every 6 to 8 hours. For more information see package insert.
For Intramuscular Administration: Add 2.5 mL of Sterile Water for Injection, SHAKE WELL to dissolve. Withdraw entire contents. Provides an approximate volume of 3.0 mL (330 mg/mL).
For Intravenous Administration: See package insert. Reconstituted solution is stable for 24 hours at room temperature or for 10 days under refrigeration (5°C or 41°F). Reconstituted solutions may range in color from pale yellow to yellow without a change in potency.
Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature]. Protect from light.
Mfd. by HIKMA FARMACÉUTICA (PORTUGAL), S.A.
Dist. by Hikma, Berkeley Heights, NJ 07922



(01)00301439924906

PLB001-WES/6

Lot: **Exp:**

NDC 0143-9924-90 Rx only

Cefazolin
for Injection, USP
1 gram* per vial
For Intravenous or
Intramuscular use

25 x 1 g Vials

Cefazolin
for Injection, USP
1 gram* per vial

To open - Cut seal along dotted line.



NDC 0143-9924-90

PLB132-WES/2
Rx only

Cefazolin
for Injection, USP

1 gram* per vial

For Intravenous or
Intramuscular use

25 x 1 g Vials

*Each vial contains sterile cefazolin sodium equivalent to 1 gram cefazolin. The sodium content is 48 mg.

Usual Adult Dosage: 250 mg to 1 gram every 6 to 8 hours. For more information see package insert.

For Intramuscular Administration: Add 2.5 mL of Sterile Water for Injection. SHAKE WELL to dissolve. Withdraw entire contents. Provides an approximate volume of 3.0 mL (330 mg/mL).

For Intravenous Administration: See package insert.

Reconstituted solution is stable for 24 hours at room temperature or for 10 days under refrigeration (5°C or 41°F). Reconstituted solutions may range in color from pale yellow to yellow without a change in potency.

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature]. Protect from light.

Mfd. by HIKMA FARMACÊUTICA (PORTUGAL), S.A.
Dist. by Hikma, Berkeley Heights, NJ 07822

hikma.

CEFAZOLIN

cefazolin injection, powder, for solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0143-9923
Route of Administration	INTRAMUSCULAR, INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CEFAZOLIN SODIUM (UNII: P380M0454Z) (CEFAZOLIN - UNII:IHS69LOY4T)	CEFAZOLIN	225 mg in 1 mL

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0143-9923-90	2.2 mL in 1 VIAL; Type 0: Not a Combination Product	09/18/2001	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA065047	09/18/2001	

CEFAZOLIN

cefazolin injection, powder, for solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0143-9924
Route of Administration	INTRAMUSCULAR, INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CEFAZOLIN SODIUM (UNII: P380M0454Z) (CEFAZOLIN - UNII:IHS69LOY4T)	CEFAZOLIN	330 mg in 1 mL

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0143-9924-90	3 mL in 1 VIAL; Type 0: Not a Combination Product	09/18/2001	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA065047	09/18/2001	

Labeler - Hikma Pharmaceuticals USA Inc. (001230762)

Establishment

Name	Address	ID/FEI	Business Operations
HIKMA FARMACEUTICA (PORTUGAL), S.A		452742943	label(0143-9923, 0143-9924) , manufacture(0143-9923, 0143-9924) , pack(0143-9923, 0143-9924)

Revised: 4/2024

Hikma Pharmaceuticals USA Inc.