

PEMETREXED- pemetrexed injection, powder, lyophilized, for solution

Bamboo US Bidco LCC

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use PEMETREXED FOR INJECTION safely and effectively. See full prescribing information for PEMETREXED FOR INJECTION.

PEMETREXED for injection, for intravenous use

Initial U.S. Approval: 2004

INDICATIONS AND USAGE

Pemetrexed for Injection is a folate analog metabolic inhibitor indicated:

- in combination with pembrolizumab and platinum chemotherapy, for the initial treatment of patients with metastatic non-squamous non-small cell lung cancer (NSCLC), with no EGFR or ALK genomic tumor aberrations. (1.1)
- in combination with cisplatin for the initial treatment of patients with locally advanced or metastatic, non-squamous NSCLC. (1.1)
- as a single agent for the maintenance treatment of patients with locally advanced or metastatic, non-squamous NSCLC whose disease has not progressed after four cycles of platinum-based first-line chemotherapy. (1.1)
- as a single agent for the treatment of patients with recurrent, metastatic non-squamous, NSCLC after prior chemotherapy. (1.1) Limitations of Use: Pemetrexed for Injection is not indicated for the treatment of patients with squamous cell, NSCLC. (1.1)
- initial treatment, in combination with cisplatin, of patients with malignant pleural mesothelioma whose disease is unresectable or who are otherwise not candidates for curative surgery. (1.2)

DOSAGE AND ADMINISTRATION

- The recommended dose of Pemetrexed for Injection administered with pembrolizumab and platinum chemotherapy in patients with a creatinine clearance (calculated by Cockcroft-Gault equation) of 45 mL/min or greater is 500 mg/m² as an intravenous infusion over 10 minutes, administered after pembrolizumab and prior to platinum chemotherapy, on Day 1 of each 21-day cycle. (2.1)
- The recommended dose of Pemetrexed for Injection, administered as a single agent or with cisplatin, in patients with creatinine clearance of 45 mL/minute or greater is 500 mg/m² as an intravenous infusion over 10 minutes on Day 1 of each 21-day cycle. (2.1, 2.2)
- Initiate folic acid 400 mcg to 1000 mcg orally, once daily, beginning 7 days prior to the first dose of Pemetrexed for Injection and continue until 21 days after the last dose of Pemetrexed for Injection. (2.4)
- Administer vitamin B₁₂, 1 mg intramuscularly, 1 week prior to the first dose of Pemetrexed for Injection and every 3 cycles. (2.4)
- Administer dexamethasone 4 mg orally, twice daily the day before, the day of, and the day after Pemetrexed for Injection administration. (2.4)

DOSAGE FORMS AND STRENGTHS

For Injection: 100 mg or 500 mg lyophilized powder in single-dose vial (3)

CONTRAINDICATIONS

History of severe hypersensitivity reaction to pemetrexed. (4)

WARNINGS AND PRECAUTIONS

- Myelosuppression: Can cause severe bone marrow suppression resulting in cytopenia and an increased risk of infection. Do not administer Pemetrexed for Injection when the absolute neutrophil count is less than 1500 cells/mm³ and platelets are less than 100,000 cells/mm³. Initiate supplementation with oral folic acid and intramuscular vitamin B₁₂ to reduce the severity of hematologic and gastrointestinal toxicity of Pemetrexed for Injection. (2.4, 5.1)
- Renal Failure: Can cause severe, and sometimes fatal, renal failure. Do not administer when creatinine clearance is less than 45 mL/min. (2.3, 5.2)
- Bullous and Exfoliative Skin Toxicity: Permanently discontinue for severe and life-threatening bullous, blistering or exfoliating skin toxicity. (5.3)
- Interstitial Pneumonitis: Withhold for acute onset of new or progressive unexplained pulmonary symptoms. Permanently discontinue if pneumonitis is confirmed. (5.4)

- Radiation Recall: Can occur in patients who received radiation weeks to years previously; permanently discontinue for signs of radiation recall. (5.5)
- Embryo-Fetal Toxicity: Can cause fetal harm. Advise patients of the potential risk to a fetus and to use effective contraception. (5.7, 8.1, 8.3)

-----ADVERSE REACTIONS-----

- The most common adverse reactions (incidence $\geq 20\%$) of Pemetrexed for Injection, when administered as a single agent are fatigue, nausea, and anorexia. (6.1)
- The most common adverse reactions (incidence $\geq 20\%$) of Pemetrexed for Injection when administered with cisplatin are vomiting, neutropenia, anemia, stomatitis/pharyngitis, thrombocytopenia, and constipation. (6.1)
- The most common adverse reactions (incidence $\geq 20\%$) of Pemetrexed for Injection when administered in combination with pembrolizumab and platinum chemotherapy are fatigue/asthenia, nausea, constipation, diarrhea, decreased appetite, rash, vomiting, cough, dyspnea, and pyrexia. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Baxter Healthcare Corporation at 1-866-888-2472 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

-----DRUG INTERACTIONS-----

Ibuprofen increased risk of Pemetrexed for Injection toxicity in patients with mild to moderate renal impairment. Modify the ibuprofen dosage as recommended for patients with a creatinine clearance between 45 mL/min and 79 mL/min. (2.5, 5.6, 7)

-----USE IN SPECIFIC POPULATIONS-----

Lactation: Advise not to breastfeed. (8.2)

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 9/2023

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

- 1.1 Non-Squamous Non-Small Cell Lung Cancer (NSCLC)
- 1.2 Mesothelioma

2 DOSAGE AND ADMINISTRATION

- 2.1 Recommended Dosage for Non-Squamous NSCLC
- 2.2 Recommended Dosage for Mesothelioma
- 2.3 Renal Impairment
- 2.4 Premedication and Concomitant Medications to Mitigate Toxicity
- 2.5 Dosage Modification of Ibuprofen in Patients with Mild to Moderate Renal Impairment Receiving Pemetrexed for Injection
- 2.6 Dosage Modifications for Adverse Reactions
- 2.7 Preparation for Administration

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Myelosuppression and Increased Risk of Myelosuppression without Vitamin Supplementation
- 5.2 Renal Failure
- 5.3 Bullous and Exfoliative Skin Toxicity
- 5.4 Interstitial Pneumonitis
- 5.5 Radiation Recall
- 5.6 Increased Risk of Toxicity with Ibuprofen in Patients with Renal Impairment
- 5.7 Embryo-Fetal Toxicity

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience

7 DRUG INTERACTIONS

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation
- 8.3 Females and Males of Reproductive Potential
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Patients with Renal Impairment

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

- 14.1 Non-Squamous NSCLC
- 14.2 Mesothelioma

15 REFERENCES

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

1.1 Non-Squamous Non-Small Cell Lung Cancer (NSCLC)

Pemetrexed for Injection is indicated:

- in combination with pembrolizumab and platinum chemotherapy, for the initial treatment of patients with metastatic non-squamous NSCLC, with no EGFR or ALK genomic tumor aberrations.
- in combination with cisplatin for the initial treatment of patients with locally advanced or metastatic, non-squamous, NSCLC.
- as a single agent for the maintenance treatment of patients with locally advanced or metastatic, non-squamous NSCLC whose disease has not progressed after four cycles of platinum-based first-line chemotherapy.
- as a single agent for the treatment of patients with recurrent, metastatic non-squamous NSCLC after prior chemotherapy.
- Limitations of Use: Pemetrexed for Injection is not indicated for the treatment of patients with squamous cell, NSCLC [see *Clinical Studies (14.1)*].

1.2 Mesothelioma

Pemetrexed for Injection is indicated, in combination with cisplatin, for the initial treatment of patients with malignant pleural mesothelioma whose disease is unresectable or who are otherwise not candidates for curative surgery.

2 DOSAGE AND ADMINISTRATION

2.1 Recommended Dosage for Non-Squamous NSCLC

- The recommended dose of Pemetrexed for Injection when administered with pembrolizumab and platinum chemotherapy for the initial treatment of metastatic non-squamous NSCLC in patients with a creatinine clearance (calculated by Cockcroft-Gault equation) of 45 mL/min or greater is 500 mg/m² as an intravenous infusion over 10 minutes administered after pembrolizumab and prior to carboplatin or cisplatin on Day 1 of each 21-day cycle for 4 cycles. Following completion of platinum-based therapy, treatment with Pemetrexed for Injection with or without pembrolizumab is administered until disease progression or unacceptable toxicity. Please refer to the full prescribing information for pembrolizumab and for carboplatin or cisplatin.
- The recommended dose of Pemetrexed for Injection when administered with cisplatin for initial treatment of locally advanced or metastatic non-squamous NSCLC in patients with a creatinine clearance (calculated by Cockcroft-Gault equation) of 45 mL/min or greater is 500 mg/m² as an intravenous infusion over 10 minutes administered prior to cisplatin on Day 1 of each 21-day cycle for up to six cycles in the absence of disease progression or unacceptable toxicity.
- The recommended dose of Pemetrexed for Injection for maintenance treatment of non-squamous NSCLC in patients with a creatinine clearance (calculated by Cockcroft-Gault equation) of 45 mL/min or greater is 500 mg/m² as an intravenous infusion over 10 minutes on Day 1 of each 21-day cycle until disease progression or unacceptable toxicity after four cycles of platinum-based first-line chemotherapy.
- The recommended dose of Pemetrexed for Injection for treatment of recurrent non-squamous NSCLC in patients with a creatinine clearance (calculated by Cockcroft-Gault equation) of 45 mL/min or greater is 500 mg/m² as an intravenous infusion over 10 minutes on Day 1 of each 21-day cycle until disease progression or unacceptable toxicity.

2.2 Recommended Dosage for Mesothelioma

- The recommended dose of Pemetrexed for Injection when administered with cisplatin in patients with a creatinine clearance (calculated by Cockcroft-Gault equation) of 45 mL/min or greater is 500 mg/m² as an intravenous infusion over 10 minutes on Day 1 of each 21-day cycle until disease progression or unacceptable toxicity.

2.3 Renal Impairment

- Pemetrexed for Injection dosing recommendations are provided for patients with a creatinine clearance (calculated by Cockcroft-Gault equation) of 45 mL/min or greater [see *Dosage and Administration (2.1, 2.2)*]. There is no recommended dose for patients whose creatinine clearance is less than 45 mL/min [see *Use in Specific Populations (8.6)*].

2.4 Premedication and Concomitant Medications to Mitigate Toxicity

Vitamin Supplementation

- Initiate folic acid 400 mcg to 1000 mcg orally once daily, beginning 7 days before the first dose of Pemetrexed for Injection and continuing until 21 days after the last dose of Pemetrexed for Injection [see *Warnings and Precautions (5.1)*].
- Administer vitamin B₁₂, 1 mg intramuscularly, 1 week prior to the first dose of Pemetrexed for Injection and every 3 cycles thereafter. Subsequent vitamin B₁₂ injections may be given the same day as treatment with Pemetrexed for Injection [see *Warnings and Precautions (5.1)*]. **Do not substitute oral vitamin B₁₂ for intramuscular vitamin B₁₂.**

Corticosteroids

- Administer dexamethasone 4 mg orally twice daily for three consecutive days, beginning the day before each Pemetrexed for Injection administration.

2.5 Dosage Modification of Ibuprofen in Patients with Mild to Moderate Renal Impairment Receiving Pemetrexed for Injection

In patients with creatinine clearances between 45 mL/min and 79 mL/min, modify administration of ibuprofen as follows [see *Warnings and Precautions (5.6)*, *Drug Interactions (7)* and *Clinical Pharmacology (12.3)*]:

- Avoid administration of ibuprofen for 2 days before, the day of, and 2 days following administration of Pemetrexed for Injection.
- Monitor patients more frequently for myelosuppression, renal, and gastrointestinal toxicity, if concomitant administration of ibuprofen cannot be avoided.

2.6 Dosage Modifications for Adverse Reactions

Obtain complete blood count on Days 1, 8, and 15 of each cycle. Assess creatinine clearance prior to each cycle. Do not administer Pemetrexed for Injection if the creatinine clearance is less than 45 mL/min.

Delay initiation of the next cycle of Pemetrexed for Injection until:

- recovery of non-hematologic toxicity to Grade 0-2,
- absolute neutrophil count (ANC) is 1500 cells/mm³ or higher, and
- platelet count is 100,000 cells/mm³ or higher.

Upon recovery, modify the dosage of Pemetrexed for Injection in the next cycle as specified in Table 1.

For dosing modifications for cisplatin, carboplatin, or pembrolizumab, refer to their prescribing information.

Table 1: Recommended Dosage Modifications for Adverse Reactions *

Toxicity in Most Recent Treatment Cycle	Pemetrexed for Injection Dose Modification for Next Cycle
Myelosuppressive toxicity [see <i>Warnings and Precautions (5.1)</i>]	
1. ANC less than 500/mm ³ and platelets greater than or equal to 50,000/mm ³	

2. <u>OR</u> 3. Platelet count less than 50,000/mm ³ without bleeding.	75% of previous dose
1. Platelet count less than 50,000/mm ³ with bleeding	50% of previous dose
1. Recurrent Grade 3 or 4 myelosuppression after 2 dose reductions	Discontinue
Non-hematologic toxicity	
1. Any Grade 3 or 4 toxicities EXCEPT mucositis or neurologic toxicity 2. <u>OR</u> 3. Diarrhea requiring hospitalization	75% of previous dose
1. Grade 3 or 4 mucositis	50% of previous dose
1. Renal toxicity [see Warnings and Precautions (5.2)]	Withhold until creatinine clearance is 45 mL/min or greater
1. Grade 3 or 4 neurologic toxicity	Permanently discontinue
1. Recurrent Grade 3 or 4 non-hematologic toxicity after 2 dose reductions	Permanently discontinue
1. Severe and life-threatening Skin Toxicity [see Warnings and Precautions (5.3)]	Permanently discontinue
1. Interstitial Pneumonitis [see Warnings and Precautions (5.4)]	Permanently discontinue

* National Cancer Institute Common Toxicity Criteria for Adverse Events version 2 (NCI CTCAE v2).

2.7 Preparation for Administration

- Pemetrexed for Injection is a hazardous drug. Follow applicable special handling and disposal procedures. ¹
- Calculate the dose of Pemetrexed for Injection and determine the number of vials needed.
- Reconstitute Pemetrexed for Injection to achieve a concentration of 25 mg/mL as follows:
 - Reconstitute each 100-mg vial with 4.2 mL of 0.9% Sodium Chloride Injection, USP (preservative-free)
 - Reconstitute each 500-mg vial with 20 mL of 0.9% Sodium Chloride Injection, USP (preservative-free)
 - Do not use calcium-containing solutions for reconstitution.
- Gently swirl each vial until the powder is completely dissolved. The resulting solution is clear and ranges in color from colorless to yellow or green-yellow. FURTHER DILUTION IS REQUIRED prior to administration.

- Store reconstituted, preservative-free product under refrigerated conditions [2-8°C (36-46°F)] for no longer than 24 hours from the time of reconstitution. Discard vial after 24 hours.
- Inspect reconstituted product visually for particulate matter and discoloration prior to further dilution. If particulate matter is observed, discard vial.
- Withdraw the calculated dose of Pemetrexed for Injection from the vial(s) and discard vial with any unused portion.
- Further dilute Pemetrexed for Injection with 0.9% Sodium Chloride Injection (preservative-free) to achieve a total volume of 100 mL for intravenous infusion.
- Store diluted, reconstituted product under refrigerated conditions [2-8°C (36-46°F)] for no more than 24 hours from the time of reconstitution. Discard after 24 hours.

3 DOSAGE FORMS AND STRENGTHS

For injection: 100 mg or 500 mg pemetrexed as a white to light-yellow or green-yellow lyophilized powder in single-dose vials for reconstitution.

4 CONTRAINDICATIONS

Pemetrexed for Injection is contraindicated in patients with a history of severe hypersensitivity reaction to pemetrexed [*see Adverse Reactions (6.1)*].

5 WARNINGS AND PRECAUTIONS

5.1 Myelosuppression and Increased Risk of Myelosuppression without Vitamin Supplementation

Pemetrexed for Injection can cause severe myelosuppression resulting in a requirement for transfusions and which may lead to neutropenic infection. The risk of myelosuppression is increased in patients who do not receive vitamin supplementation. In Study JMCH, incidences of Grade 3-4 neutropenia (38% versus 23%), thrombocytopenia (9% versus 5%), febrile neutropenia (9% versus 0.6%), and neutropenic infection (6% versus 0) were higher in patients who received Pemetrexed for Injection plus cisplatin without vitamin supplementation as compared to patients who were fully supplemented with folic acid and vitamin B₁₂ prior to and throughout Pemetrexed for Injection plus cisplatin treatment.

Initiate supplementation with oral folic acid and intramuscular vitamin B₁₂ prior to the first dose of Pemetrexed for Injection; continue vitamin supplementation during treatment and for 21 days after the last dose of Pemetrexed for Injection to reduce the severity of hematologic and gastrointestinal toxicity of Pemetrexed for Injection [*see Dosage and Administration (2.4)*]. Obtain a complete blood count at the beginning of each cycle. Do not administer Pemetrexed for Injection until the ANC is at least 1500 cells/mm³ and platelet count is at least 100,000 cells/mm³. Permanently reduce Pemetrexed for Injection in patients with an ANC of less than 500 cells/mm³ or platelet count of less than 50,000 cells/mm³ in previous cycles [*see Dosage and Administration (2.6)*].

In Studies JMDB and JMCH, among patients who received vitamin supplementation, incidence of Grade 3-4 neutropenia was 15% and 23%, the incidence of Grade 3-4

anemia was 6% and 4%, and incidence of Grade 3-4 thrombocytopenia was 4% and 5%, respectively. In Study JMCH, 18% of patients in the Pemetrexed for Injection arm required red blood cell transfusions compared to 7% of patients in the cisplatin arm [see *Adverse Reactions (6.1)*]. In Studies JMEN, PARAMOUNT, and JMEI, where all patients received vitamin supplementation, incidence of Grade 3-4 neutropenia ranged from 3% to 5%, and incidence of Grade 3-4 anemia ranged from 3% to 5%.

5.2 Renal Failure

Pemetrexed for Injection can cause severe, and sometimes fatal, renal toxicity. The incidences of renal failure in clinical studies in which patients received Pemetrexed for Injection with cisplatin were: 2.1% in Study JMDB and 2.2% in Study JMCH. The incidence of renal failure in clinical studies in which patients received Pemetrexed for Injection as a single agent ranged from 0.4% to 0.6% (Studies JMEN, PARAMOUNT, and JMEI [see *Adverse Reactions (6.1)*]. Determine creatinine clearance before each dose and periodically monitor renal function during treatment with Pemetrexed for Injection. Withhold Pemetrexed for Injection in patients with a creatinine clearance of less than 45 mL/minute [see *Dosage and Administration (2.3)*].

5.3 Bullous and Exfoliative Skin Toxicity

Serious and sometimes fatal, bullous, blistering and exfoliative skin toxicity, including cases suggestive of Stevens-Johnson Syndrome/Toxic epidermal necrolysis can occur with Pemetrexed for Injection. Permanently discontinue Pemetrexed for Injection for severe and life-threatening bullous, blistering or exfoliating skin toxicity.

5.4 Interstitial Pneumonitis

Serious interstitial pneumonitis, including fatal cases, can occur with Pemetrexed for Injection treatment. Withhold Pemetrexed for Injection for acute onset of new or progressive unexplained pulmonary symptoms such as dyspnea, cough, or fever pending diagnostic evaluation. If pneumonitis is confirmed, permanently discontinue Pemetrexed for Injection.

5.5 Radiation Recall

Radiation recall can occur with Pemetrexed for Injection in patients who have received radiation weeks to years previously. Monitor patients for inflammation or blistering in areas of previous radiation treatment. Permanently discontinue Pemetrexed for Injection for signs of radiation recall.

5.6 Increased Risk of Toxicity with Ibuprofen in Patients with Renal Impairment

Exposure to Pemetrexed for Injection is increased in patients with mild to moderate renal impairment who take concomitant ibuprofen, increasing the risks of adverse reactions of Pemetrexed for Injection. In patients with creatinine clearances between 45 mL/min and 79 mL/min, avoid administration of ibuprofen for 2 days before, the day of, and 2 days following administration of Pemetrexed for Injection. If concomitant ibuprofen use cannot be avoided, monitor patients more frequently for Pemetrexed for Injection adverse reactions, including myelosuppression, renal, and gastrointestinal toxicity [see *Dosage and Administration (2.5)*, *Drug Interactions (7)*, and *Clinical*

Pharmacology (12.3)] .

5.7 Embryo-Fetal Toxicity

Based on findings from animal studies and its mechanism of action, Pemetrexed for Injection can cause fetal harm when administered to a pregnant woman. In animal reproduction studies, intravenous administration of pemetrexed to pregnant mice during the period of organogenesis was teratogenic, resulting in developmental delays and increased malformations at doses lower than the recommended human dose of 500 mg/m². Advise pregnant women of the potential risk to the fetus. Advise females of reproductive potential to use effective contraception during treatment with Pemetrexed for Injection and for 6 months after the last dose. Advise males with female partners of reproductive potential to use effective contraception during treatment with Pemetrexed for Injection and for 3 months after the last dose [*see Use in Specific Populations (8.1, 8.3) and Clinical Pharmacology (12.1)] .*

6 ADVERSE REACTIONS

The following adverse reactions are discussed in greater detail in other sections of the labeling:

- Myelosuppression [*see Warnings and Precautions (5.1)]*
- Renal failure [*see Warnings and Precautions (5.2)]*
- Bullous and exfoliative skin toxicity [*see Warning and Precautions (5.3)]*
- Interstitial pneumonitis [*see Warnings and Precautions (5.4)]*
- Radiation recall [*see Warnings and Precautions (5.5)]*

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reactions rates cannot be directly compared to rates in other clinical trials and may not reflect the rates observed in clinical practice.

In clinical trials, the most common adverse reactions (incidence $\geq 20\%$) of Pemetrexed for Injection, when administered as a single agent, are fatigue, nausea, and anorexia. The most common adverse reactions (incidence $\geq 20\%$) of Pemetrexed for Injection, when administered in combination with cisplatin are vomiting, neutropenia, anemia, stomatitis/pharyngitis, thrombocytopenia, and constipation. The most common adverse reactions (incidence $\geq 20\%$) of Pemetrexed for Injection, when administered in combination with pembrolizumab and platinum chemotherapy, are fatigue/asthenia, nausea, constipation, diarrhea, decreased appetite, rash, vomiting, cough, dyspnea, and pyrexia.

Non-Squamous NSCLC

First-line Treatment of Metastatic Non-squamous NSCLC with Pembrolizumab and Platinum Chemotherapy

The safety of Pemetrexed for Injection, in combination with pembrolizumab and investigator's choice of platinum (either carboplatin or cisplatin), was investigated in Study KEYNOTE-189, a multicenter, double-blind, randomized (2:1), active-controlled trial in patients with previously untreated, metastatic non-squamous NSCLC with no EGFR or ALK genomic tumor aberrations. A total of 607 patients received Pemetrexed for Injection, pembrolizumab, and platinum every 3 weeks for 4 cycles followed by

Pemetrexed for Injection and pembrolizumab (n=405), or placebo, Pemetrexed for Injection, and platinum every 3 weeks for 4 cycles followed by placebo and Pemetrexed for Injection (n=202). Patients with autoimmune disease that required systemic therapy within 2 years of treatment; a medical condition that required immunosuppression; or who had received more than 30 Gy of thoracic radiation within the prior 26 weeks were ineligible [see *Clinical Studies (14.1)*].

The median duration of exposure to Pemetrexed for Injection was 7.2 months (range: 1 day to 1.7 years). Seventy-two percent of patients received carboplatin. The study population characteristics were: median age of 64 years (range: 34 to 84), 49% age 65 years or older, 59% male, 94% White and 3% Asian, and 18% with history of brain metastases at baseline.

Pemetrexed for Injection was discontinued for adverse reactions in 23% of patients in the Pemetrexed for Injection, pembrolizumab, and platinum arm. The most common adverse reactions resulting in discontinuation of Pemetrexed for Injection in this arm were acute kidney injury (3%) and pneumonitis (2%). Adverse reactions leading to interruption of Pemetrexed for Injection occurred in 49% of patients in the Pemetrexed for Injection, pembrolizumab, and platinum arm. The most common adverse reactions or laboratory abnormalities leading to interruption of Pemetrexed for Injection in this arm ($\geq 2\%$) were neutropenia (12%), anemia (7%), asthenia (4%), pneumonia (4%), thrombocytopenia (4%), increased blood creatinine (3%), diarrhea (3%), and fatigue (3%).

Table 2 summarizes the adverse reactions that occurred in $\geq 20\%$ of patients treated with Pemetrexed for Injection, pembrolizumab, and platinum.

Table 2: Adverse Reactions Occurring in $\geq 20\%$ of Patients in KEYNOTE-189

Adverse Reaction	Pemetrexed for Injection Pembrolizumab Platinum Chemotherapy n=405		Placebo Pemetrexed for Injection Platinum Chemotherapy n=202	
	All Grades* (%)	Grade 3-4 (%)	All Grades (%)	Grade 3-4 (%)
Gastrointestinal Disorders				
1. Nausea	56	3.5	52	3.5
1. Constipation	35	1.0	32	0.5
1. Diarrhea	31	5	21	3.0
1. Vomiting	24	3.7	23	3.0
General Disorders and Administration Site Conditions				
1. Fatigue †	56	12	58	6
1. Pyrexia	20	0.2	15	0

Metabolism and Nutrition Disorders				
1. Decreased appetite	28	1.5	30	0.5
Skin and Subcutaneous Tissue Disorders				
1. Rash ‡	25	2.0	17	2.5
Respiratory, Thoracic and Mediastinal Disorders				
1. Cough	21	0	28	0
1. Dyspnea	21	3.7	26	5

* Graded per NCI CTCAE version 4.03.

† Includes asthenia and fatigue.

‡ Includes genital rash, rash, rash generalized, rash macular, rash maculo-papular, rash papular, rash pruritic, and rash pustular.

Table 3 summarizes the laboratory abnormalities that worsened from baseline in at least 20% of patients treated with Pemetrexed for Injection, pembrolizumab, and platinum.

Table 3: Laboratory Abnormalities Worsened from Baseline in $\geq 20\%$ of Patients in KEYNOTE-189

Laboratory Test*	Pemetrexed for Injection Pembrolizumab Platinum Chemotherapy		Placebo Pemetrexed for Injection Platinum Chemotherapy	
	All Grades[†] %	Grades 3- 4 %	All Grades %	Grades 3- 4 %
Chemistry				
1. Hyperglycemia	63	9	60	7
1. Increased ALT	47	3.8	42	2.6
1. Increased AST	47	2.8	40	1.0
1. Hypoalbuminemia	39	2.8	39	1.1
1. Increased creatinine	37	4.2	25	1.0
1. Hyponatremia	32	7	23	6
1. Hypophosphatemia	30	10	28	14
1. Increased alkaline phosphatase	26	1.8	29	2.1

1. Hypocalcemia	24	2.8	17	0.5
1. Hyperkalemia	24	2.8	19	3.1
1. Hypokalemia	21	5	20	5
Hematology				
1. Anemia	85	17	81	18
1. Lymphopenia	64	22	64	25
1. Neutropenia	48	20	41	19
1. Thrombocytopenia	30	12	29	8

* Each test incidence is based on the number of patients who had both baseline and at least one on-study laboratory measurement available: Pemetrexed for Injection/pembrolizumab/platinum chemotherapy (range: 381 to 401 patients) and placebo/Pemetrexed for Injection/platinum chemotherapy (range: 184 to 197 patients).

† Graded per NCI CTCAE version 4.03.

Initial Treatment in Combination with Cisplatin

The safety of Pemetrexed for Injection was evaluated in Study JMDB, a randomized (1:1), open-label, multicenter trial conducted in chemotherapy-naïve patients with locally advanced or metastatic NSCLC. Patients received either Pemetrexed for Injection 500 mg/m² intravenously and cisplatin 75 mg/m² intravenously on Day 1 of each 21-day cycle (n=839) or gemcitabine 1250 mg/m² intravenously on Days 1 and 8 and cisplatin 75 mg/m² intravenously on Day 1 of each 21-day cycle (n=830). All patients were fully supplemented with folic acid and vitamin B₁₂.

Study JMDB excluded patients with an Eastern Cooperative Oncology Group Performance Status (ECOG PS of 2 or greater), uncontrolled third-space fluid retention, inadequate bone marrow reserve and organ function, or a calculated creatinine clearance less than 45 mL/min. Patients unable to stop using aspirin or other non-steroidal anti-inflammatory drugs or unable to take folic acid, vitamin B₁₂ or corticosteroids were also excluded from the study.

The data described below reflect exposure to Pemetrexed for Injection plus cisplatin in 839 patients in Study JMDB. Median age was 61 years (range 26-83 years); 70% of patients were men; 78% were White, 16% were Asian, 2.9% were Hispanic or Latino, 2.1% were Black or African American, and <1% were other ethnicities; 36% had an ECOG PS 0. Patients received a median of 5 cycles of Pemetrexed for Injection.

Table 4 provides the frequency and severity of adverse reactions that occurred in ≥5% of 839 patients receiving Pemetrexed for Injection in combination with cisplatin in Study JMDB. Study JMDB was not designed to demonstrate a statistically significant reduction in adverse reaction rates for Pemetrexed for Injection, as compared to the control arm, for any specified adverse reaction listed in Table 4.

Table 4: Adverse Reactions Occurring in ≥5% of Fully Vitamin-Supplemented Patients Receiving Pemetrexed for Injection in Combination with Cisplatin

Chemotherapy in Study JMDB

Adverse Reaction*	Pemetrexed for Injection/ Cisplatin (N=839)		Gemcitabine/Cisplatin (N=830)	
	All Grades (%)	Grade 3-4 (%)	All Grades (%)	Grade 3-4 (%)
All adverse reactions	90	37	91	53
Laboratory				
Hematologic				
1. Anemia	33	6	46	10
1. Neutropenia	29	15	38	27
1. Thrombocytopenia	10	4	27	13
Renal				
1. Elevated creatinine	10	1	7	1
Clinical				
Constitutional symptoms				
1. Fatigue	43	7	45	5
Gastrointestinal				
1. Nausea	56	7	53	4
1. Vomiting	40	6	36	6
1. Anorexia	27	2	24	1
1. Constipation	21	1	20	0
1. Stomatitis/pharyngitis	14	1	12	0
1. Diarrhea	12	1	13	2
1. Dyspepsia/heartburn	5	0	6	0
Neurology				
1. Sensory neuropathy	9	0	12	1
1. Taste disturbance	8	0	9	0
Dermatology/Skin				
1. Alopecia	12	0	21	1

1. Rash/Desquamation	7	0	8	1
----------------------	---	---	---	---

* NCI CTCAE version 2.0.

The following additional adverse reactions of Pemetrexed for Injection were observed.

1. **Incidence 1% to <5%**

- Body as a Whole*— febrile neutropenia, infection, pyrexia
- General Disorders*— dehydration
- Metabolism and Nutrition*— increased AST, increased ALT
- Renal*— renal failure
- Eye Disorder*— conjunctivitis

2. **Incidence <1%**

- Cardiovascular*— arrhythmia
- General Disorders*— chest pain
- Metabolism and Nutrition*— increased GGT
- Neurology*— motor neuropathy

Maintenance Treatment Following First-line Non-Pemetrexed for Injection Containing Platinum-Based Chemotherapy

In Study JMEN, the safety of Pemetrexed for Injection was evaluated in a randomized (2:1), placebo-controlled, multicenter trial conducted in patients with non-progressive locally advanced or metastatic NSCLC following four cycles of a first-line, platinum-based chemotherapy regimen. Patients received either Pemetrexed for Injection 500 mg/m² or matching placebo intravenously every 21 days until disease progression or unacceptable toxicity. Patients in both study arms were fully supplemented with folic acid and vitamin B₁₂.

Study JMEN excluded patients with an ECOG PS of 2 or greater, uncontrolled third-space fluid retention, inadequate bone marrow reserve and organ function, or a calculated creatinine clearance less than 45 mL/min. Patients unable to stop using aspirin or other non-steroidal anti-inflammatory drugs or unable to take folic acid, vitamin B₁₂ or corticosteroids were also excluded from the study.

The data described below reflect exposure to Pemetrexed for Injection in 438 patients in Study JMEN. Median age was 61 years (range 26-83 years), 73% of patients were men; 65% were White, 31% were Asian, 2.9% were Hispanic or Latino, and <2% were other ethnicities; 39% had an ECOG PS 0. Patients received a median of 5 cycles of Pemetrexed for Injection and a relative dose intensity of Pemetrexed for Injection of 96%. Approximately half the patients (48%) completed at least six, 21-day cycles and 23% completed ten or more 21-day cycles of Pemetrexed for Injection.

Table 5 provides the frequency and severity of adverse reactions reported in ≥5% of the 438 Pemetrexed for Injection-treated patients in Study JMEN.

Table 5: Adverse Reactions Occurring in ≥5% of Patients Receiving Pemetrexed for Injection in Study JMEN

Adverse Reaction*	Pemetrexed for Injection (N=438)		Placebo (N=218)	
	All Grades (%)	Grade 3-4 (%)	All Grades	Grade 3-4 (%)

			(%)	
All adverse reactions	66	16	37	4
1. Laboratory				
1. Hematologic				
1. Anemia	15	3	6	1
1. Neutropenia	6	3	0	0
1. Hepatic				
1. Increased ALT	10	0	4	0
1. Increased AST	8	0	4	0
1. Clinical				
1. Constitutional symptoms				
1. Fatigue	25	5	11	1
1. Gastrointestinal				
1. Nausea	19	1	6	1
1. Anorexia	19	2	5	0
1. Vomiting	9	0	1	0
1. Mucositis/stomatitis	7	1	2	0
1. Diarrhea	5	1	3	0
1. Infection	5	2	2	0
1. Neurology				
1. Sensory neuropathy	9	1	4	0
1. Dermatology/Skin				
1. Rash/desquamation	10	0	3	0

* NCI CTCAE version 3.0.

The requirement for transfusions (9.5% versus 3.2%), primarily red blood cell

transfusions, and for erythropoiesis stimulating agents (5.9% versus 1.8%) were higher in the Pemetrexed for Injection arm compared to the placebo arm.

The following additional adverse reactions were observed in patients who received Pemetrexed for Injection.

1. Incidence 1% to <5%

Dermatology/Skin— alopecia, pruritus/itching

Gastrointestinal— constipation

General Disorders— edema, fever

Hematologic— thrombocytopenia

Eye Disorder— ocular surface disease (including conjunctivitis), increased lacrimation

2. Incidence <1%

Cardiovascular— supraventricular arrhythmia

Dermatology/Skin— erythema multiforme

General Disorders— febrile neutropenia, allergic reaction/hypersensitivity

Neurology— motor neuropathy

Renal— renal failure

Maintenance Treatment Following First-line Pemetrexed for Injection Plus Platinum Chemotherapy

The safety of Pemetrexed for Injection was evaluated in PARAMOUNT, a randomized (2:1), placebo-controlled study conducted in patients with non-squamous NSCLC with non-progressive (stable or responding disease) locally advanced or metastatic NSCLC following four cycles of Pemetrexed for Injection in combination with cisplatin as first-line therapy for NSCLC. Patients were randomized to receive Pemetrexed for Injection 500 mg/m² or matching placebo intravenously on Day 1 of each 21-day cycle until disease progression or unacceptable toxicity. Patients in both study arms received folic acid and vitamin B₁₂ supplementation.

PARAMOUNT excluded patients with an ECOG PS of 2 or greater, uncontrolled third-space fluid retention, inadequate bone marrow reserve and organ function, or a calculated creatinine clearance less than 45 mL/min. Patients unable to stop using aspirin or other non-steroidal anti-inflammatory drugs or unable to take folic acid, vitamin B₁₂ or corticosteroids were also excluded from the study.

The data described below reflect exposure to Pemetrexed for Injection in 333 patients in PARAMOUNT. Median age was 61 years (range 32 to 83 years); 58% of patients were men; 94% were White, 4.8% were Asian, and <1% were Black or African American; 36% had an ECOG PS 0. The median number of maintenance cycles was 4 for Pemetrexed for Injection and placebo arms. Dose reductions for adverse reactions occurred in 3.3% of patients in the Pemetrexed for Injection arm and 0.6% in the placebo arm. Dose delays for adverse reactions occurred in 22% of patients in the Pemetrexed for Injection arm and 16% in the placebo arm.

Table 6 provides the frequency and severity of adverse reactions reported in ≥5% of the 333 Pemetrexed for Injection-treated patients in PARAMOUNT.

Table 6: Adverse Reactions Occurring in ≥5% of Patients Receiving Pemetrexed for Injection in PARAMOUNT

	Pemetrexed for Injection (N=333)	Placebo (N=167)
--	---	------------------------

Adverse Reaction*	Injection (N=333)		(N=167)	
	All Grades (%)	Grade 3-4 (%)	All Grades (%)	Grades 3-4 (%)
All adverse reactions	53	17	34	4.8
1. Laboratory				
1. Hematologic				
1. Anemia	15	4.8	4.8	0.6
1. Neutropenia	9	3.9	0.6	0
Clinical				
1. Constitutional symptoms				
1. Fatigue	18	4.5	11	0.6
1. Gastrointestinal				
1. Nausea	12	0.3	2.4	0
1. Vomiting	6	0	1.8	0
1. Mucositis/stomatitis	5	0.3	2.4	0
1. General disorders				
1. Edema	5	0	3.6	0

* NCI CTCAE version 3.0.

The requirement for red blood cell (13% versus 4.8%) and platelet (1.5% versus 0.6%) transfusions, erythropoiesis stimulating agents (12% versus 7%), and granulocyte colony stimulating factors (6% versus 0%) were higher in the Pemetrexed for Injection arm compared to the placebo arm.

The following additional Grade 3 or 4 adverse reactions were observed more frequently in the Pemetrexed for Injection arm.

1. Incidence 1% to <5%

Blood/Bone Marrow— thrombocytopenia

General Disorders— febrile neutropenia

2. Incidence <1%

Cardiovascular— ventricular tachycardia, syncope

General Disorders— pain

Gastrointestinal— gastrointestinal obstruction

Neurologic— depression

Renal— renal failure

Vascular— pulmonary embolism

Treatment of Recurrent Disease After Prior Chemotherapy

The safety of Pemetrexed for Injection was evaluated in Study JMEI, a randomized (1:1), open-label, active-controlled trial conducted in patients who had progressed following platinum-based chemotherapy. Patients received Pemetrexed for Injection 500 mg/m² intravenously or docetaxel 75 mg/m² intravenously on Day 1 of each 21-day cycle. All patients on the Pemetrexed for Injection arm received folic acid and vitamin B₁₂ supplementation.

Study JMEI excluded patients with an ECOG PS of 3 or greater, uncontrolled third-space fluid retention, inadequate bone marrow reserve and organ function, or a calculated creatinine clearance less than 45 mL/min. Patients unable to discontinue aspirin or other non-steroidal anti-inflammatory drugs or unable to take folic acid, vitamin B₁₂ or corticosteroids were also excluded from the study.

The data described below reflect exposure to Pemetrexed for Injection in 265 patients in Study JMEI. Median age was 58 years (range 22 to 87 years); 73% of patients were men; 70% were White, 24% were Asian, 2.6% were Black or African American, 1.8% were Hispanic or Latino, and <2% were other ethnicities; 19% had an ECOG PS 0.

Table 7 provides the frequency and severity of adverse reactions reported in ≥5% of the 265 Pemetrexed for Injection-treated patients in Study JMEI. Study JMEI is not designed to demonstrate a statistically significant reduction in adverse reaction rates for Pemetrexed for Injection, as compared to the control arm, for any specified adverse reaction listed in the Table 7 below.

Table 7: Adverse Reactions Occurring in ≥5% of Fully Supplemented Patients Receiving Pemetrexed for Injection in Study JMEI

Adverse Reaction*	Pemetrexed for Injection (N=265)		Docetaxel (N=276)	
	All Grades (%)	Grades 3-4 (%)	All Grade (%)	Grades 3-4 (%)
1. Laboratory				
1. Hematologic				
1. Anemia	19	4	22	4
1. Neutropenia	11	5	45	40
1. Thrombocytopenia	8	2	1	0
1. Hepatic				
1. Increased ALT	8	2	1	0
1. Increased AST	7	1	1	0
1. Clinical				

1. Gastrointestinal				
1. Nausea	31	3	17	2
1. Anorexia	22	2	24	3
1. Vomiting	16	2	12	1
1. Stomatitis/pharyngitis	15	1	17	1
1. Diarrhea	13	0	24	3
1. Constipation	6	0	4	0
1. Constitutional symptoms				
1. Fatigue	34	5	36	5
1. Fever	8	0	8	0
1. Dermatology/Skin				
1. Rash/desquamation	14	0	6	0
1. Pruritus	7	0	2	0
1. Alopecia	6	1	38	2

* NCI CTCAE version 2.0.

The following additional adverse reactions were observed in patients assigned to receive Pemetrexed for Injection.

1. Incidence 1% to <5%

Body as a Whole— abdominal pain, allergic reaction/hypersensitivity, febrile neutropenia, infection

Dermatology/Skin— erythema multiforme

Neurology— motor neuropathy, sensory neuropathy

2. Incidence <1%

Cardiovascular— supraventricular arrhythmias *Renal*— renal failure

Mesothelioma

The safety of Pemetrexed for Injection was evaluated in Study JMCH, a randomized (1:1), single-blind study conducted in patients with MPM who had received no prior chemotherapy for MPM. Patients received Pemetrexed for Injection 500 mg/m² intravenously in combination with cisplatin 75 mg/m² intravenously on Day 1 of each 21-day cycle or cisplatin 75 mg/m² intravenously on Day 1 of each 21-day cycle administered until disease progression or unacceptable toxicity. Safety was assessed in

226 patients who received at least one dose of Pemetrexed for Injection in combination with cisplatin and 222 patients who received at least one dose of cisplatin alone. Among 226 patients who received Pemetrexed for Injection in combination with cisplatin, 74% (n=168) received full supplementation with folic acid and vitamin B₁₂ during study therapy, 14% (n=32) were never supplemented, and 12% (n=26) were partially supplemented.

Study JMCH excluded patients with Karnofsky Performance Scale (KPS) of less than 70, inadequate bone marrow reserve and organ function, or a calculated creatinine clearance less than 45 mL/min. Patients unable to stop using aspirin or other non-steroidal anti-inflammatory drugs were also excluded from the study.

The data described below reflect exposure to Pemetrexed for Injection in 168 patients that were fully supplemented with folic acid and vitamin B₁₂. Median age was 60 years (range 19 to 85 years); 82% were men; 92% were White, 5% were Hispanic or Latino, 3.0% were Asian, and <1% were other ethnicities; 54% had KPS of 90-100. The median number of treatment cycles administered was 6 in the Pemetrexed for Injection/cisplatin fully supplemented group and 2 in the Pemetrexed for Injection/cisplatin never supplemented group. Patients receiving Pemetrexed for Injection in the fully supplemented group had a relative dose intensity of 93% of the protocol-specified Pemetrexed for Injection dose intensity. The most common adverse reaction resulting in dose delay was neutropenia.

Table 8 provides the frequency and severity of adverse reactions $\geq 5\%$ in the subgroup of Pemetrexed for Injection-treated patients who were fully vitamin supplemented in Study JMCH. Study JMCH was not designed to demonstrate a statistically significant reduction in adverse reaction rates for Pemetrexed for Injection, as compared to the control arm, for any specified adverse reaction listed in the table below.

Table 8: Adverse Reactions Occurring in $\geq 5\%$ of Fully Supplemented Subgroup of Patients Receiving Pemetrexed for Injection/Cisplatin in Study JMCH *

Adverse Reaction [†]	Pemetrexed for Injection/cisplatin (N=168)		Cisplatin (N=163)	
	All Grades (%)	Grade 3-4 (%)	All Grades (%)	Grade 3-4 (%)
1. Laboratory				
1. Hematologic				
1. Neutropenia	56	23	13	3
1. Anemia	26	4	10	0
1. Thrombocytopenia	23	5	9	0
1. Renal				

1. Elevated creatinine	11	1	10	1
1. Decreased creatinine clearance	16	1	18	2
1. Clinical				
1. Eye Disorder				
1. Conjunctivitis	5	0	1	0
1. Gastrointestinal				
1. Nausea	82	12	77	6
1. Vomiting	57	11	50	4
1. Stomatitis/pharyngitis	23	3	6	0
1. Anorexia	20	1	14	1
1. Diarrhea	17	4	8	0
1. Constipation	12	1	7	1
1. Dyspepsia	5	1	1	0
1. Constitutional Symptoms				
1. Fatigue	48	10	42	9
1. Metabolism and Nutrition				
1. Dehydration	7	4	1	1
1. Neurology				
1. Sensory neuropathy	10	0	10	1
1. Taste disturbance	8	0	6	0
1. Dermatology/Skin				
1. Rash	16	1	5	0
1. Alopecia	11	0	6	0

* In Study JMCH, 226 patients received at least one dose of Pemetrexed for Injection in combination with cisplatin and 222 patients received at least one dose of cisplatin. Table 8 provides the ADRs

for subgroup of patients treated with Pemetrexed for Injection in combination with cisplatin (168 patients) or cisplatin alone (163 patients) who received full supplementation with folic acid and vitamin B 12 during study therapy.

† NCI CTCAE version 2.0.

The following additional adverse reactions were observed in patients receiving Pemetrexed for Injection plus cisplatin:

1. **Incidence 1% to <5%**

Body as a Whole— febrile neutropenia, infection, pyrexia

Dermatology/Skin— urticaria

General Disorders— chest pain

Metabolism and Nutrition— increased AST, increased ALT, increased GGT

Renal— renal failure

2. **Incidence <1%**

Cardiovascular— arrhythmia

Neurology— motor neuropathy

Exploratory Subgroup Analyses based on Vitamin Supplementation

Table 9 provides the results of exploratory analyses of the frequency and severity of NCI CTCAE Grade 3 or 4 adverse reactions reported in more Pemetrexed for Injection-treated patients who did not receive vitamin supplementation (never supplemented) as compared with those who received vitamin supplementation with daily folic acid and vitamin B₁₂ from the time of enrollment in Study JMCH (fully-supplemented).

Table 9: Exploratory Subgroup Analysis of Selected Grade 3/4 Adverse Reactions Occurring in Patients Receiving Pemetrexed for Injection in Combination with Cisplatin with or without Full Vitamin Supplementation in Study JMCH *

Grade 3-4 Adverse Reactions	Fully Supplemented Patients N=168 (%)	Never Supplemented Patients N=32 (%)
Neutropenia	23	38
Thrombocytopenia	5	9
Vomiting	11	31
Febrile neutropenia	1	9
Infection with Grade 3/4 neutropenia	0	6
Diarrhea	4	9

* NCI CTCAE version 2.0.

The following adverse reactions occurred more frequently in patients who were fully vitamin supplemented than in patients who were never supplemented:

- hypertension (11% versus 3%),
- chest pain (8% versus 6%),
- thrombosis/embolism (6% versus 3%).

Additional Experience Across Clinical Trials

1. *Sepsis, with or without neutropenia, including fatal cases:*1%
Severe esophagitis, resulting in hospitalization: <1%

6.2 Postmarketing Experience

The following adverse reactions have been identified during post-approval use of Pemetrexed for Injection. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

1. *Blood and Lymphatic System*— immune-mediated hemolytic anemia
Gastrointestinal— colitis, pancreatitis
General Disorders and Administration Site Conditions— edema
Injury, poisoning, and procedural complications— radiation recall
Respiratory— interstitial pneumonitis
Skin— Serious and fatal bullous skin conditions, Stevens-Johnson syndrome, and toxic epidermal necrolysis

7 DRUG INTERACTIONS

Effects of Ibuprofen on Pemetrexed

Ibuprofen increases exposure (AUC) of pemetrexed [see *Clinical Pharmacology (12.3)*]. In patients with creatinine clearance between 45 mL/min and 79 mL/min:

- Avoid administration of ibuprofen for 2 days before, the day of, and 2 days following administration of Pemetrexed for Injection [see *Dosage and Administration (2.5)*].
- Monitor patients more frequently for myelosuppression, renal, and gastrointestinal toxicity, if concomitant administration of ibuprofen cannot be avoided.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

Based on findings from animal studies and its mechanism of action, Pemetrexed for Injection can cause fetal harm when administered to a pregnant woman [see *Clinical Pharmacology (12.1)*]. There are no available data on Pemetrexed for Injection use in pregnant women. In animal reproduction studies, intravenous administration of pemetrexed to pregnant mice during the period of organogenesis was teratogenic, resulting in developmental delays and malformations at doses lower than the recommended human dose of 500 mg/m² [see *Data*]. Advise pregnant women of the potential risk to a fetus [see *Use in Special Populations (8.3)*].

In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2 to 4% and 15 to 20%, respectively.

Data

Animal Data

Pemetrexed was teratogenic in mice. Daily dosing of pemetrexed by intravenous injection to pregnant mice during the period of organogenesis increased the incidence of fetal malformations (cleft palate; protruding tongue; enlarged or misshaped kidney; and

fused lumbar vertebra) at doses (based on BSA) 0.03 times the human dose of 500 mg/m². At doses, based on BSA, greater than or equal to 0.0012 times the 500 mg/m² human dose, pemetrexed administration resulted in dose-dependent increases in developmental delays (incomplete ossification of talus and skull bone; and decreased fetal weight).

8.2 Lactation

Risk Summary

There is no information regarding the presence of pemetrexed or its metabolites in human milk, the effects on the breastfed infant, or the effects on milk production. Because of the potential for serious adverse reactions in breastfed infants from Pemetrexed for Injection, advise women not to breastfeed during treatment with Pemetrexed for Injection and for one week after last dose.

8.3 Females and Males of Reproductive Potential

Based on animal data Pemetrexed for Injection can cause malformations and developmental delays when administered to a pregnant woman [see *Use in Specific Populations (8.1)*].

1. Pregnancy Testing

Verify pregnancy status of females of reproductive potential prior to initiating Pemetrexed for Injection [see *Use in Specific Populations (8.1)*].

Contraception

Females

Because of the potential for genotoxicity, advise females of reproductive potential to use effective contraception during treatment with Pemetrexed for Injection for 6 months after the last dose.

Males

Because of the potential for genotoxicity, advise males with female partners of reproductive potential to use effective contraception during treatment with Pemetrexed for Injection and for 3 months after the last dose [see *Nonclinical Toxicology (13.1)*].

Infertility

Males

Pemetrexed for Injection may impair fertility in males of reproductive potential. It is not known whether these effects on fertility are reversible [see *Nonclinical Toxicology (13.1)*].

8.4 Pediatric Use

The safety and effectiveness of Pemetrexed for Injection in pediatric patients have not been established. The safety and pharmacokinetics of Pemetrexed for Injection were evaluated in two clinical studies conducted in pediatric patients with recurrent solid tumors (NCT00070473 N=32 and NCT00520936 N=72). Patients in both studies received concomitant vitamin B₁₂ and folic acid supplementation and dexamethasone.

No tumor responses were observed. Adverse reactions observed in pediatric patients were similar to those observed in adults.

Single-dose pharmacokinetics of Pemetrexed for Injection were evaluated in 22 patients

age 4 to 18 years enrolled in NCT00070473 were within range of values in adults.

8.5 Geriatric Use

Of the 3,946 patients enrolled in clinical studies of Pemetrexed for Injection, 34% were 65 and over and 4% were 75 and over. No overall differences in effectiveness were observed between these patients and younger patients. The incidences of Grade 3-4 anemia, fatigue, thrombocytopenia, hypertension, and neutropenia were higher in patients 65 years of age and older as compared to younger patients: in at least one of five randomized clinical trials. [see *Adverse Reactions (6.1)* and *Clinical Studies (14.1, 14.2)*].

8.6 Patients with Renal Impairment


Pemetrexed for Injection is primarily excreted by the kidneys. Decreased renal function results in reduced clearance and greater exposure (AUC) to Pemetrexed for Injection compared with patients with normal renal function [*Warnings and Precautions (5.2, 5.6)* and *Clinical Pharmacology (12.3)*] . No dose is recommended for patients with creatinine clearance less than 45 mL/min [see *Dosage and Administration (2.3)*].

10 OVERDOSAGE

No drugs are approved for the treatment of Pemetrexed for Injection overdose. Based on animal studies, administration of leucovorin may mitigate the toxicities of Pemetrexed for Injection overdose. It is not known whether pemetrexed is dialyzable.

11 DESCRIPTION

Pemetrexed for Injection is a folate analog metabolic inhibitor. The drug substance, pemetrexed disodium heptahydrate, has the chemical name L-glutamic acid, *N*-[4-[2-(2-amino-4,7-dihydro-4-oxo-1 *H*-pyrrolo[2,3-*d*]pyrimidin-5-yl)ethyl]benzoyl]-, disodium salt, heptahydrate with a molecular formula of $C_{20}H_{19}N_5Na_2O_6 \cdot 7H_2O$ and a molecular weight of 597.49. The structural formula is as follows:



Pemetrexed for Injection is a sterile white-to-light yellow or green-yellow lyophilized powder in single-dose vials to be reconstituted for intravenous infusion. Each 100-mg vial of Pemetrexed for Injection contains 100 mg pemetrexed (equivalent to 139.8 mg pemetrexed disodium heptahydrate) and 106 mg mannitol. Each 500-mg vial of

Pemetrexed for Injection contains 500 mg pemetrexed (equivalent to 699 mg pemetrexed disodium heptahydrate) and 500 mg mannitol. Hydrochloric acid and/or sodium hydroxide may have been added to adjust pH.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Pemetrexed for Injection is a folate analog metabolic inhibitor that disrupts folate-dependent metabolic processes essential for cell replication. In vitro studies show that pemetrexed inhibits thymidylate synthase (TS), dihydrofolate reductase, and glycinamide ribonucleotide formyltransferase (GARFT), which are folate-dependent enzymes involved in the de novo biosynthesis of thymidine and purine nucleotides. Pemetrexed is taken into cells by membrane carriers such as the reduced folate carrier and membrane folate binding protein transport systems. Once in the cell, pemetrexed is converted to polyglutamate forms by the enzyme folylpolyglutamate synthetase. The polyglutamate forms are retained in cells and are inhibitors of TS and GARFT.

12.2 Pharmacodynamics

Pemetrexed inhibited the in vitro growth of mesothelioma cell lines (MSTO-211H, NCI-H2052) and showed synergistic effects when combined with cisplatin.

Based on population pharmacodynamic analyses, the depth of the absolute neutrophil counts (ANC) nadir correlates with the systemic exposure to pemetrexed and supplementation with folic acid and vitamin B₁₂. There is no cumulative effect of pemetrexed exposure on ANC nadir over multiple treatment cycles.

12.3 Pharmacokinetics

Absorption

The pharmacokinetics of pemetrexed when Pemetrexed for Injection was administered as a single agent in doses ranging from 0.2 to 838 mg/m² infused over a 10-minute period have been evaluated in 426 cancer patients with a variety of solid tumors. Pemetrexed total systemic exposure (AUC) and maximum plasma concentration (C_{max}) increased proportionally with increase of dose. The pharmacokinetics of pemetrexed did not change over multiple treatment cycles.

Distribution

Pemetrexed has a steady-state volume of distribution of 16.1 liters. In vitro studies indicated that pemetrexed is 81% bound to plasma proteins.

Elimination

The total systemic clearance of pemetrexed is 91.8 mL/min and the elimination half-life of pemetrexed is 3.5 hours in patients with normal renal function (creatinine clearance of 90 mL/min). As renal function decreases, the clearance of pemetrexed decreases and exposure (AUC) of pemetrexed increases.

Metabolism

Pemetrexed is not metabolized to an appreciable extent.

Excretion

Pemetrexed is primarily eliminated in the urine, with 70% to 90% of the dose recovered

unchanged within the first 24 hours following administration. In vitro studies indicated that pemetrexed is a substrate of OAT3 (organic anion transporter 3), a transporter that is involved in the active secretion of pemetrexed.

Specific Populations

Age (26 to 80 years) and sex had no clinically meaningful effect on the systemic exposure of pemetrexed based on population pharmacokinetic analyses.

Racial Groups

The pharmacokinetics of pemetrexed were similar in Whites and Blacks or African Americans. Insufficient data are available for other ethnic groups.

Patients with Hepatic Impairment

Pemetrexed has not been formally studied in patients with hepatic impairment. No effect of elevated AST, ALT, or total bilirubin on the PK of pemetrexed was observed in clinical studies.

Patients with Renal Impairment

Pharmacokinetic analyses of pemetrexed included 127 patients with impaired renal function. Plasma clearance of pemetrexed decreases as renal function decreases, with a resultant increase in systemic exposure. Patients with creatinine clearances of 45, 50, and 80 mL/min had 65%, 54%, and 13% increases, respectively in systemic exposure (AUC) compared to patients with creatinine clearance of 100 mL/min [see *Dosage and Administration (2.3)* and *Warnings and Precautions (5.2)*].

Third-Space Fluid

The pemetrexed plasma concentrations in patients with various solid tumors with stable, mild to moderate third-space fluid were comparable to those observed in patients without third space fluid collections. The effect of severe third space fluid on pharmacokinetics is not known.

Drug Interaction Studies

Drugs Inhibiting OAT3 Transporter

Ibuprofen, an OAT3 inhibitor, administered at 400 mg four times a day decreased the clearance of pemetrexed and increased its exposure (AUC) by approximately 20% in patients with normal renal function (creatinine clearance >80 mL/min).

In Vitro Studies

Pemetrexed is a substrate for OAT3. Ibuprofen, an OAT3 inhibitor inhibited the uptake of pemetrexed in OAT3-expressing cell cultures with an average $[I_{50}]/IC_{50}$ ratio of 0.38. In vitro data predict that at clinically relevant concentrations, other NSAIDs (naproxen, diclofenac, celecoxib) would not inhibit the uptake of pemetrexed by OAT3 and would not increase the AUC of pemetrexed to a clinically significant extent. [see *Drug Interactions (7)*].

Pemetrexed is a substrate for OAT4. In vitro, ibuprofen and other NSAIDs (naproxen, diclofenac, celecoxib) are not inhibitors of OAT4 at clinically relevant concentrations.

Aspirin

Aspirin, administered in low to moderate doses (325 mg every 6 hours), does not affect the pharmacokinetics of pemetrexed.

Cisplatin

Cisplatin does not affect the pharmacokinetics of pemetrexed and the pharmacokinetics

of total platinum are unaltered by pemetrexed.

Vitamins

Neither folic acid nor vitamin B₁₂ affect the pharmacokinetics of pemetrexed.

Drugs Metabolized by Cytochrome P450 Enzymes

In vitro studies suggest that pemetrexed does not inhibit the clearance of drugs metabolized by CYP3A, CYP2D6, CYP2C9, and CYP1A2.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

No carcinogenicity studies have been conducted with pemetrexed. Pemetrexed was clastogenic in an in vivo micronucleus assay in mouse bone marrow but was not mutagenic in multiple in vitro tests (Ames assay, Chinese Hamster Ovary cell assay).

Pemetrexed administered intraperitoneally at doses of ≥ 0.1 mg/kg/day to male mice (approximately 0.0006 times the recommended human dose based on BSA) resulted in reduced fertility, hypospermia, and testicular atrophy.

14 CLINICAL STUDIES

14.1 Non-Squamous NSCLC

Initial Treatment in Combination with Pembrolizumab and Platinum

The efficacy of Pemetrexed for Injection in combination with pembrolizumab and platinum chemotherapy was investigated in Study KEYNOTE-189 (NCT02578680), a randomized, multicenter, double-blind, active-controlled trial conducted in patients with metastatic non-squamous NSCLC, regardless of PD-L1 tumor expression status, who had not previously received systemic therapy for metastatic disease and in whom there were no EGFR or ALK genomic tumor aberrations. Patients with autoimmune disease that required systemic therapy within 2 years of treatment; a medical condition that required immunosuppression; or who had received more than 30 Gy of thoracic radiation within the prior 26 weeks were ineligible. Randomization was stratified by smoking status (never versus former/current), choice of platinum (cisplatin versus carboplatin), and tumor PD-L1 status (TPS <1% [negative] versus TPS $\geq 1\%$). Patients were randomized (2:1) to one of the following treatment arms:

- Pemetrexed for Injection 500 mg/m², pembrolizumab 200 mg, and investigator's choice of cisplatin 75 mg/m² or carboplatin AUC 5 mg/mL/min intravenously on Day 1 of each 21-day cycle for 4 cycles followed by Pemetrexed for Injection 500 mg/m² and pembrolizumab 200 mg intravenously every 3 weeks. Pemetrexed for Injection was administered after pembrolizumab and prior to platinum chemotherapy on Day 1.
- Placebo, Pemetrexed for Injection 500 mg/m², and investigator's choice of cisplatin 75 mg/m² or carboplatin AUC 5 mg/mL/min intravenously on Day 1 of each 21-day cycle for 4 cycles followed by placebo and Pemetrexed for Injection 500 mg/m² intravenously every 3 weeks.

Treatment with Pemetrexed for Injection continued until RECIST v1.1 (modified to follow

a maximum of 10 target lesions and a maximum of 5 target lesions per organ)-defined progression of disease as determined by the investigator or unacceptable toxicity. Patients randomized to placebo, Pemetrexed for Injection, and platinum chemotherapy were offered pembrolizumab as a single agent at the time of disease progression.

Assessment of tumor status was performed at Week 6, Week 12, and then every 9 weeks thereafter. The main efficacy outcome measures were OS and PFS as assessed by BICR RECIST v1.1, modified to follow a maximum of 10 target lesions and a maximum of five target lesions per organ. Additional efficacy outcome measures were ORR and duration of response, as assessed by the BICR according to RECIST v1.1, modified to follow a maximum of 10 target lesions and a maximum of 5 target lesions per organ.

A total of 616 patients were randomized: 410 patients to the Pemetrexed for Injection, pembrolizumab, and platinum chemotherapy arm and 206 to the placebo, Pemetrexed for Injection, and platinum chemotherapy arm. The study population characteristics were: median age of 64 years (range: 34 to 84); 49% age 65 or older; 59% male; 94% White and 3% Asian; 56% ECOG performance status of 1; and 18% with history of brain metastases. Thirty-one percent had tumor PD-L1 expression TPS <1%. Seventy-two percent received carboplatin and 12% were never smokers. A total of 85 patients in the placebo, Pemetrexed for Injection, and chemotherapy arm received an anti-PD-1/PD-L1 monoclonal antibody at the time of disease progression.

The trial demonstrated a statistically significant improvement in OS and PFS for patients randomized to Pemetrexed for Injection in combination with pembrolizumab and platinum chemotherapy compared with placebo, Pemetrexed for Injection, and platinum chemotherapy (see Table 10 and Figure 1).

Table 10: Efficacy Results of KEYNOTE-189

Endpoint	Pemetrexed for Injection Pembrolizumab Platinum Chemotherapy n=410	Placebo Pemetrexed for Injection Platinum Chemotherapy n=206
OS		
1. Number (%) of patients with event	127 (31%)	108 (52%)
1. Median in months (95% CI)	NR (NR, NR)	11.3 (8.7, 15.1)
1. Hazard ratio *(95% CI)	0.49 (0.38, 0.64)	
1. p-value	<0.0001	
PFS		
1. Number of patients with event (%)	245 (60%)	166 (81%)
1. Median in months (95% CI)	8.8 (7.6, 9.2)	4.9 (4.7, 5.5)

1. Hazard ratio (95% CI)	0.52 (0.43, 0.64)	
1. p-value †	<0.0001	
ORR		
1. Overall response rate ‡(95% CI)	48% (43, 53)	19% (14, 25)
1. Complete response	0.5%	0.5%
1. Partial response	47%	18%
1. p-value §	<0.0001	
Duration of Response		
1. Median in months (range)	11.2 (1.1+, 18.0+)	7.8 (2.1+, 16.4+)


* Based on the stratified Cox proportional hazard model.

† Based on stratified log-rank test.

‡ Response: Best objective response as confirmed complete response or partial response.

§ Based on Miettinen and Nurminen method stratified by PD-L1 status, platinum chemotherapy and smoking status. NR = not reached

At the protocol specified final OS analysis, the median in the Pemetrexed for Injection in combination with pembrolizumab and platinum chemotherapy arm was 22.0 months (95% CI: 19.5, 24.5) compared to 10.6 months (95% CI: 8.7, 13.6) in the placebo with Pemetrexed for Injection and platinum chemotherapy arm, with an HR of 0.56 (95% CI: 0.46, 0.69).



Number at Risk		Time in Months							
	0	3	6	9	12	15	18	21	
PI+P+C	410	377	347	278	163	71	18	0	
PI+C	206	183	149	104	59	25	8	0	

PI+P+C = Pemetrexed for Injection + pembrolizumab + platinum chemotherapy.
 PI+C = Pemetrexed for Injection + platinum chemotherapy + placebo.

*Based on the protocol-specified final OS analysis

Initial Treatment in Combination with Cisplatin

The efficacy of Pemetrexed for Injection was evaluated in Study JMDB (NCT00087711), a multi-center, randomized (1:1), open-label study conducted in 1725 chemotherapy-naive patients with Stage IIIb/IV NSCLC. Patients were randomized to receive Pemetrexed for Injection with cisplatin or gemcitabine with cisplatin. Randomization was stratified by Eastern Cooperative Oncology Group Performance Status (ECOG PS 0 versus 1), gender, disease stage, basis for pathological diagnosis (histopathological/cytopathological), history of brain metastases, and investigative center. Pemetrexed for Injection was administered intravenously over 10 minutes at a dose of 500 mg/m² on Day 1 of each 21-day cycle. Cisplatin was administered intravenously at a dose of 75 mg/m² approximately 30 minutes after Pemetrexed for Injection administration on Day 1 of each cycle, gemcitabine was administered at a dose of 1250 mg/m² on Day 1 and Day 8, and cisplatin was administered intravenously at a dose of 75 mg/m² approximately 30 minutes after administration of gemcitabine, on Day

1 of each 21-day cycle. Treatment was administered up to a total of 6 cycles; patients in both arms received folic acid, vitamin B₁₂, and dexamethasone [see *Dosage and Administration (2.4)*]. The primary efficacy outcome measure was overall survival.

A total of 1725 patients were enrolled with 862 patients randomized to Pemetrexed for Injection in combination with cisplatin and 863 patients to gemcitabine in combination with cisplatin. The median age was 61 years (range 26-83 years), 70% were male, 78% were White, 17% were Asian, 2.9% were Hispanic or Latino, and 2.1% were Black or African American, and <1% were other ethnicities. Among patients for whom ECOG PS (n=1722) and smoking history (n=1516) were collected, 65% had an ECOG PS of 1, 36% had an ECOG PS of 0, and 84% were smokers. For tumor characteristics, 73% had non-squamous NSCLC and 27% had squamous NSCLC; 76% had Stage IV disease. Among 1252 patients with non-squamous NSCLC histology, 68% had a diagnosis of adenocarcinoma, 12% had large cell histology and 20% had other histologic subtypes.


Efficacy results in Study JMDB are presented in Table 11 and Figure 2.

Table 11: Efficacy Results in Study JMDB

Efficacy Parameter	Pemetrexed for Injection plus Cisplatin (N=862)	Gemcitabine plus Cisplatin (N=863)
Overall Survival		
1. Median (months) (95% CI)	10.3 (9.8-11.2)	10.3 (9.6-10.9)
1. Hazard ratio (HR) *† (95% CI)	0.94 (0.84-1.05)	
Progression-Free Survival		
1. Median (months) (95% CI)	4.8 (4.6-5.3)	5.1 (4.6-5.5)
1. Hazard ratio (HR) ^{a,b} (95% CI)	1.04 (0.94-1.15)	
Overall Response Rate (95% CI)	27.1% (24.2-30.1)	24.7% (21.8-27.6)

* Unadjusted for multiple comparisons.

† Adjusted for gender, stage, basis of diagnosis, and performance status.



Patients at Risk		Survival Time (months)									
	0	3	6	9	12	15	18	21	24	27	30
PIC	862	737	598	458	341	235	146	88	45	10	0
GC	863	731	590	456	327	209	139	78	34	14	0

In pre-specified analyses assessing the impact of NSCLC histology on overall survival, clinically relevant differences in survival according to histology were observed. These subgroup analyses are shown in Table 12 and Figure 3 and 4. This difference in treatment effect for Pemetrexed for Injection based on histology demonstrating a lack of efficacy in squamous cell histology was also observed in Studies JMEN and JMEI.


Table 12: Overall Survival in NSCLC Histologic Subgroups in Study JMDB

Histologic Subgroups	Pemetrexed for Injection plus Cisplatin (N=862)	Gemcitabine plus Cisplatin (N=863)
Non-squamous NSCLC (N=1252)		
1. Median (months) (95% CI)	11.0 (10.1-12.5)	10.1 (9.3-10.9)
1. HR *† (95% CI)	0.84 (0.74-0.96)	
Adenocarcinoma (N=847)		
1. Median (months) (95% CI)	12.6 (10.7-13.6)	10.9 (10.2-11.9)
1. HR *† (95% CI)	0.84 (0.71-0.99)	

Large Cell (N=153)		
1. Median (months) (95% CI)	10.4 (8.6-14.1)	6.7 (5.5-9.0)
1. HR ^{*†} (95% CI)	0.67 (0.48-0.96)	
Non-squamous, not otherwise specified (N=252)		
1. Median (months) (95% CI)	8.6 (6.8-10.2)	9.2 (8.1-10.6)
1. HR ^{*†} (95% CI)	1.08 (0.81-1.45)	
Squamous Cell (N=473)		
1. Median (months) (95% CI)	9.4 (8.4-10.2)	10.8 (9.5-12.1)
1. HR ^{*†} (95% CI)	1.23 (1.00-1.51)	


* Unadjusted for multiple comparisons.

† Adjusted for ECOG PS, gender, disease stage, and basis for pathological diagnosis (histopathological/cytopathological).



Patients at Risk

PIC	618	533	437	341	264	188	118	72	37	8	0
GC	634	542	435	339	240	151	101	55	26	10	0



Patients at Risk		0	3	6	9	12	15	18	21	24	27	30
PIC		244	204	161	117	77	47	28	16	8	2	0
GC		229	189	155	117	87	58	38	23	8	4	0

Maintenance Treatment Following First-line Non-Pemetrexed for Injection Containing Platinum-Based Chemotherapy

The efficacy of Pemetrexed for Injection as maintenance therapy following first-line platinum-based chemotherapy was evaluated in Study JMEN (NCT00102804), a multicenter, randomized (2:1), double-blind, placebo-controlled study conducted in 663 patients with Stage IIIb/IV NSCLC who did not progress after four cycles of platinum-based chemotherapy. Patients were randomized to receive Pemetrexed for Injection 500 mg/m² intravenously every 21 days or placebo until disease progression or intolerable toxicity. Patients in both study arms received folic acid, vitamin B₁₂, and dexamethasone [see *Dosage and Administration (2.4)*]. Randomization was carried out using a minimization approach [Pocock and Simon (1975)] using the following factors: gender, ECOG PS (0 versus 1), response to prior chemotherapy (complete or partial response versus stable disease), history of brain metastases (yes versus no), non-platinum component of induction therapy (docetaxel versus gemcitabine versus paclitaxel), and disease stage (IIIb versus IV). The major efficacy outcome measures were progression-free survival based on assessment by independent review and overall survival; both were measured from the date of randomization in Study JMEN.

A total of 663 patients were enrolled with 441 patients randomized to Pemetrexed for Injection and 222 patients randomized to placebo. The median age was 61 years (range 26-83 years); 73% were male; 65% were White, 32% were Asian, 2.9% were Hispanic or Latino, and <2% were other ethnicities; 60% had an ECOG PS of 1; and 73% were current or former smokers. Median time from initiation of platinum-based chemotherapy to randomization was 3.3 months (range 1.6 to 5.1 months) and 49% of the population achieved a partial or complete response to first-line, platinum-based chemotherapy. With


regard to tumor characteristics, 81% had Stage IV disease, 73% had non-squamous NSCLC and 27% had squamous NSCLC. Among the 481 patients with non-squamous NSCLC, 68% had adenocarcinoma, 4% had large cell, and 28% had other histologies.

Efficacy results are presented in Table 13 and Figure 5.

Table 13: Efficacy Results in Study JMEN

Efficacy Parameter	Pemetrexed for Injection	Placebo
Overall survival	N=441	N=222
1. Median (months) (95% CI)	13.4 (11.9-15.9)	10.6 (8.7-12.0)
1. Hazard ratio * (95% CI)	0.79 (0.65-0.95)	
1. p-value	p=0.012	
Progression-free survival per independent review	N=387	N=194
1. Median (months) (95% CI)	4.0 (3.1-4.4)	2.0 (1.5-2.8)
1. Hazard ratio * (95% CI)	0.60 (0.49-0.73)	
1. p-value	p<0.00001	

* Hazard ratios are adjusted for multiplicity but not for stratification variables.



Patients at Risk

PI	441	396	340	274	221	179	141	97	63	45	29	19	11	2	0
Placebo	222	200	160	119	93	76	60	40	29	20	13	6	4	0	0

The results of pre-specified subgroup analyses by NSCLC histology are presented in Table 14 and Figure 6 and 7.


Table 14: Efficacy Results in Study JMEN by Histologic Subgroup

Efficacy Parameter	Overall Survival		Progression-Free Survival Per Independent Review	
	Pemetrexed for Injection (N=441)	Placebo (N=222)	Pemetrexed for Injection (N=387)	Placebo (N=194)
Non-squamous NSCLC (n=481)				
1. Median (months)	15.5	10.3	4.4	1.8
1. HR *	0.70		0.47	
1. (95% CI)	(0.56-0.88)		(0.37-0.60)	
Adenocarcinoma (n=328)				
1. Median (months)	16.8	11.5	4.6	2.7
1. HR *	0.73		0.51	
1. (95% CI)	(0.56-0.96)		(0.38-0.68)	

Large cell carcinoma (n=20)				
1. Median (months)	8.4	7.9	4.5	1.5
1. HR *	0.98		0.40	
1. (95% CI)	(0.36-2.65)		(0.12-1.29)	
Other †(n=133)				
1. Median (months)	11.3	7.7	4.1	1.6
1. HR *	0.61		0.44	
1. (95% CI)	(0.40-0.94)		(0.28-0.68)	
Squamous cell NSCLC (n=182)				
1. Median (months)	9.9	10.8	2.4	2.5
1. HR *	1.07		1.03	
1. (95% CI)	(0.77-1.50)		(0.71-1.49)	


* Hazard ratios are not adjusted for multiplicity

† Primary diagnosis of NSCLC not specified as adenocarcinoma, large cell carcinoma, or squamous cell carcinoma.



Patients at Risk

PI	325	302	265	216	178	141	117	82	51	38	25	15	9	2	0
Placebo	156	140	112	80	63	52	42	28	20	11	7	4	3	0	0



Patients at Risk		0	3	6	9	12	15	18	21	24	27	30	33	36	39
PI	116	94	75	58	43	38	24	15	12	7	4	4	2	0	0
Placebo	66	60	48	39	30	24	18	12	9	9	6	2	1	0	0

Maintenance Treatment Following First-line Pemetrexed for Injection Plus Platinum Chemotherapy

The efficacy of Pemetrexed for Injection as maintenance therapy following first-line platinum-based chemotherapy was also evaluated in PARAMOUNT (NCT00789373), a multi-center, randomized (2:1), double-blind, placebo-controlled study conducted in patients with Stage IIIb/IV non-squamous NSCLC who had completed four cycles of Pemetrexed for Injection in combination with cisplatin and achieved a complete response (CR) or partial response (PR) or stable disease (SD). Patients were required to have an ECOG PS of 0 or 1. Patients were randomized to receive Pemetrexed for Injection 500 mg/m² intravenously every 21 days or placebo until disease progression. Randomization was stratified by response to Pemetrexed for Injection in combination with cisplatin induction therapy (CR or PR versus SD), disease stage (IIIb versus IV), and ECOG PS (0 versus 1). Patients in both arms received folic acid, vitamin B₁₂, and dexamethasone. The main efficacy outcome measure was investigator-assessed progression-free survival (PFS) and an additional efficacy outcome measure was overall survival (OS); PFS and OS were measured from the time of randomization.

A total of 539 patients were enrolled with 359 patients randomized to Pemetrexed for Injection and 180 patients randomized to placebo. The median age was 61 years (range 32 to 83 years); 58% were male; 95% were White, 4.5% were Asian, and <1% were Black or African American; 67% had an ECOG PS of 1; 78% were current or former smokers; and 43% of the population achieved a partial or complete response to first-line, platinum-based chemotherapy. With regard to tumor characteristics, 91% had Stage IV disease, 87% had adenocarcinoma, 7% had large cell, and 6% had other histologies.


Efficacy results for PARAMOUNT are presented in Table 15 and Figure 8.

Table 15: Efficacy Results in PARAMOUNT

Efficacy Parameter	Pemetrexed for Injection (N=359)	Placebo (N=180)
Overall survival		
Median (months) (95% CI)	13.9 (12.8-16.0)	11.0 (10.0-12.5)
Hazard ratio (HR) * (95% CI)	0.78 (0.64-0.96)	
p-value	p=0.02	
Progression-free survival†		
Median (months) (95% CI)	4.1 (3.2-4.6)	2.8 (2.6-3.1)
Hazard ratio (HR) * (95% CI)	0.62 (0.49-0.79)	
p-value	p<0.0001	

* Hazard ratios are adjusted for multiplicity but not for stratification variables.

† Based on investigator’s assessment.



Patients at Risk

PI	359	333	272	235	200	166	138	105	79	43	15	2	0
Placebo	180	169	131	103	78	65	49	35	23	12	8	3	0

Treatment of Recurrent Disease After Prior Chemotherapy

The efficacy of Pemetrexed for Injection was evaluated in Study JMEI (NCT00004881), a multicenter, randomized (1:1), open-label study conducted in patients with Stage III or IV NSCLC that had recurred or progressed following one prior chemotherapy regimen for

advanced disease. Patients were randomized to receive Pemetrexed for Injection 500 mg/m² intravenously or docetaxel 75 mg/m² as a 1-hour intravenous infusion once every 21 days. Patients randomized to Pemetrexed for Injection also received folic acid and vitamin B₁₂. The study was designed to show that overall survival with Pemetrexed for Injection was non-inferior to docetaxel, as the major efficacy outcome measure, and that overall survival was superior for patients randomized to Pemetrexed for Injection compared to docetaxel, as a secondary outcome measure.

A total of 571 patients were enrolled with 283 patients randomized to Pemetrexed for Injection and 288 patients randomized to docetaxel. The median age was 58 years (range 22 to 87 years); 72% were male; 71% were White, 24% were Asian, 2.8% were Black or African American, 1.8% were Hispanic or Latino, and <2% were other ethnicities; 88% had an ECOG PS of 0 or 1. With regard to tumor characteristics, 75% had Stage IV disease; 53% had adenocarcinoma, 30% had squamous histology; 8% large cell; and 9% had other histologic subtypes of NSCLC.

The efficacy results in the overall population and in subgroup analyses based on histologic subtype are provided in Tables 16 and 17, respectively. Study JMEI did not show an improvement in overall survival in the intent-to-treat population. In subgroup analyses, there was no evidence of a treatment effect on survival in patients with squamous NSCLC; the absence of a treatment effect in patients with NSCLC of squamous histology was also observed Studies JMDB and JMEN [see *Clinical Studies (14.1)*].

Table 16: Efficacy Results in Study JMEI

Efficacy Parameter	Pemetrexed for Injection (N=283)	Docetaxel (N=288)
Overall survival		
1. Median (months) (95% CI)	8.3 (7.0-9.4)	7.9 (6.3-9.2)
1. Hazard ratio * (95% CI)	0.99 (0.82-1.20)	
Progression-free survival		
1. Median (months) (95% CI)	2.9 (2.4-3.1)	2.9 (2.7-3.4)
1. Hazard ratio ^a (95% CI)	0.97 (0.82-1.16)	
1. Overall response rate (95% CI)	8.5% (5.2-11.7)	8.3% (5.1-11.5)

* Hazard ratios are not adjusted for multiplicity or for stratification variables.

Table 17: Exploratory Efficacy Analyses by Histologic Subgroup in Study JMEI

Histologic Subgroups	Pemetrexed for Injection (N=283)	Docetaxel (N=288)
Non-squamous NSCLC (N=399)		
1. Median (months) (95% CI)	9.3 (7.8-9.7)	8.0 (6.3-9.3)
1. HR * (95% CI)	0.89 (0.71-1.13)	
Adenocarcinoma (N=301)		
1. Median (months) (95% CI)	9.0 (7.6-9.6)	9.2 (7.5-11.3)
1. HR * (95% CI)	1.09 (0.83-1.44)	
Large Cell (N=47)		
1. Median (months) (95% CI)	12.8 (5.8-14.0)	4.5 (2.3-9.1)
1. HR * (95% CI)	0.38 (0.18-0.78)	
Other †(N=51)		
1. Median (months) (95% CI)	9.4 (6.0-10.1)	7.9 (4.0-8.9)
1. HR * (95% CI)	0.62 (0.32-1.23)	
Squamous NSCLC (N=172)		
1. Median (months) (95% CI)	6.2 (4.9-8.0)	7.4 (5.6-9.5)
1. HR * (95% CI)	1.32 (0.93-1.86)	

* Hazard ratio unadjusted for multiple comparisons.

† Primary diagnosis of NSCLC not specified as adenocarcinoma, large cell carcinoma, or squamous cell carcinoma.

14.2 Mesothelioma

The efficacy of Pemetrexed for Injection was evaluated in Study JMCH (NCT00005636), a multicenter, randomized (1:1), single-blind study conducted in patients with MPM who had received no prior chemotherapy. Patients were randomized (n=456) to receive Pemetrexed for Injection 500 mg/m² intravenously over 10 minutes followed 30 minutes

later by cisplatin 75 mg/m² intravenously over two hours on Day 1 of each 21-day cycle or to receive cisplatin 75 mg/m² intravenously over 2 hours on Day 1 of each 21-day cycle; treatment continued until disease progression or intolerable toxicity. The study was modified after randomization and treatment of 117 patients to require that all patients receive folic acid 350 mcg to 1000 mcg daily beginning 1 to 3 weeks prior to the first dose of Pemetrexed for Injection and continuing until 1 to 3 weeks after the last dose, vitamin B₁₂ 1000 mcg intramuscularly 1 to 3 weeks prior to first dose of Pemetrexed for Injection and every 9 weeks thereafter, and dexamethasone 4 mg orally, twice daily, for 3 days starting the day prior to each Pemetrexed for Injection dose. Randomization was stratified by multiple baseline variables including KPS, histologic subtype (epithelial, mixed, sarcomatoid, other), and gender. The major efficacy outcome measure was overall survival and additional efficacy outcome measures were time to disease progression, overall response rate, and response duration.

A total of 448 patients received at least one dose of protocol-specified therapy; 226 patients were randomized to and received at least one dose of Pemetrexed for Injection plus cisplatin, and 222 patients were randomized to and received cisplatin. Among the 226 patients who received cisplatin with Pemetrexed for Injection, 74% received full supplementation with folic acid and vitamin B₁₂ during study therapy, 14% were never supplemented, and 12% were partially supplemented. Across the study population, the median age was 61 years (range: 20 to 86 years); 81% were male; 92% were White, 5% were Hispanic or Latino, 3.1% were Asian, and <1% were other ethnicities; and 54% had a baseline KPS score of 90-100% and 46% had a KPS score of 70-80%. With regard to tumor characteristics, 46% had Stage IV disease, 31% Stage III, 15% Stage II, and 7% Stage I disease at baseline; the histologic subtype of mesothelioma was epithelial in 68% of patients, mixed in 16%, sarcomatoid in 10% and other histologic subtypes in 6%. The baseline demographics and tumor characteristics of the subgroup of fully supplemented patients was similar to the overall study population.


The efficacy results from Study JMCH are summarized in Table 18 and Figure 9.

Table 18: Efficacy Results in Study JMCH

Efficacy Parameter	All Randomized and Treated Patients (N=448)		Fully Supplemented Patients (N=331)	
	Pemetrexed for Injection/Cisplatin (N=226)	Cisplatin (N=222)	Pemetrexed for Injection/Cisplatin (N=168)	Cisplatin (N=163)
Median overall survival (months)	12.1	9.3	13.3	10.0
(95% CI)	(10.0-14.4)	(7.8-10.7)	(11.4-14.9)	(8.4-11.9)
Hazard ratio *	0.77		0.75	
Log rank p-value	0.020		NA †	

* Hazard ratios are not adjusted for stratification variables.

† Not a pre-specified analysis.



Patients at Risk		Survival Time (months)										
		0	3	6	9	12	15	18	21	24	27	30
PI + Cisplatin	226	201	166	128	84	50	32	17	8	4	0	0
Cisplatin	222	195	153	104	63	31	21	14	3	1	0	0

Based upon prospectively defined criteria (modified Southwest Oncology Group methodology) the objective tumor response rate for Pemetrexed for Injection plus cisplatin was greater than the objective tumor response rate for cisplatin alone. There was also improvement in lung function (forced vital capacity) in the Pemetrexed for Injection plus cisplatin arm compared to the control arm.

15 REFERENCES

1. "OSHA Hazardous Drugs." *OSHA*. [<https://www.osha.gov/hazardous-drugs>]

16 HOW SUPPLIED/STORAGE AND HANDLING

How Supplied

Pemetrexed for Injection is a white-to-light yellow or green-yellow lyophilized powder supplied in single-dose vials for reconstitution for intravenous infusion.

NDC 0338-0720-01: Carton containing one (1) single-dose vial of 100 mg pemetrexed.

NDC 0338-0722-01: Carton containing one (1) single-dose vial of 500 mg pemetrexed.

Storage and Handling

Store at 25°C (77°F); excursions permitted to 15-30°C (59-86°F) [see USP Controlled Room Temperature].

Pemetrexed for Injection is a hazardous drug. Follow applicable special handling and disposal procedures. ¹

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Patient Information).

Premedication and Concomitant Medication: Instruct patients to take folic acid as directed and to keep appointments for vitamin B₁₂ injections to reduce the risk of treatment-related toxicity. Instruct patients of the requirement to take corticosteroids to reduce the risks of treatment-related toxicity [see *Dosage and Administration (2.4)* and *Warnings and Precautions (5.1)*].

Myelosuppression: Inform patients of the risk of low blood cell counts and instruct them to immediately contact their physician for signs of infection, fever, bleeding, or symptoms of anemia [see *Warnings and Precautions (5.1)*].

Renal Failure: Inform patients of the risks of renal failure, which may be exacerbated in patients with dehydration arising from severe vomiting or diarrhea. Instruct patients to immediately contact their healthcare provider for a decrease in urine output [see *Warnings and Precautions (5.2)*].

Bullous and Exfoliative Skin Disorders: Inform patients of the risks of severe and exfoliative skin disorders. Instruct patients to immediately contact their healthcare provider for development of bullous lesions or exfoliation in the skin or mucous membranes [see *Warnings and Precautions (5.3)*].

Interstitial Pneumonitis: Inform patients of the risks of pneumonitis. Instruct patients to immediately contact their healthcare provider for development of dyspnea or persistent cough [see *Warnings and Precautions (5.4)*].

Radiation Recall: Inform patients who have received prior radiation of the risks of radiation recall. Instruct patients to immediately contact their healthcare provider for development of inflammation or blisters in an area that was previously irradiated [see *Warnings and Precautions (5.5)*].

Increased Risk of Toxicity with Ibuprofen in Patients with Renal Impairment: Advise patients with mild to moderate renal impairment of the risks associated with concomitant ibuprofen use and instruct them to avoid use of all ibuprofen containing products for 2 days before, the day of, and 2 days following administration of Pemetrexed for Injection [see *Dosage and Administration (2.5)*, *Warnings and Precautions (5.6)*, and *Drug Interactions (7)*].

Embryo-Fetal Toxicity: Advise females of reproductive potential and males with female partners of reproductive potential of the potential risk to a fetus [see *Warnings and Precautions (5.7)* and *Use in Specific Populations (8.1)*]. Advise females of reproductive potential to use effective contraception during treatment with Pemetrexed for Injection and for 6 months after the last dose. Advise females to inform their prescriber of a known or suspected pregnancy. Advise males with female partners of reproductive potential to use effective contraception during treatment with Pemetrexed for Injection and for 3 months after the last dose [see *Warnings and Precautions (5.7)* and *Use in Specific Populations (8.3)*].

Lactation: Advise women not to breastfeed during treatment with Pemetrexed for Injection and for 1 week after the last dose [see *Use in Specific Populations (8.2)*].

Manufactured for:

Baxter Logo

**Baxter Healthcare Corporation
Deerfield, IL 60015 USA**

Made in Germany

HA-30-02-266

Baxter is a registered trademark of Baxter International Inc.

Revised: 09/2023

PATIENT INFORMATION
Pemetrexed for Injection
for Intravenous Use

What is Pemetrexed for Injection?

Pemetrexed for Injection is a prescription medicine used to treat:

- **a kind of lung cancer called non-squamous non-small cell lung cancer (NSCLC).** Pemetrexed for Injection is used:
 - as the first treatment in combination with pembrolizumab and platinum chemotherapy when your lung cancer with no abnormal EGFR or ALK gene has spread (advanced NSCLC).
 - as the first treatment in combination with cisplatin when your lung cancer has spread (advanced NSCLC).
 - alone as maintenance treatment after you have received 4 cycles of chemotherapy that contains platinum for first treatment of your advanced NSCLC and your cancer has not progressed.
 - alone when your lung cancer has returned or spread after prior chemotherapy.
- Pemetrexed for Injection is not for use for the treatment of people with squamous cell NSCL.
- **a kind of cancer called malignant pleural mesothelioma.** This cancer affects the lining of the lungs and chest wall. Pemetrexed for Injection is used in combination with cisplatin as the first treatment for malignant pleural mesothelioma that cannot be removed by surgery or you are not able to have surgery.

Pemetrexed for Injection has not been shown to be safe and effective in children.

Do not take Pemetrexed for Injection: if you have had a severe allergic reaction to any medicine that contains pemetrexed.

Before taking Pemetrexed for Injection, tell your healthcare provider about all of your medical conditions, including if you:

- have kidney problems.
- have had radiation therapy.
- are pregnant or plan to become pregnant .Pemetrexed for Injection can harm your unborn baby.
 - **Females who are able to become pregnant:**
Your healthcare provider will check to see if you are pregnant before you start treatment with Pemetrexed for Injection.
You should use effective birth control (contraception) during treatment with Pemetrexed for Injection and for 6 months after the last does. Tell your healthcare provider right away if you become pregnant or think you are pregnant during treatment with Pemetrexed for Injection.
 - **Males** with female partners who are able to become pregnant should use effective birth control (contraception) during treatment with Pemetrexed for

Injection and for 3 months after the last dose.

- Are breastfeeding or plan to breastfeed .It is not known if Pemetrexed for Injection passes into breast milk. Do not breastfeed during treatment with Pemetrexed for Injection and for 1 week after the last dose.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Tell your healthcare provider if you have kidney problems and take a medicine that contains ibuprofen. You should avoid taking ibuprofen for 2 days before, the day of, and 2 days after receiving treatment with Pemetrexed for Injection.

How is Pemetrexed for Injection given?

- **It is very important to take folic acid and vitamin B₁₂ during your treatment with Pemetrexed for Injection to lower your risk of harmful side effects.**
 - Take folic acid exactly as prescribed by your healthcare provider 1 time a day, beginning 7 days (1 week) before your first dose of Pemetrexed for Injection and continue taking folic acid until 21 days (3 weeks) after your last dose of Pemetrexed for Injection.
 - Your healthcare provider will give you vitamin B₁₂ injections during treatment with Pemetrexed for Injection. You will get your first vitamin B₁₂ injection 7 days (1 week) before your first dose of Pemetrexed for Injection, and then every 3 cycles.
- Your healthcare provider will prescribe a medicine called corticosteroid for you to take 2 times a day for 3 days, beginning the day before each treatment with Pemetrexed for Injection.
- Pemetrexed for Injection is given to you by intravenous (IV) infusion into your vein. The infusion is given over 10 minutes.
- Pemetrexed for Injection is usually given once every 21 days (3 weeks).

What are the possible side effects of Pemetrexed for Injection?

Pemetrexed for Injection can cause serious side effects, including:

- **Low blood cell counts.** Low blood cell counts can be severe, including low white blood cell counts (neutropenia), low platelet counts (thrombocytopenia), and low red blood cell counts (anemia). Your healthcare provider will do blood test to check your blood cell counts regularly during your treatment with Pemetrexed for Injection. **Tell your healthcare provider right away if you have any signs of infection, fever, bleeding, or severe tiredness during your treatment with Pemetrexed for Injection.**
- **Kidney problems, including kidney failure.** Pemetrexed for Injection can cause severe kidney problems that can lead to death. Severe vomiting or diarrhea can lead to loss of fluids (dehydration) which may cause kidney problems to become worse. Tell your healthcare provider right away if you have a decrease in amount of urine.
- **Severe skin reactions.** Severe skin reactions that may lead to death can happen with Pemetrexed for Injection. Tell your healthcare provider right away if you develop blisters, skin sores, skin peeling, or painful sores, or ulcers in your mouth, nose, throat or genital area.
- **Lung problems (pneumonitis).** Pemetrexed for Injection can cause serious lung problems that can lead to death. Tell your healthcare provider right away if you get any new or worsening symptoms of shortness of breath, cough, or fever.

- **Radiation recall.** Radiation recall is a skin reaction that can happen in people who have received radiation treatment in the past and are treated with Pemetrexed for Injection. Tell your healthcare provider if you get swelling, blistering, or a rash that looks like a sunburn in an area that was previously treated with radiation.

The most common side effects of Pemetrexed for Injection when given alone are:

• tiredness	• nausea	• loss of appetite
-------------	----------	--------------------

The most common side effects of Pemetrexed for Injection when given with cisplatin are:

• vomiting	• low white blood cell counts (neutropenia)
• swelling or sores in your mouth or sore throat	• low platelet counts (thrombocytopenia)
• constipation	• low red blood cell counts (anemia)

The most common side effects of Pemetrexed for Injection when given with pembrolizumab and platinum chemotherapy are:

• tiredness/weakness	• nausea
• constipation	• diarrhea
• loss of appetite	• rash
• vomiting	• cough
• shortness of breath	• fever

Pemetrexed for Injection may cause fertility problems in males. This may affect your ability to father a child. It is not known if these effects are reversible. Talk to your healthcare provider if this is a concern for you.

Your healthcare provider will do blood test to check for side effects during treatment with Pemetrexed for Injection. Your healthcare provider may change your dose of Pemetrexed for Injection, delay treatment, or stop treatment if you have certain side effects.

Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

These are not all the side effects of Pemetrexed for Injection. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

General information about the safe and effective use of Pemetrexed for Injection.

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet.

You can ask your pharmacist or healthcare provider for information about Pemetrexed for Injection that is written for health professionals.

What are the ingredients in Pemetrexed for Injection?

Active ingredient: pemetrexed

Inactive ingredients: mannitol, hydrochloric acid and/or sodium hydroxide may have been added to adjust pH.

Manufactured for:

Baxter Logo

Baxter Healthcare Corporation

Deerfield, IL 60015 USA

This Patient Information has been approved by the U.S. Food and Drug Administration.
Revised: September 2023

PACKAGE/LABEL PRINCIPAL DISPLAY PANEL

Single-Dose Vial NDC 0338-0720-01
Rx only

Pemetrexed
for Injection, USP
100 mg per vial
For intravenous use only.
Discard unused portion.
Baxter

Caution: Cytotoxic Agent
Each vial contains pemetrexed disodium equivalent to 100 mg pemetrexed, 106 mg of mannitol. Hydrochloric acid and/or sodium hydroxide may have been added to adjust pH.
Storage: Store powder at 25°C (77°F); excursions permitted to 15-30°C (59-86°F) [see USP Controlled Room Temperature].
See accompanying literature for storage of reconstituted and infusion solutions.
Manufactured for: Baxter Healthcare Corporation, Deerfield, IL 60015 USA. Product of Germany.

LOT/EXP:

(01)00303380720019
USA WA-65-01-887 C55

LOT/EXP:

Single-Dose Vial

Rx only

NDC 0338-0720-01

Pemetrexed

for Injection, USP

100 mg per vial

For intravenous use only.

Discard unused portion.

Baxter Logo

Caution: Cytotoxic Agent

Each vial contains pemetrexed disodium

equivalent to 100 mg pemetrexed, 106 mg of mannitol. Hydrochloric acid and/or sodium hydroxide may have been added to adjust pH. Storage: Store powder at 25°C (77°F); excursions permitted to 15-30°C (59-86°F) [see USP Controlled Room Temperature]. See accompanying literature for storage of reconstituted and infusion solutions.


Manufactured for: **Baxter Healthcare Corporation**, Deerfield, IL 60015 USA. Product of Germany .

Barcode

(01)00303380720019

2D Barcode

USA HA-65-01-837 C 55



Pemetrexed
for Injection, USP
100 mg per vial

Barcode

(01) 20303380720013

Baxter is a registered trademark of
Baxter International Inc.

GTIN:

SN:

LOT:

EXP:

Single-Dose Vial
NDC 0338-0720-01

Pemetrexed
for Injection, USP
100 mg per vial

Barcode

HA-80-03-188
USA

Pemetrexed
for Injection, USP

Each vial contains pemetrexed disodium equivalent to 100 mg pemetrexed, 106 mg of mannitol. Hydrochloric acid and/or sodium hydroxide may have been added to adjust pH.

To Reconstitute: Add 4.2 mL of 0.9% Sodium Chloride Injection (preservative free) to make a solution containing 25 mg/mL pemetrexed. Reconstituted solution must be further diluted before use (see accompanying literature). Administer infusion solution within 24 hours after initial reconstitution. Discard unused portion. See accompanying literature for dosage.

C
498

Single-Dose Vial
NDC 0338-0720-01

Pemetrexed
for Injection, USP
100 mg per vial

For intravenous use only.

Rx only

Baxter Logo

2335B0115

Barcode

Pemetrexed

for Injection, USP

Storage: Store powder at 25°C (77°F); excursions permitted to 15-30°C (59-86°F) [see USP Controlled Room Temperature]. Store reconstituted and infusion solutions at refrigerated, 2-8°C (36-46°F).

Caution: Cytotoxic Agent

Manufactured for:

Baxter Healthcare Corporation

Deerfield, IL 60015 USA

Product of Germany

Single-Dose Vial NDC 0338-0722-01
Rx only

Pemetrexed
for Injection, USP

500 mg per vial

For intravenous use only.
Discard unused portion.

Baxter

Caution: Cytotoxic Agent

Each vial contains pemetrexed disodium equivalent to 500 mg pemetrexed, 500 mg of mannitol. Hydrochloric acid and/or sodium hydroxide may have been added to adjust pH.

Storage: Store powder at 25°C (77°F); excursions permitted to 15-30°C (59-86°F) [see USP Controlled Room Temperature]. See accompanying literature for storage of reconstituted and infusion solutions.

Manufactured for: Baxter Healthcare Corporation, Deerfield, IL 60015 USA. Product of Germany.

LOT/EXP:
USA 184-05-01-088 C 57

LOT/EXP:

Single-Dose Vial

Rx only

NDC 0338-0722-01

Pemetrexed

for Injection, USP

500 mg per vial

For intravenous use only.

Discard unused portion.

Baxter Logo

Caution: Cytotoxic Agent

Each vial contains pemetrexed disodium equivalent

to 500 mg pemetrexed, 500 mg of mannitol.
Hydrochloric acid and/or sodium hydroxide may
have been added to adjust pH.


Storage: Store powder at 25°C (77°F);
excursions permitted to 15-30°C (59-86°F) [see USP Controlled
Room Temperature]. See accompanying literature
for storage of reconstituted and infusion solutions.
Manufactured for: **Baxter Healthcare Corporation,**
Deerfield, IL 60015 USA. Product of Germany .

Barcode

(01)00303380722013

2D Barcode

USA HA-65-01-838 C 57



Pemetrexed
for Injection, USP
500 mg per vial

Barcode
(01) 20303380722017

Baxter is a registered trademark of
Baxter International Inc.

GTIN:
SN:
LOT:
EXP:

Single-Dose Vial
NDC 0338-0722-01

Pemetrexed
for Injection, USP

500 mg per vial

Barcode

HA-80-02-963
USA

Pemetrexed
for Injection, USP

Each vial contains Pemetrexed disodium equivalent to 500 mg pemetrexed, 500 mg of mannitol. Hydrochloric acid and/or sodium hydroxide may have been added to adjust pH.

To Reconstitute:Add 20 mL of 0.9% Sodium Chloride Injection (preservative free) to make a solution containing 25 mg/mL pemetrexed. Reconstituted solution must be further diluted before use (see accompanying literature).

Administer infusion solution within 24 hours after initial reconstitution. Discard unused portion.

See accompanying literature for dosage.

C
477

Single-Dose Vial
NDC 0338-0722-01

Pemetrexed
for Injection, USP

500 mg per vial

For intravenous use only.

Rx only

Baxter Logo

2335B0115

Barcode

Pemetrexed
for Injection, USP

Storage:Store powder at 25°C (77°F); excursions permitted to 15-30°C (59-86°F) [see USP Controlled Room Temperature].

Store reconstituted and infusion solutions at refrigerated, 2-8°C (36-46°F).

Caution: Cytotoxic Agent

Manufactured for:
Baxter Healthcare Corporation
Deerfield, IL 60015 USA

Product of Germany

PEMETREXED

pemetrexed injection, powder, lyophilized, for solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:83703-561
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PEMETREXED DISODIUM HEPTAHYDRATE (UNII: 9T47E4OM16) (PEMETREXED - UNII:04Q9AIZ7NO)	PEMETREXED	100 mg in 10 mL

Inactive Ingredients

Ingredient Name	Strength
MANNITOL (UNII: 3OWL53L36A)	106 mg in 10 mL
HYDROCHLORIC ACID (UNII: QTT17582CB)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83703-561-01	1 in 1 CARTON	08/18/2022	
1		10 mL in 1 VIAL, SINGLE-DOSE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA214436	08/18/2022	

PEMETREXED

pemetrexed injection, powder, lyophilized, for solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:83703-562
Route of Administration	INTRAVENTRICULAR		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PEMETREXED DISODIUM HEPTAHYDRATE (UNII: 9T47E4OM16) (PEMETREXED - UNII:04Q9AIZ7NO)	PEMETREXED	500 mg in 50 mL

Inactive Ingredients

Ingredient Name	Strength
MANNITOL (UNII: 3OWL53L36A)	500 mg in 50 mL
HYDROCHLORIC ACID (UNII: QTT17582CB)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83703-562-01	1 in 1 CARTON	08/18/2022	
1		50 mL in 1 VIAL, SINGLE-DOSE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA214436	08/18/2022	

Labeler - Bamboo US Bidco LCC (119087615)

Establishment

Name	Address	ID/FEI	Business Operations
Baxter Oncology GmbH		344276063	analysis(83703-561, 83703-562) , manufacture(83703-561, 83703-562) , label(83703-561, 83703-562) , pack(83703-561, 83703-562) , sterilize(83703-561, 83703-562)

Revised: 7/2024

Bamboo US Bidco LCC