

BETR COLD AND FLU NIGHTTIME RELIEF- acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hydrochloride tablet, film coated

Live Betr LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Live Betr LLC Cold & Flu Nighttime Relief Drug Facts

Active ingredients (in each caplet)

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Doxylamine succinate 6.25 mg

Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Nasal decongestant

Uses

- temporarily relieves common cold/flu symptoms:
- nasal congestion
- sinus congestion and pressure
- cough due to minor throat and bronchial irritation
- cough to help you sleep
- minor aches and pains
- headache
- fever
- sore throat
- runny nose and sneezing
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if

you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- **do not use more than directed**

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed – see overdose warning
- do not exceed 8 caplets per 24 hrs

adults & children 12 yrs & over	2 caplets with water every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Inactive ingredients

crospovidone, D&C yellow #10 aluminum lake, FD&C blue #1 aluminum lake, FD&C blue #2 aluminum lake, FD&C yellow #6 aluminum lake, flavor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, propylene glycol, silicon dioxide, stearic acid, sucralose, talc, titanium dioxide

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

Betr™

Get Betr, Give Back.

cold & flu nighttime relief

ACETAMINOPHEN, PHENYLEPHRINE HCl, DOXYLAMINE SUCCINATE,
DEXTROMETHORPHAN HBr, CAPLETS

Pain Reliever | Fever Reducer | Antihistamine

Nasal Decongestant | Cough Suppressant

Actual Size

-Relief Of: Minor Aches & Pains; Nasal Congestion & Sinus Pressure; Sneezing; Runny
Nose; Cough

24 CAPLETS

Your purchase helped someone in need.



Questions or comments? 1-800-719-9260

Inactive ingredients (continued)
 FD&C Blue #1 aluminum lake, FD&C Yellow #6 aluminum lake, FD&C Blue #2 aluminum lake, FD&C Yellow #5 aluminum lake, FD&C Yellow #10 aluminum lake, FD&C Yellow #12 aluminum lake, FD&C Yellow #13 aluminum lake, FD&C Yellow #15 aluminum lake, FD&C Yellow #17 aluminum lake, FD&C Yellow #18 aluminum lake, FD&C Yellow #19 aluminum lake, FD&C Yellow #2 aluminum lake, FD&C Yellow #21 aluminum lake, FD&C Yellow #22 aluminum lake, FD&C Yellow #23 aluminum lake, FD&C Yellow #24 aluminum lake, FD&C Yellow #25 aluminum lake, FD&C Yellow #26 aluminum lake, FD&C Yellow #27 aluminum lake, FD&C Yellow #28 aluminum lake, FD&C Yellow #29 aluminum lake, FD&C Yellow #3 aluminum lake, FD&C Yellow #30 aluminum lake, FD&C Yellow #31 aluminum lake, FD&C Yellow #32 aluminum lake, FD&C Yellow #33 aluminum lake, FD&C Yellow #34 aluminum lake, FD&C Yellow #35 aluminum lake, FD&C Yellow #36 aluminum lake, FD&C Yellow #37 aluminum lake, FD&C Yellow #38 aluminum lake, FD&C Yellow #39 aluminum lake, FD&C Yellow #40 aluminum lake, FD&C Yellow #41 aluminum lake, FD&C Yellow #42 aluminum lake, FD&C Yellow #43 aluminum lake, FD&C Yellow #44 aluminum lake, FD&C Yellow #45 aluminum lake, FD&C Yellow #46 aluminum lake, FD&C Yellow #47 aluminum lake, FD&C Yellow #48 aluminum lake, FD&C Yellow #49 aluminum lake, FD&C Yellow #50 aluminum lake, FD&C Yellow #51 aluminum lake, FD&C Yellow #52 aluminum lake, FD&C Yellow #53 aluminum lake, FD&C Yellow #54 aluminum lake, FD&C Yellow #55 aluminum lake, FD&C Yellow #56 aluminum lake, FD&C Yellow #57 aluminum lake, FD&C Yellow #58 aluminum lake, FD&C Yellow #59 aluminum lake, FD&C Yellow #60 aluminum lake, FD&C Yellow #61 aluminum lake, FD&C Yellow #62 aluminum lake, FD&C Yellow #63 aluminum lake, FD&C Yellow #64 aluminum lake, FD&C Yellow #65 aluminum lake, FD&C Yellow #66 aluminum lake, FD&C Yellow #67 aluminum lake, FD&C Yellow #68 aluminum lake, FD&C Yellow #69 aluminum lake, FD&C Yellow #70 aluminum lake, FD&C Yellow #71 aluminum lake, FD&C Yellow #72 aluminum lake, FD&C Yellow #73 aluminum lake, FD&C Yellow #74 aluminum lake, FD&C Yellow #75 aluminum lake, FD&C Yellow #76 aluminum lake, FD&C Yellow #77 aluminum lake, FD&C Yellow #78 aluminum lake, FD&C Yellow #79 aluminum lake, FD&C Yellow #80 aluminum lake, FD&C Yellow #81 aluminum lake, FD&C Yellow #82 aluminum lake, FD&C Yellow #83 aluminum lake, FD&C Yellow #84 aluminum lake, FD&C Yellow #85 aluminum lake, FD&C Yellow #86 aluminum lake, FD&C Yellow #87 aluminum lake, FD&C Yellow #88 aluminum lake, FD&C Yellow #89 aluminum lake, FD&C Yellow #90 aluminum lake, FD&C Yellow #91 aluminum lake, FD&C Yellow #92 aluminum lake, FD&C Yellow #93 aluminum lake, FD&C Yellow #94 aluminum lake, FD&C Yellow #95 aluminum lake, FD&C Yellow #96 aluminum lake, FD&C Yellow #97 aluminum lake, FD&C Yellow #98 aluminum lake, FD&C Yellow #99 aluminum lake, FD&C Yellow #100 aluminum lake.

Directions (continued)

Take only as directed—see overdose warning.
 Do not exceed 8 caplets per 24 hrs.
 Add to children 12 yrs & over: 2 caplets with water every 4 hrs.
 Children 4 to under 12 yrs: ask a doctor.
 Children under 4 yrs: do not use.

Drug Facts

Purpose
 Pain reliever/fever reducer
 Cough suppressant
 Antihistamine
 Nasal decongestant
 Cough suppressant

Active ingredients (in each caplet)
 Acetaminophen HBr 10 mg
 Doxylamine succinate 6.25 mg
 Phenylephrine HCl 5 mg

Uses
 Temporarily relieves common cold/flu symptoms:
 nasal congestion
 sinus congestion
 and pressure
 cough due to minor throat and bronchial irritation
 cough to help you sleep
 minor aches and pains
 fever
 sore throat
 runny nose and sneezing
 reduces swelling of nasal passages
 temporarily restores free breathing through the nose
 promotes nasal and/or sinus drainage

Warnings
 This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours. With other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product. Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
 skin reddening
 blisters
 rash
 If a skin reaction occurs, stop use and seek medical help right away.
 Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. If you have ever had an allergic reaction to this product or any of its ingredients:
 Ask a doctor before use if you have:
 liver disease
 heart disease
 high blood pressure
 thyroid disease
 diabetes
 glaucoma
 cough that occurs with too much phlegm (mucus)
 a breathing problem such as emphysema or chronic bronchitis
 persistent or chronic cough such as occurs with smoking, asthma, or emphysema
 trouble urinating due to an enlarged prostate gland
 Ask a doctor or pharmacist before use if you are:
 taking sedatives or tranquilizers
 taking the blood thinning drug warfarin
 When using this product:
 do not use more than directed
 excitability may occur, especially in children
 marked drowsiness may occur
 avoid alcoholic drinks
 be careful when driving a motor vehicle or operating machinery
 alcohol, sedatives, and tranquilizers may increase drowsiness
 Stop use and ask a doctor if:
 you get nervous, dizzy or sleepless
 pain, nasal congestion, or cough gets worse or lasts more than 7 days
 fever gets worse or lasts more than 3 days
 redness or swelling is present
 new symptoms occur
 cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
 If pregnant or breast-feeding, ask a health professional before use.
 Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

COOLIE ARIEVA

72V62 CD C3



DO NOT USE IF BLISTER UNIT IS BROKEN OR TORN

Distributed by
 Betr Remedies, LLC
 730 Arizona Ave
 Santa Monica, CA 90401
 www.betrremedies.com



STORE AT 20-25°C (68-77°F).

PARENTS:
 Learn about teen medicine abuse
www.StopMedicineAbuse.org

Betr

Get Betr, Give Back.




cold & flu
 nighttime relief

ACETAMINOPHEN, PHENYLEPHRINE HCl, DOXYLAMINE
 SUCCINATE, DEXTROMETHORPHAN HBr, CAPLETS

Pain Reliever	Fever Reducer	Antihistamine
Nasal Decongestant	Cough Suppressant	

Relief Of: Minor Aches & Pains; Fever; Nasal Congestion
 & Sinus Pressure; Sneezing, Runny Nose; Cough

Actual Size



24 CAPLETS

BETR COLD AND FLU NIGHTTIME RELIEF

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hydrochloride tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:80267-721
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
DOXYLAMINE SUCCINATE (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
CROSPVIDONE (15 MPA.S AT 5%) (UNII: 68401960MK)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	GREEN	Score	no score
Shape	OVAL	Size	19mm
Flavor		Imprint Code	L72V
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:80267-721-62	12 in 1 CARTON	10/19/2021	
1		2 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	10/19/2021	

Labeler - Live Betr LLC (117423281)

Revised: 9/2022

Live Betr LLC