

ASPIRIN 81MG- aspirin 81mg tablet, chewable
Lodaa LLC

316 - Chewable Cherry Low Dose Aspirin 81mg Tablet

Active ingredient(s)

Aspirin 81 mg (NSAID*)
*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Use(s)

for the temporary relief of minor aches and pains or as recommended by your doctor

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include hives, facial swelling, asthma (wheezing), shock.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Do not use

if you are allergic to aspirin or any other pain reliever/fever reducer this product for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic you have asthma

Ask a doctor or pharmacist before use if

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away
- you experience any of the following signs of stomach bleeding: feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or loss of hearing occurs

If pregnant or breastfeeding,

ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

Keep out of reach of children

In case of accidental overdose, get medical help or contact a Poison Control Center(1-800-222-1222) immediately.

Directions

- drink a full glass of water with each dose
- **adults and children 12 years and over:** take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- **children under 12 years:** consult a doctor

Other information

store at room temperature.

do not use if imprinted safety seal under cap is broken or missing

Inactive ingredients

corn starch, dextrans, FD&C red # 40 aluminum lake, flavor, saccharin sodium

Questions/Comments

Call 1-888-952-0050 (Mon-Fri 9AM-5PM EST)

Principal Display Panel

AARDVARK PHARMA

Compare To Bayer® Chewable Low Dose Aspirin active Ingredient**

Adult Chewable Low Dose

Aspirin

Pain Reliever (NSAID) 81 mg

CHEWABLE CHERRY FLAVORED

Talk to your doctor or other healthcare provider before using this product for your heart.

36 CHEWABLE Tablets



Drug Facts

Active ingredient (in each tablet)
Aspirin 81 mg (NSAID*)Pain reliever *nonsteroidal anti-inflammatory drug

Purpose
for the temporary relief of minor aches and pains or as recommended by your doctor

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Aspirin may cause a severe allergic reaction which may include hives, facial swelling, asthma (wheezing), shock.

LB2185 R0224

Drug facts continued on inside panel

Distributed by: Aardvark Pharma LLC 1415 W 22nd St, Tower Floor Oak Brook, IL USA



7 26872 06267 0

Lot No:
Exp Date

PRINT AREA

Drug Facts (continued)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer or this product for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor or if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- gout
- diabetes
- arthritis

Drug Facts (continued)

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away
- you experience any of the following signs of stomach bleeding: feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or loss of hearing occurs

if pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

Other information

- store at room temperature
- do not use if imprinted safety seal under cap is broken or missing

Inactive ingredients
corn starch, dextrans, FD&C red # 40 aluminum lake, flavor, saccharin sodium

Questions or comments?
1-888-952-0050 (Mon-Fri 9AM-5PM EST)

**This product is not manufactured or distributed by Bayer HealthCare LLC., owner of the registered trademark Bayer® Chewable Low Dose Aspirin.

Aspirin 81mg Cherry Flavor

ASPIRIN 81MG

aspirin 81mg tablet, chewable

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:83139-316

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
DEXTRATES (UNII: G263MI44RU)	
FD&C RED NO. 40 ALUMINUM LAKE (UNII: 6T47AS764T)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	

Product Characteristics

Color	pink	Score	no score
Shape	ROUND	Size	7mm
Flavor	CHERRY	Imprint Code	AZ013
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83139-316-36	36 in 1 BOTTLE; Type 0: Not a Combination Product	06/04/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	06/04/2024	

Labeler - Lodaat LLC (011845714)

Revised: 6/2024

Lodaat LLC