

**MUCUS RELIEF NIGHTTIME CONGESTION COUGH MAXIMUM STRENGTH-
acetaminophen diphenhydramine hci phenylephrine hci liquid
EQUATE (Wal-Mart Stores, Inc.) (see also WAL-MART INC)**

Drug Facts

Active ingredients (in each 20 mL)

Acetaminophen 650 mg

Diphenhydramine HCl 25 mg

Phenylephrine HCl 10 mg

Purposes

Pain reliever

Antihistamine/cough suppressant

Nasal decongestant

Uses

- temporarily relieves
 - nasal congestion
 - headache
 - minor aches and pain
 - cough
 - sinus congestion and pressure
 - runny nose and sneezing
- temporarily promotes nasal and/or sinus drainage
- controls cough to help you get to sleep

Warnings

Liver warning: this product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen ask a doctor or pharmacist.
- with other drug containing diphenhydramine, even one used on the skin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- for children under 12 years of age

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- diabetes
- thyroid disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

ask a doctor or pharmacist before use if

- you are taking the blood thinning drug warfarin
- you are taking sedatives or tranquilizers

When using this product

- **do not use more than directed**
- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedative, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness or sleeplessness occur
- pain, nasal congestion, or cough gets worse, or lasts more than 7 days
- fever gets worse, or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms

Directions

- **do not take more than directed (see Overdose warning)**
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL=milliliter
- dose as follows or as directed by a doctor
- adults and children 12 years and older: 20 mL every 4 hours while symptoms last
- children under 12 years of age: do not use

Other information

- **each 20 mL contains:** sodium 12 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Questions or comments?

1-888-287-1915

Principal Display Panel

Compare to Maximum Strength Mucinex® Sinus-Max® Night time Congestion & Cough active ingredients*

NIGHTTIME

Sinus Relief Congestion & Cough

Acetaminophen - Pain Reliever

Diphenhydramine HCL - Antihistamine/Cough Suppressant

Phenylephrine HCL - Nasal Decongestant

MAXIMUM STRENGTH

Multi-Symptom Relief

- Clear sinus congestion

- Relieves headache
- relieves runny nose & sneezing
- Control Cough

For ages 12+

FL OZ (mL)

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.

*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® Sinus-Max® Night Time Congestion & Cough.

DISTRIBUTED BY: Walmart Inc.,

Bentonville, AR 72716

Package Label

NDC 49035-461-06

equate™



NIGHTTIME

Sinus Relief Congestion & Cough

Acetaminophen - Pain reliever
Diphenhydramine HCl - Antihistamine/Cough suppressant
Phenylephrine HCl - Nasal decongestant

MAXIMUM STRENGTH

Multi-Symptom Relief

- Clears sinus congestion
- Relieves headache
- Relieves runny nose & sneezing
- Controls cough

6 FL OZ (177mL)

For Ages 12+

P.L.D.-B378C LB006396

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.

Satisfaction guaranteed - Or we'll replace it or give you your money back. For questions or comments please call **1-888-287-1915**.

DISTRIBUTED BY: Walmart Inc., Bentonville, AR 72716
*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® Sinus-Max® Night Time Congestion & Cough.



PLD-B378C LB006396

PEEL CORNER FOR DRUG FACTS

Drug Facts	
Active ingredients (in each 20 mL)	Purposes
Acetaminophen 650 mg.....	Pain reliever
Diphenhydramine HCl 25 mg.....
.....	Antihistamine/cough suppressant
Phenylephrine HCl 10 mg.....	Nasal decongestant

Uses	
<ul style="list-style-type: none"> temporarily relieves nasal congestion minor aches and pains sinus congestion and pressure runny nose and sneezing temporarily promotes nasal and/or sinus drainage controls cough to help you get to sleep 	<ul style="list-style-type: none"> headache cough

Warnings	
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:	
<ul style="list-style-type: none"> more than 4,000 mg of acetaminophen in 24 hours with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product 	
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:	
<ul style="list-style-type: none"> skin reddening blisters rash 	

Drug Facts (continued)

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other drug containing diphenhydramine, even one used on the skin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- for children under 12 years of age

Ask a doctor before use if you have

liver disease	heart disease
high blood pressure	diabetes
thyroid disease	glaucoma

- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Drug Facts (continued)

Ask a doctor or pharmacist before use if

- you are taking the blood thinning drug warfarin
- you are taking sedatives or tranquilizers

When using this product

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse, or lasts more than 7 days
- fever gets worse, or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or

Drug Facts (continued)

contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- mL = milliliter
- keep dosing cup with product
- dose as follows or as directed by a doctor
- adults and children 12 years and older: 20 mL every 4 hours while symptoms last
- children under 12 years of age: do not use

Other information

- each 20 mL contains: sodium 12 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Questions or comments?
1-888-287-1915

PEEL CORNER FOR MORE DRUG FACTS

EQUATE Nighttime Sinus Relief Congestion & Cough

MUCUS RELIEF NIGHTTIME CONGESTION COUGH MAXIMUM STRENGTH

acetaminophen diphenhydramine hci phenylephrine hci liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:49035-461
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 20 mL
DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	25 mg in 20 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 20 mL

Inactive Ingredients	
Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYL GALLATE (UNII: 8D45NN7V92)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49035-461-06	177 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	04/30/2018	04/30/2025

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/30/2018	04/30/2025

Labeler - EQUATE (Wal-Mart Stores, Inc.) (see also WAL-MART INC) (051957769)

Revised: 12/2023

EQUATE (Wal-Mart Stores, Inc.) (see also WAL-MART INC)