

**MOTION SICKNESS- meclizine hcl tablet**  
**ARMY AND AIR FORCE EXCHANGE SERVICE**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Exchange Select 44-403**

***Active ingredient (in each tablet)***

Meclizine HCl 25 mg

***Purpose***

Antiemetic

***Uses***

for the prevention and treatment of nausea, vomiting, or dizziness associated with motion sickness

***Warnings***

**Do not use**

for children under 12 years of age unless directed by a doctor

**Ask a doctor before use if you have**

- glaucoma
- difficulty in urination due to enlargement of the prostate gland
- a breathing problem such as emphysema or chronic bronchitis

**Ask a doctor or pharmacist before use if you are**

taking sedatives or tranquilizers.

**When using this product**

- drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives, and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

***Directions***

- to prevent motion sickness, take the first dose one hour before starting activity
- to prevent or treat motion sickness: 1 to 2 tablets once daily for adults and children 12 years and over, or as directed by a doctor

**Other information**

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- protect from heat and humidity
- see end flap for expiration date and lot number

**Inactive ingredients**

corn starch, D&C yellow #10 aluminum lake, lactose, magnesium stearate, silica gel

**Questions of comments?**

**1-800-426-9391**

**Principal Display Panel**

exchange √ select™

**Compare to The Active Ingredient of Dramamine® Less Drowsy Formula\***

LESS DROWSY FORMULA

**MOTION SICKNESS**

Meclizine HCl Tablets 25mg / Antiemetic

**Helps prevent Nausea and Dizziness due to Motion Sickness for up to 24 hours**

**16**

**TABLETS**

Actual Size

quality value

\*This product is not manufactured or distributed by Prestige Brands, Inc., owner of the registered trademark Dramamine® Less Drowsy Formula.

50844 REV0715J40321

**"SATISFACTION GUARANTEED OR YOUR MONEY BACK"**


Manufactured For Your Military Exchanges

Distributed by: LNK International, Inc.

Hauppauge, NY 11788

1-800-426-9391

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING**



# MOTION SICKNESS

Helps prevent Nausea and Dizziness due to Motion Sickness for up to 24 hours

exchange select™

Compare To The Active Ingredient of Dramamine® Less Drowsy Formula\*

## LESS DROWSY FORMULA MOTION SICKNESS

Meclizine HCl Tablets 25mg / Antiemetic

Helps prevent Nausea and Dizziness due to Motion Sickness for up to 24 hours

16 Actual Size  
Tablets

quality  
value

LESS DROWSY FORMULA  
MOTION SICKNESS

exchange  
select™



TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

OMIT D

exchange select™

## LESS DROWSY FORMULA MOTION SICKNESS

B-0006E-403-21-R  
REV0715J40321

6

50844 REV0715J40321  
Less Drowsy Formula.  
This product is not manufactured or distributed by Prestige Brands, Inc., owner of the registered trademark Dramamine®

Manufactured For Your Military Exchanges  
Distributed by: LNK International, Inc.  
Hauptaage, NY 11788  
1-800-426-9391

“SATISFACTION GUARANTEED OR YOUR MONEY BACK”  
**Questions or comments?** 1-800-426-9391  
**Inactive ingredients** com starch, D&C yellow #10 aluminum lake, lactose, magnesium stearate, silica gel  
■ see end flap for expiration date and lot number  
■ protect from heat and humidity  
■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)  
■ **OPENED OR BLISTER IS TORN OR BROKEN**  
■ **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS**

**Other information**  
■ **difficultly in urination** due to enlargement of the prostate gland  
■ a breathing problem such as emphysema or chronic bronchitis  
■ **Ask a doctor or pharmacist before use if you are taking** sedatives or tranquilizers.  
■ **When using this product** ■ avoid alcoholic beverages ■ drowsiness may occur ■ drowsiness may increase alcohol, sedatives, and tranquilizers may increase drowsiness  
■ use caution when driving a motor vehicle or operating machinery  
■ **if pregnant or breast-feeding, ask a health professional** before use.

**Warnings**  
Do not use for children under 12 years of age unless directed by a doctor.  
■ **Ask a doctor before use if you have** ■ glaucoma  
■ **Uses** for the prevention and treatment of nausea, vomiting, or dizziness associated with motion sickness.  
■ **Directions**  
■ to prevent motion sickness, take the first dose one hour before starting activity  
■ to prevent or treat motion sickness: 1 to 2 tablets once daily for adults and children 12 years and over, or as directed by a doctor  
■ **Drug Facts (continued)**  
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Drug Facts**  
KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

**Active ingredient (in each tablet)** Purpose  
Meclizine HCl 25 mg, Antiemetic

**Directions**  
■ to prevent motion sickness, take the first dose one hour before starting activity  
■ to prevent or treat motion sickness: 1 to 2 tablets once daily for adults and children 12 years and over, or as directed by a doctor  
■ **Other information**  
■ **difficultly in urination** due to enlargement of the prostate gland  
■ a breathing problem such as emphysema or chronic bronchitis  
■ **Ask a doctor or pharmacist before use if you are taking** sedatives or tranquilizers.  
■ **When using this product** ■ avoid alcoholic beverages ■ drowsiness may occur ■ drowsiness may increase alcohol, sedatives, and tranquilizers may increase drowsiness  
■ use caution when driving a motor vehicle or operating machinery  
■ **if pregnant or breast-feeding, ask a health professional** before use.

LEBG149F A1494A

Exchange Select 44-403

**MOTION SICKNESS**

meclizine hcl tablet

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:55301-403
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
MECLIZINE HYDROCHLORIDE (UNII: HDP7W44CIO) (MECLIZINE - UNII:3L5TQ84570)	MECLIZINE HYDROCHLORIDE	25 mg

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
STARCH, CORN (UNII: O8232NY3SJ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
LACTOSE (UNII: J2B2A4N98G)	

**Product Characteristics**

<b>Color</b>	YELLOW	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	9mm
<b>Flavor</b>		<b>Imprint Code</b>	44;403
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55301-403-21	2 in 1 CARTON	06/24/2002	
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		

**Marketing Information**

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC MONOGRAPH FINAL	part336	06/24/2002	

**Labeler** - ARMY AND AIR FORCE EXCHANGE SERVICE (001695568)

<b>Establishment</b>			
<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
LNK International, Inc.		038154464	PACK(55301-403)

<b>Establishment</b>			
<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
LNK International, Inc.		832867894	MANUFACTURE(55301-403)

Revised: 2/2017

ARMY AND AIR FORCE EXCHANGE SERVICE