

**KOPERTOX- copper naphthenate liquid**  
**Zoetis**

-----  
**KOPERTOX®**

(copper naphthenate)

**WATER-RESISTANT PROTECTION WITHOUT BANDAGING**

**RECOMMENDED** as an aid in treating horses and ponies with thrush due to organisms susceptible to copper naphthenate.

**Do not use in horses intended for human consumption.**

**CAUTION: COMBUSTIBLE MIXTURE**

**ACTIVE INGREDIENT:**

Copper Naphthenate ..... 37.5% w/w

**INACTIVE INGREDIENTS:**

..... 62.5% w/w  
Total .....100.0%

**FOR EXTERNAL USE ONLY**

**Not for Use in Humans.**

**Keep Out of Reach of Children.**

**Indicated in the treatment of thrush in horses and ponies**

**CAUTION:** Do not use in horses intended for human consumption.

**CAUTION: COMBUSTIBLE MIXTURE**

Use in a well-ventilated place. Avoid fire, flame, sparks or heaters.

If swallowed, do not induce vomiting, call physician immediately. Avoid breathing vapor. Avoid contact with skin and eyes.

**Keep out of reach of children and pets.**

**GENERAL DIRECTIONS:**

Clean the hoof thoroughly, removing debris and necrotic material prior to application of KOPERTOX®. Apply daily to affected hoofs with a narrow paint brush (about 1") until fully healed. Caution: Do not allow runoff of excess KOPERTOX® onto hair since contact with KOPERTOX® may cause some hair loss. Do not contaminate feed.

Store at controlled room temperature 15° to 30°C (59° to 86°F). Keep container tightly closed when not in use.

**NOTE:** KOPERTOX® is easily removed from hands, clothing and surfaces with light grade fuel oil or any type of lighter fluid.

**CONTAINS FOIL SEAL - REMOVE BEFORE USE.**

**SHAKE WELL BEFORE USE.**

**CONTACT INFORMATION:**

To report suspected adverse drug events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS) contact Zoetis Inc. at 1-888-963-8471. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>



236 mL (8 oz) Rev. 03/2023 5164100

473 mL (16 oz) Rev. 03/2023 5161100

Approved by FDA under ANADA # 200-304



Distributed by:  
Zoetis Inc.  
Kalamazoo, MI 49007

Manufactured by:  
First Priority, Inc.  
Elgin, IL 60123

**236 mL (8 oz)**

# KOPERTOX<sup>®</sup>

## (copper naphthenate)

WATER-RESISTANT PROTECTION  
WITHOUT BANDAGING

**RECOMMENDED**  
as an aid in treating horses and  
ponies with thrush due to organisms  
susceptible to copper naphthenate.

**Do not use in horses intended  
for human consumption.**

**CAUTION: COMBUSTIBLE MIXTURE**

**ACTIVE INGREDIENT:**

Copper Naphthenate .....37.5% w/w

**INACTIVE INGREDIENTS:** .....62.5% w/w

Total.....100.0%

**FOR EXTERNAL USE ONLY**

**Not for Use in Humans. Keep Out of Reach of Children.**

**236 mL (8 oz)**

Approved by FDA under ANADA # 200-304

Manufactured by:  
First Priority, Inc.  
Elgin, IL 60123

**zoetis**  
Distributed by:  
Zoetis Inc.  
Kalamazoo, MI 49007

# KOPERTOX<sup>®</sup>

## (copper naphthenate)

**Indicated in the treatment of  
thrush in horses and ponies**

**CAUTION:** Do not use in horses intended for human consumption.

**CAUTION: COMBUSTIBLE MIXTURE**

Use in a well-ventilated place. Avoid fire, flame, sparks or heaters.  
If swallowed, do not induce vomiting, call physician immediately.  
Avoid breathing vapor. Avoid contact with skin and eyes.

**Keep out of reach of children and pets.**

**GENERAL DIRECTIONS:** Clean the hoof thoroughly, removing  
debris and necrotic material prior to application of KOPERTOX<sup>®</sup>.  
Apply daily to affected hoofs with a narrow paint brush (about 1")  
until fully healed. Caution: Do not allow runoff of excess  
KOPERTOX<sup>®</sup> onto hair since contact with KOPERTOX<sup>®</sup> may  
cause some hair loss. Do not contaminate feed.

Store at controlled room temperature 15° to 30°C (59° to 86°F).  
Keep container tightly closed when not in use.

**NOTE:** KOPERTOX<sup>®</sup> is easily removed from hands, clothing and  
surfaces with light grade fuel oil or any type of lighter fluid.

**CONTAINS FOIL SEAL – REMOVE BEFORE USE**

**SHAKE WELL BEFORE USE**

**CONTACT INFORMATION:** To report suspected adverse  
drug events, for technical assistance or to obtain a copy of the  
Safety Data Sheet (SDS) contact Zoetis Inc. at 1-888-963-8471.  
For additional information about adverse drug experience  
reporting for animal drugs, contact FDA at 1-888-FDA-VETS or  
<http://www.fda.gov/reportanimalae>

Revised: March 2023

51641000



Lot No.:

Exp. Date:

**NO  
VARNISH**



**473 mL (16 oz)**

# KOPERTOX<sup>®</sup>

## (copper naphthenate)

WATER-RESISTANT PROTECTION  
WITHOUT BANDAGING

**RECOMMENDED**  
as an aid in treating horses and  
ponies with thrush due to organisms  
susceptible to copper naphthenate.

**Do not use in horses intended  
for human consumption.**

### CAUTION: COMBUSTIBLE MIXTURE

#### ACTIVE INGREDIENT:

Copper Naphthenate ..... 37.5% w/w

INACTIVE INGREDIENTS: ..... 62.5% w/w

Total.....100.0%

### FOR EXTERNAL USE ONLY

**Not for Use in Humans.  
Keep Out of Reach of Children.**

**473 mL (16 oz)**

Approved by FDA under ANADA # 200-304

Manufactured by:  
First Priority, Inc.  
Elgin, IL 60123

**zoetis**  
Distributed by:  
Zoetis Inc.  
Kalamazoo, MI 49007

# KOPERTOX<sup>®</sup>

## (copper naphthenate)

Indicated in the treatment of  
thrush in horses and ponies

**CAUTION:** Do not use in horses intended for human  
consumption.

### CAUTION: COMBUSTIBLE MIXTURE

Use in a well-ventilated place. Avoid fire, flame, sparks or  
heaters.

If swallowed, do not induce vomiting, call physician  
immediately. Avoid breathing vapor. Avoid contact with skin  
and eyes.

**Keep out of reach of children and pets.**

**GENERAL DIRECTIONS:** Clean the hoof thoroughly,  
removing debris and necrotic material prior to application of  
KOPERTOX<sup>®</sup>. Apply daily to affected hoofs with a narrow  
paint brush (about 1") until fully healed. Caution: Do not allow  
runoff of excess KOPERTOX<sup>®</sup> onto hair since contact with  
KOPERTOX<sup>®</sup> may cause some hair loss. Do not contaminate  
feed.

Store at controlled room temperature 15° to 30°C (59° to 86°F).  
Keep container tightly closed when not in use.

**NOTE:** KOPERTOX<sup>®</sup> is easily removed from hands, clothing  
and surfaces with light grade fuel oil or any type of lighter fluid.

**CONTAINS FOIL SEAL – REMOVE BEFORE USE**

**SHAKE WELL BEFORE USE**

**CONTACT INFORMATION:** To report suspected adverse  
drug events, for technical assistance or to obtain a copy  
of the Safety Data Sheet (SDS) contact Zoetis Inc. at  
1-888-963-8471. For additional information about adverse  
drug experience reporting for animal drugs, contact FDA at  
1-888-FDA-VETS or <http://www.fda.gov/reportanimalae>

Revised: March 2023

51641100



Lot No.:

Exp. Date:

**NO  
VARNISH**



## KOPERTOX

copper naphthenate liquid

### Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:54771-1100
Route of Administration	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
COPPER NAPHTHENATE (UNII: 9J2IBN2H70) (COPPER - UNII: 789U1901C5)	COPPER NAPHTHENATE	0.825 g in 1 mL

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:54771-1100-1	12 in 1 CASE		
1		236 mL in 1 BOTTLE, DISPENSING		
2	NDC:54771-1100-2	12 in 1 CASE		
2		473 mL in 1 BOTTLE, DISPENSING		
<b>Marketing Information</b>				
<b>Marketing Category</b>		<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
ANADA		ANADA200304	10/12/2023	

**Labeler -** Zoetis (828851555)

**Establishment**

Name	Address	ID/FEI	Business Operations
FIRST PRIORITY INCORPORATED		179925722	manufacture

Revised: 12/2023

Zoetis