

**MENSTRUAL PAIN RELIEF COMPLETE- acetaminophen, pyrilamine maleate,
caffeine tablet, film coated
Shield Pharmaceuticals Corp**

Menstrual Pain Relief Complete

Drug Facts

Active ingredient (in each liquid-filled capsule)

Acetaminophen 500 mg

Caffeine 60 mg

Pyrilamine maleate 15 mg

Purpose

Pain reliever

Diuretic

Antihistamine

Uses

for the temporary relief of these symptoms associated with menstrual periods:

- cramps
- bloating
- water
- weight gain
- headache
- backache
- muscle aches
- fatigue

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash
- hives
- facial swelling
- asthma (wheezing)

- shock

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- if you have ever had an allergic reaction to this product or any of its ingredients
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if you have

- difficulty in urination due to enlargement of the prostate gland
- liver disease _ glaucoma
- a breathing problem such as emphysema or chronic

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- avoid alcoholic beverages
- excitability may occur, especially in children
- be careful when driving a motor vehicle or operating machiner
- alcohol, sedatives, and tranquilizers may increase drowsiness
- you may get drowsy
- limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat. The recommended dose of this product contains about as much caffeine as a cup of coffee.

Stop use and ask a doctor if

- new symptoms occur _ redness or swelling is present
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

adults and children 12 years and over:

- **do not take more than the recommended dose**
- adults and children 12 years and over: take 2 caplets with water
- repeat every 6 hours, as needed
- do not exceed 6 caplets per day

- children under 12 years: ask a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Inactive ingredients

aerosil, calcium phosphate, colloidal silicon dioxide, cross carmelloose sodium, FD&C yellow #6, hypromellose, isopropyl alcohol, magnesium stearate, methlyene chloride, microcrystalline cellulose, pregelatinized starch, polyethylene glycols, poly vinyl pyrrolidone, povidone, stearic acid, talc, titanium dioxide

Questions or comments?

1-800-373-6981 (toll-free)

Distributed by:

Shield Pharmaceuticals Corp.

Hauppauge, NY 11788

PRINCIPAL DISPLAY PANEL

Menstrual Pain Relief Complete

Acetaminophen/Caffeine/Pyrilamine maleate

Pain reliever/Diuretic/Antihistamine

Carton



Label

Drug Facts (continued)
product because too much caffeine may cause nervousness, irritability, sleeplessness, and a racing heart beat. The recommended dose of this product contains about as much caffeine as a cup of coffee.

Stop use and ask a doctor if:

- redness or swelling is present ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days

The seal design is a serious condition.
If pregnant or breast feeding, ask a health professional before use.

Keep out of reach of children

In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

do not take more than the recommended dose

- adults and children 12 years and older: take 2 caplets with water
- repeat every 6 hours as needed
- do not exceed 6 caplets per day
- children under 12 years: ask a doctor

Other Information

■ store at 25°C (77°F), exclusions permitted between 15°–30°C (59°–86°F)
inactive ingredients
See caution for complete list of inactive ingredients.

Questions or comments?

1-800-373-8881 (toll-free)



DO NOT USE IF SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Purpose

Pain reliever

Diacetaminophen

Antihistamine

Active ingredients (in each caplet)

Acetaminophen 500 mg

Caffeine 60 mg

Pyrilamine maleate 15 mg

Uses

for the temporary relief of these symptoms associated with menstrual periods: ■ cramps ■ bloating ■ water-weight gain
■ headache ■ backache ■ muscle aches

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more acetaminophen products every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash ■ hives
■ facial swelling ■ asthma (wheezing) ■ shock
If a skin reaction occurs, stop use and seek medical help right away.

Do not use: ■ if you have ever had an allergic reaction to this product or any of its ingredients ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
Ask a doctor before use if you have: ■ difficulty in urination due to enlargement of the prostate gland ■ liver disease ■ glaucoma ■ a breathing problem such as emphysema or chronic bronchitis
Ask a doctor or pharmacist before use if you are: ■ taking the blood thinner dicumarol ■ taking sedatives or tranquilizers
When using this product: ■ avoid alcohol, beverages ■ eat fatty foods, especially in children ■ be careful when driving a motor vehicle or operating machinery ■ avoid sedatives and tranquilizers which may increase drowsiness ■ you may get drowsy ■ limit the use of caffeine-containing medications, foods, or beverages while taking this product

Drug Facts (continued under label)

Distributed by:
Shield Pharmaceuticals Corp.
Hauppauge, NY 11788
LOT: SLD24002
EXP.: 04/2025



ORG. 0124
Made in India

MENSTRUAL PAIN RELIEF COMPLETE

acetaminophen, pyrilamine maleate, caffeine tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:83059-0006
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg
CAFFEINE (UNII: 3G6A5W38E) (CAFFEINE - UNII:3G6A5W38E)	CAFFEINE	60 mg
PYRILAMINE MALEATE (UNII: R35D29L3ZA) (PYRILAMINE - UNII:HPE317O9TL)	PYRILAMINE MALEATE	15 mg

Inactive Ingredients

Ingredient Name	Strength

SILICON DIOXIDE (UNII: ETJ7Z6XBU4)
CALCIUM PHOSPHATE (UNII: 97Z1W3NDX)
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)
HYPROMELLOSES (UNII: 3NXW29V3WO)
ISOPROPYL ALCOHOL (UNII: ND2M416302)
MAGNESIUM STEARATE (UNII: 70097M6I30)
METHYLENE CHLORIDE (UNII: 588X2YUY0A)
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)
STARCH, CORN (UNII: O8232NY3SJ)
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)
PYRROLIDONE (UNII: KKL5D39EOL)
POVIDONE (UNII: FZ989GH94E)
STEARIC ACID (UNII: 4ELV7Z65AP)
TALC (UNII: 7SEV7J4R1U)
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)

Product Characteristics

Color	white	Score	no score
Shape	CAPSULE	Size	17mm
Flavor		Imprint Code	S17;M7
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83059-0006-1	1 in 1 CARTON	01/16/2024	
1		15 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	01/16/2024	

Labeler - Shield Pharmaceuticals Corp (118724924)

Revised: 1/2024

Shield Pharmaceuticals Corp