# CHILDRENS IBUPROFEN- ibuprofen tablet, chewable CVS Pharmacy

-----

CVS Pharmacy, Inc. Children's Ibuprofen Drug Facts

# Active ingredient (in each chewable tablet)

Ibuprofen 100 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

# **Purposes**

Pain reliever/fever reducer

### Uses

temporarily:

- reduces fever
- relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

# Warnings

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Sore throat warning:** Severe or persistent sore throat or sore throat accompanied by high fever,

headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless

### Do not use

- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

## Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic

# Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

## When using this product

- mouth or throat burning may occur; give with food or water
- take with food or milk if stomach upset occurs

## Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding:
- feels faint
- vomits blood
- has bloody or black stools
- has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
- chest pain
- trouble breathing
- weakness in one part or side of body
- slurred speech
- leg swelling
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

# Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

## **Directions**

- · chew or crush tablets completely before swallowing
- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- find right dose on chart below. If possible, use weight to dose; otherwise use age.
- if needed, repeat dose every **6-8 hours**
- do not use more than 4 times a day

Dosing Chart			
Weight (lb)	Age (yr)	Tablets	
under 24	under 2	ask a doctor	
24-35	2-3	1	
36-47	4-5	1 ½	
48-59	6-8	2	
60-71	9-10	2 ½	
72-95	11	3	

### Other information

- phenylketonurics: contains phenylalanine 6 mg per tablet
- store between  $20-25^{\circ}$ C (68-77°F)
- do not use if printed seal under cap is broken or missing

# **Inactive ingredients**

acesulfame potassium, ammonium glycyrrhizin, aspartame, carnauba wax, croscarmellose sodium, D&C red no. 27 aluminum lake, FD&C blue no. 1 aluminum lake, hypromellose, magnesium stearate, mannitol, natural and artificial flavors, silicon dioxide, sodium lauryl sulfate, soybean oil, succinic acid

# Questions or comments?

1-800-719-9260

# Package/Label Principal Display Panel

Compare to the active ingredient in Children's Motrin®

Chewables

For ages 2-11 / Children's

Ibuprofen CHEWABLE TABLETS, 100 mg

LASTS UP TO 8 HOURS

Pain reliever; Fever reducer (NSAID)

Chew or crush tablets completely before swallowing
Grape Flavor
Actual Size
24 CHEWABLE TABLETS







### KEEP CARTON FOR REFERENCE

### Drug Facts

Active ingredient (in each chewable tablet) Purposes buprofen 100 mg (NSAID)\*..... \*nonsteroidal anti-inflammatory drug

### Uses temporarily: ■ reduces fever

■ relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

Warnings Allergy alert: buprofen may cause a severe allergic reaction,

Allergy alert buprolen may cause a severe allergic reaction, especially in people allergic baptin. Symptoms may include:

This may be allergic baptin. Symptoms may include:

This may be allergic may be allergic may be allergic reaction occurs stop use and seek medical help right away. Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:
This had a broach users or bleeding problems

Tables a blood thinghout (affice and and or stemid drug.)

In has had stomach ubers or bleeding problems
It takes a blood thinning (anticoagulant) or steroid drug
Takes other drugs containing pressription romoprescription
NSADs (aspirin, ibuprofen, naproxen, or others)
Takes more or for a longer time from directed
Heart attackand stroke warming: NSADs, except aspirin, increase
the risk of heart stack, heart failure, and stroke. These can be fatal. The
risk is higher if you use more than directed or for longer than directed
than than the manning Sovere or existent serve than at or sere threat. sax is inject injective in the customers of no foliar installations. Somethinast warning. Severe or persistent one throat or some throat accompanied by high fever headsche, nausea, and vomitting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless diecked by doctor.

### Drug Facts (continued)

### Do not use

■ if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer

■ right before or after heart surgery

### Ask a doctor before use if

ask a outour eeure use ii ■ stomach bleeding warning applies to your child ■ child has a history of stomach problems, such as heartburn ■ child has problems or serious side effects from taking pain relievers or fever reducers
■ child has not been drinking fluids
■ child has lost a lot of fluid due to vomiting or diarrhea

■ child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke

■ child has asthma ■ child is taking a diuretic

Ask a doctor or pharmacist before use if the child is under a doctor's care for any serious condition

■ taking any other drug

When using this product ■ mouthor throat burning may occur, give with food or water ■ take with food or milk if stomach upset occurs

Stopuse and ask adoctor if

■ Child experiences any of the following signs of stomach bleeding:
■ feels faint ■ vomits blood ■ has bloody or black stools
■ has stomach pain fractioes not get better

child has symptoms of heart problems or stroke:

chest pain trouble breathing

weakness in one part or side of body
surredspeech legswelling

### Drug Facts (continued)

- the child does not get any relief within first day (24 hours) of treatment fever or pain gets worse or lasts more than 3 days redness or swelling is present in the painful area

- any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-900-222-1222)

chewor crush tablets completely before swallowing
this product does not contain directions or complete warnings

for adult use
do not give more than directed
find right dose on chart below. If possible, use weight to dose; otherwise useage.

If needed, repeat dose every 6-8 hours
donotuse more than 4 times a day

### Docina Chart

Weight (lb)	Age (yr)	Tablets
under 24	under 2	ask a doctor
24-35	2-3	1
36-47	4-5	11/2
48-59	6-8	2
60-71	9-10	21/2
72-95	11	3

Other Into m atton

phenyket ourts: contain phenylalarine 6 mg pertédiet

sche between 3-0.5°C (186.7°P F)

do not u.e. if printed seal und e cap istreken or missing

Irac f vel Ingredients acesultane potas ium, ammonium
obysyrition aspateme, careatheway, rocsamebbes solum,
0.8% redno.27 duminum bke, 10.8°C three.1 duminum late,
hypromelbes, magnas ium sebaste, mannitu hate' at and artibid
flavor salicondioi des cdium buyl stitite, a yrben oʻl, succinicadd Questors ar comments ?1-800-719-9260

Drug Facts (portinued)

# **CHILDRENS IBUPROFEN**

ibuprofen tablet, chewable

### **Product Information**

Product Type HUMAN OTC DRUG Item Code (Source) NDC:69842-521

Route of Administration ORAL

# **Active Ingredient/Active Moiety**

	Ingr	edient Name	Basis of Strength	Strength
IBUPRO FEN	(UNII: WK2XYI10QM) (IBU	PROFEN - UNII:WK2XYI10QM)	IBUPROFEN	100 mg

Inactive Ingredients

Ingredient Name

Strength

ACESULFAME POTASSIUM (UNII: 230 V73Q5G9)

AMMO NIUM GLYCYRRHIZATE (UNII: 3VRD35U26C)

ASPARTAME (UNII: Z0H242BBR1)

CARNAUBA WAX (UNII: R12CBM0EIZ)

CROSCARMELLOSE SODIUM (UNII: M28 O L 1 HH48)

**D&C RED NO. 27** (UNII: 2LRS 185U6K)

FD&C BLUE NO. 1 (UNII: H3R47K3TBD)

HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)

MAGNESIUM STEARATE (UNII: 70097M6I30)

MANNITOL (UNII: 30WL53L36A)

 $\textbf{SILICON DIO XIDE} \ (\text{UNII: ETJ7Z6 XBU4})$ 

SODIUM LAURYL SULFATE (UNII: 368GB5141J)

SOYBEAN OIL (UNII: 241ATL177A)

SUCCINIC ACID (UNII: AB6MNQ6J6L)

## **Product Characteristics**

Color	PURPLE (Lavender)	Score	2 pieces
Shape	ROUND	Size	12mm
Flavor	GRAPE	Imprint Code	L521
Contains			

Ρ	ac	kag	ing
_	-		~~~~

#	Item Code	Package Description	<b>Marketing Start Date</b>	Marketing End Date
1	NDC:69842-521-62	1 in 1 CARTON	05/29/2019	
1		24 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:69842-521-67	2 in 1 CARTON	05/29/2019	
2		24 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076359	05/29/2019	

# Labeler - CVS Pharmacy (062312574)

Revised: 6/2020 CVS Pharmacy