

**NAPROXEN SODIUM- naproxen sodium tablet, film coated  
Preferred Pharmaceuticals Inc.**

-----  
**Perrigo Naproxen Sodium Tablets 220 mg Drug Facts**

**Active ingredient (in each tablet)**

Naproxen sodium 220 mg  
(naproxen 200 mg) (NSAID)\*

\*nonsteroidal anti-inflammatory drug

**Purposes**

Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis
- muscular aches
- backache
- menstrual cramps
- headache
- toothache
- the common cold
- temporarily reduces fever

**Warnings**

**Allergy alert:** Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

### **Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

### **Ask a doctor before use if**

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

### **Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

### **When using this product**

- take with food or milk if stomach upset occurs

### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- pain gets worse or lasts more than 10 days

- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- **do not take more than directed**
- **the smallest effective dose should be used**
- drink a full glass of water with each dose

Adults and children 12 years and older	<ul style="list-style-type: none"> <li>• take 1 tablet every 8 to 12 hours while symptoms last</li> <li>• for the first dose you may take 2 tablets within the first hour</li> <li>• do not exceed 2 tablets in any 8- to 12-hour period</li> <li>• do not exceed 3 tablets in a 24-hour period</li> </ul>
Children under 12 years	<ul style="list-style-type: none"> <li>• ask a doctor</li> </ul>

**Other information**

- **each tablet contains:** sodium 22 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

**Inactive ingredients**

FD&C blue no. 2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

**Questions or comments?**

**1-800-719-9260**

# Principal Display Panel

Compare to Aleve® active ingredient

Naproxen Sodium Tablets 220 mg

Pain Reliever/Fever Reducer (NSAID)

Actual Size

Strength To Last 12 Hours

Repackaged By: Preferred Pharmaceuticals Inc.


**Naproxen Sodium Tab USP 220mg**

Generic for Aleve


Each tablet contains: Naproxen sodium 220mg g...pain reliever / (naproxen 200mg) (NSAID).. .fever reducer

**Pkg Size:** Exp Date:  
Lot#: Lot#: Batch#: Ins:  
Mfg: Padagis Pharmaceuticals Ltd  
Prod#: Prod#: Warning

Allergy alert: naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Stomach bleeding warning: this product contains an NSAID, which may cause severe stomach bleeding. Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer or right before or after heart surgery. Ask a doctor or pharmacist before use if you are under a doctor's care for any serious condition, or taking any other drug. Keep out of the reach of children. If pregnant or breast-feeding, ask a health professional before use. Store at 20°-25°C (68°-77°F). Avoid high humidity and excessive heat above 40°C (104°F). Tablet is round, blue, and imprinted with L490



CAUTION: Federal law PROHIBITS transfer of this drug to any person other than the patient for whom it was prescribed



Naproxen Sodium Tab USP 220mg  
Qty: Ins:  
Lot#: Bat#:  
Prod# (NDC):

Naproxen Sodium Tab USP 220mg  
Qty: Ins:  
Lot#: Bat#:  
Prod# (NDC):

Naproxen Sodium Tab USP 220mg  
Qty:  
Insurance NDC:  
Lot#: Bat#:

Naproxen Sodium Tab USP 220mg  
Qty: Ins:  
Lot#: Bat#:  
Prod# (NDC):

Directions English

Take \_\_\_ Tablet(s) \_\_\_ time(s) a day

Instrucciones Espanol:

Cada \_\_\_ Tableta(s) \_\_\_ vece(s) al dia

Log  
Chart  
Billing  
Patient

<b>NAPROXEN SODIUM</b> naproxen sodium tablet, film coated			
Product Information			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:68788-8447(NDC:45802-490)
<b>Route of Administration</b>	ORAL		
Active Ingredient/Active Moiety			
<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>	
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ)	NAPROXEN SODIUM	220 mg	
Inactive Ingredients			
<b>Ingredient Name</b>	<b>Strength</b>		
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)			
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)			
MAGNESIUM STEARATE (UNII: 70097M6I30)			
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)			
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)			
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)			

**TALC** (UNII: 7SEV7J4R1U)

**TITANIUM DIOXIDE** (UNII: 15FIX9V2JP)

### Product Characteristics

<b>Color</b>	BLUE (Light Blue)	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	L490
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68788-8447-2	20 in 1 BOTTLE; Type 0: Not a Combination Product	05/17/2023	
2	NDC:68788-8447-3	30 in 1 BOTTLE; Type 0: Not a Combination Product	05/17/2023	
3	NDC:68788-8447-4	40 in 1 BOTTLE; Type 0: Not a Combination Product	05/17/2023	
4	NDC:68788-8447-6	60 in 1 BOTTLE; Type 0: Not a Combination Product	05/17/2023	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074661	05/17/2023	

**Labeler** - Preferred Pharmaceuticals Inc. (791119022)

**Registrant** - Preferred Pharmaceuticals Inc. (791119022)

### Establishment

Name	Address	ID/FEI	Business Operations
Preferred Pharmaceuticals Inc.		791119022	REPACK(68788-8447)

Revised: 5/2023

Preferred Pharmaceuticals Inc.